

Tackling delirium: a crucial target for improving clinical outcomes

ABSTRACT

Delirium is a common and debilitating syndrome in hospitalized patients, and its impact on mortality, morbidity and duration of hospital admission is increasingly apparent. Delirium is a complex phenomenon, for which there is no specific treatment, but research over the last decade has revealed contributing factors, many of which are modifiable, and preventative strategies have demonstrated benefit. This review highlights the importance of reducing the impact of delirium on hospitalized patients, and summarizes the current evidence for strategies to achieve this. Current recommendations focus on the pre-emptive implementation of multi-modal non-pharmacological interventions to reduce the occurrence of delirium in the first place, and routine monitoring using validated tools to identify delirious patients early. Potential therapies for established delirium remain controversial.

Delirium is a common affliction, its presence detected in approximately one fifth of general hospital inpatients (Ryan et al, 2013), and its incidence increasing to 30% in the elderly (Potter et al, 2006) and 80% of the critically unwell (Ely et al, 2001). Perhaps because of its ubiquity, it is easy to dismiss as merely an unavoidable consequence of illness, but delirium is independently associated with increased mortality, as well as longer stays in hospital, and on the intensive care unit (Ely et al, 2004). Each additional day of delirium is associated with an average of 10 extra days in hospital and a 10% increased risk of death, as well as an association with long-term cognitive impairment (Jones, 2007). Magnetic resonance imaging has shown an association between the duration of delirium and cerebral atrophy and white matter disruption (Reade and Finfer, 2014).

Various theories have been proposed to describe the pathophysiology underlying delirium, including cerebral inflammation altering the blood–brain barrier and changes in neurotransmitter levels, such as reduced cholinergic activity or increased catecholamine serotonergic activity (Maclullich et al, 2013).

Dr Hanieh Asadi, Clinical Fellow, Intensive Care Unit, Royal Free Hospital, London

Dr Daniel Martin, Consultant in Anaesthesia and Intensive Care Medicine, Intensive Care Unit, Royal Free Hospital, London and Reader, Division of Surgery and Interventional Science, University College London, London

Dr Helen McKenna, Research Fellow, Division of Surgery and Interventional Science, University College London, Royal Free Hospital, London NW3 2QG

Correspondence to: Dr H McKenna (helen.mckenna.15@ucl.ac.uk)

More recently, it has been proposed that the features of delirium may be a clinical manifestation of circadian rhythm disruption (Madrid-Navarro et al, 2015). Both illness and the hospital environment itself disrupt the usual 24-hour cycles that govern human behaviour (such as sleeping, waking, eating and physical activity), physiological indices (temperature, heart rate, urine output) and intracellular processes (gene expression, enzyme activity and mitochondrial function) (McKenna et al, 2017). The circadian oscillation in these diverse functions is synchronized to the time of day through the light–dark cycle and other cues such as regular feeding times. One of the early signs of delirium is inversion of the sleep–wake cycle, and this has led to the hypothesis that delirium may represent the observable signs of underlying circadian dysrhythmia, the impact of which extends much deeper, affecting a multitude of cellular and organ system functions. Prevention, identification and treatment of delirium are key priorities for improving clinical outcomes for all patients admitted to hospital. This article reviews the current evidence for strategies to reduce the impact of delirium on hospital patients.

Diagnosing delirium

According to the Diagnostic and Statistical Manual of Mental Disorders (DSM-5) delirium is a disturbance in attention (reduced ability to focus, sustain and shift attention), awareness and cognition over a short period of time (hours to a few days) and fluctuates in severity during the course of the day (Pisani et al, 2006). Delirium can manifest in three different ways:

1. Hyperactive (agitated)
2. Hypoactive (withdrawn)
3. Mixed (alternating periods of hyperactive and hypoactive).

The most common type is mixed followed by hypoactive delirium (Page and Gough, 2010). The form of delirium most likely to go unrecognized is hypoactive delirium, and this is associated with the worst prognosis (Robinson et al, 2011). The presence of delirium is consistently underdiagnosed, with detection rates by clinical staff shown to be as low as 25% (Voyer et al, 2012). Diagnosis requires regular and repeated screening as, without it, the majority of delirium cases go unnoticed (Ryan et al, 2013). The National Institute of Health and Care Excellence (2010) guideline recommends that all patients are observed at least daily for recent changes or fluctuations of usual behaviour, and if present, a formal assessment must be carried out to confirm the diagnosis. All patients on an intensive care unit

should be formally assessed for delirium at least once every 12 hours using a validated screening tool (Hayhurst et al, 2016). The Confusion Assessment Method (CAM) is the most widely used in clinical practice (De and Wand, 2015), and has been adapted for the intensive care unit (CAM-ICU) (Hayhurst et al, 2016). Other screening tools were reviewed by De and Wand (2015), including the Delirium Rating Scale (found to be most useful for psychiatric and geriatric patients) and the Nurses' Delirium Screening Checklist (best for recovery room setting).

Primary prevention of delirium

The most effective strategy for reducing the impact of delirium is to prevent its occurrence in the first place. Delirium is precipitated by multiple factors, all of which must be addressed. Risk factors for delirium are divided into four domains (Van Rompaey et al, 2009):

1. Patient characteristics
2. Chronic pathology
3. Acute illness
4. The environment.

The first two – patient characteristics (such as age, alcohol use, smoking) and chronic pathology (such as pre-existing cognitive impairment) – are important for identifying those most at risk, for prioritization in any delirium prevention strategy. The latter two – acute illness and the environment – may be amenable to modification and offer a target for reducing delirium and improving clinical outcomes.

Multicomponent strategies which minimize these risk factors have consistently demonstrated efficacy in reducing incidence of delirium. A systematic review and meta-analysis including 14 interventional studies concluded that multi-modal non-pharmacological interventions can be effective in preventing delirium and falls during a hospital admission in the elderly, with 11 studies showing decreases in incidence of delirium (odds ratio 0.47, 95% confidence intervals 0.42–0.76), and rate of falls reduced by 64% in two randomized or matched trials (Hshieh et al, 2015). These bundles typically include measures to ensure adequate oxygenation, hydration, nutrition and analgesia. They prompt staff to facilitate mobilization, as well as provide necessary aides for vision and hearing.

Modifications to the environment may help in reducing circadian disruption, by promoting daytime cues such as adequately bright light to simulate the light intensity of outdoors and supporting physical activity and social interaction in the day. Relatives should be actively encouraged to visit and bring cues such as photographs and familiar music. Patients should be reminded verbally of the time, date and location. Provision of easily visible clocks and calendars aid in active reorientation.

At night, reducing sleep interference by minimizing bright light, loud noises and avoidable clinical interventions will promote sleep. In a cohort study in a mixed intensive care unit, the implementation of a bundle of environmental interventions to promote sleep led to an increase in qualitative as well as quantitative measures of sleep, which

“ The most effective strategy for reducing the impact of delirium is to prevent its occurrence in the first place. ”


in turn led to a reduction in the incidence and duration of delirium from 33% to 14% (Patel et al, 2014). The bundle consisted of reducing the monitor alarms, telephone ringtones, providing eye masks, dimming the lights during night-time, minimizing non-clinical discussions around patients' bed spaces, orientating the patients to time, place and date, documenting regular CAM-ICU assessments, addressing pain, assessing patients for sedation hold and setting sedation targets. *Figure 1* summarizes the key components of an evidence-based delirium prevention strategy.

As yet, there is insufficient evidence of benefit from any prophylactic pharmacological strategies. Only two trials have demonstrated reductions in postoperative incidence of delirium with prophylactic administration of antipsychotics, such as intravenous haloperidol (0.5 mg bolus followed by 0.1 mg/hr infusion for 12 hours) (Wang et al, 2012), and oral risperidone (0.5 mg twice daily) (Hakim et al, 2012) but trial results have been inconsistent. Administration of the exogenous form of the circadian hormone melatonin (0.5 mg once nightly) was associated with a lower incidence of delirium (12%) in elderly medical patients compared to placebo (31%) (Al-Aama et al, 2011) but again, results from other studies have been conflicting (Oh et al, 2017).

Treatment options for established delirium

Despite the success of non-pharmacological strategies for the prevention of delirium, very few studies have assessed their efficacy in the treatment of established delirium. The first priority is to identify and treat any modifiable organic causes such as pain, hypoxia, acidemia and other metabolic derangements, infection and haemodynamic instability (Oh et al, 2017). Other organic causes, non-clinical seizures, hypertensive encephalopathy and

Figure 1. Components of a multi-faceted delirium prevention strategy.



KEY POINTS

- Delirium is associated with a higher mortality, morbidity and longer stays in hospital, with negative impact on quality of life long after discharge.
- Delirium is grossly underdiagnosed, and must be screened for daily in hospital patients.
- Multicomponent, non-pharmacological interventions have proven effective in decreasing the occurrence of delirium.
- Evidence for the benefit of pharmacological approaches to delirium prevention or treatment remain limited.

intracerebral haemorrhage should be ruled out on the basis of history, examination and investigation when appropriate. It is important to review all drug side effects and drug interactions, or the impact of withdrawal of drugs, particularly in the context of pre-existing alcohol and/or substance abuse. The following classes of drugs have been implicated in contributing to delirium: opioids, antidepressants, anticonvulsants, antihistamines, antipsychotics, antimuscarinics and steroids. Less familiar but also implicated are assorted agents such as furosemide, ranitidine, atenolol and digoxin (Hayhurst et al, 2016). Although often a necessary evil in the intensive care unit, sedative medications and deeper levels of sedation are associated with increased risk of delirium, with benzodiazepines appearing to be the worst culprits (Nguyen and Pon, 2016).

Pharmacological approaches to treating delirium remain controversial. Antipsychotics are most commonly used in patients with hyperactive delirium where there is a risk to safety, but the side effects of the proposed drugs must be balanced against any benefit they may provide. In delirium as a result of alcohol or benzodiazepine withdrawal, benzodiazepines offer a specific treatment, but otherwise it is advised to avoid these drugs as they can contribute to development of delirium. Although antipsychotics, such as haloperidol, are included in guidelines as potential therapy for established delirium, a systematic review of the use of antipsychotics (including haloperidol, risperidone and olanzapine) for the treatment of delirium in elderly patients demonstrated no improvement in delirium duration, severity, mortality or hospital length of stay in the 19 identified studies (Neufeld et al, 2016). In one randomized controlled trial of 247 patients in the palliative care setting, those taking haloperidol or risperidone had greater delirium scores compared to those receiving placebo, and mortality rates were higher in those receiving haloperidol (Agar et al, 2017).

In the intensive care setting, dexmedetomidine, a selective alpha₂-adrenoceptor agonist with analgesic and sedative effects, has shown promise in the treatment of hyperactive delirium. In a randomized controlled trial of 74 mechanically ventilated patients, those receiving dexmedetomidine had a shorter duration of delirium and increased number of ventilator-free hours at day 7, compared to those receiving placebo (Reade et al, 2016).

It is possible that this effect was a result of the decrease in the amount of other sedative agents (propofol, opioids and midazolam) required by the treatment group compared to placebo. This finding has been corroborated by the significantly lower prevalence of delirium in mechanically ventilated patients receiving dexmedetomidine for sedation rather than midazolam (54% compared to 77%) in a multicentre international prospective trial of 375 patients (Riker et al, 2009). Patients in the dexmedetomidine group also had more delirium-free days, and were extubated sooner than the midazolam group.

Conclusions

Illness and admission to hospital place a patient at risk of developing delirium, which has stark consequences for the individual, reducing his/her likelihood of survival, and diminishing the quality of life long after hospital discharge. Once established, delirium is difficult to treat, but there is convincing evidence from clinical trials for the effectiveness of comprehensive bundles of care in preventing its onset in the first place. Detection and prevention of delirium should be a key priority for all health-care professionals. The authors recommend auditing incidence of delirium, on the wards and in the intensive care unit, and for multidisciplinary clinical teams to take shared responsibility in the implementation of strategies to improve delirium detection and prevention. **BJHM**

Conflict of interest: none.

- Agar MR, Lawlor PG, Quinn S et al (2017) Efficacy of oral risperidone, haloperidol, or placebo for symptoms of delirium among patients in palliative care: a randomized clinical trial. *JAMA Intern Med* **177**(1): 34–42. <https://doi.org/10.1001/jamainternmed.2016.7491>
- Al-Aama T, Brymer C, Gutmanis I, Woolmore-Goodwin SM, Esbaugh J, Dasgupta M (2011) Melatonin decreases delirium in elderly patients: a randomized, placebo-controlled trial. *Int J Geriatr Psychiatry* **26**(7): 687–694. <https://doi.org/10.1002/gps.2582>
- De J, Wand AP (2015) Delirium screening: a systematic review of delirium screening tools in hospitalized patients. *Gerontologist* **55**(6): 1079–1099. <https://doi.org/10.1093/geront/gnv100>
- Ely EW, Inouye SK, Bernard GR et al (2001) Delirium in mechanically ventilated patients: validity and reliability of the confusion assessment method for the intensive care unit (CAM-ICU). *JAMA* **286**(21): 2703–2710. <https://doi.org/10.1001/jama.286.21.2703>
- Ely EW, Shintani A, Truman B et al (2004) Delirium as a predictor of mortality in mechanically ventilated patients in the intensive care unit. *JAMA* **291**(14): 1753–1762. <https://doi.org/10.1001/jama.291.14.1753>
- Hakim SM, Othman AI, Naoum DO (2012) Early treatment with risperidone for subsyndromal delirium after on-pump cardiac surgery in the elderly: a randomized trial. *Anesthesiology* **116**(5): 987–997. <https://doi.org/10.1097/ALN.0b013e31825153cc>
- Hayhurst CJ, Pandharipande PP, Hughes CG (2016) Intensive care unit delirium: a review of diagnosis, prevention, and treatment. *Anesthesiology* **125**(6): 1229–1241. <https://doi.org/10.1097/ALN.0000000000001378>
- Hsieh TT, Yue J, Oh E, Puella M, Dowal S, Travison T, Inouye SK (2015) Effectiveness of multicomponent nonpharmacological delirium interventions: a meta-analysis. *JAMA Intern Med* **175**(4): 512–520. <https://doi.org/10.1001/jamainternmed.2014.7779>
- Jones C (2007) Aftermath of intensive care: the scale of the problem. *Br J Hosp Med (Lond)* **68**(9): 464–466. <https://doi.org/10.12968/hmed.2007.68.9.27163>
- Maclullich AM, Anand A, Davis DH et al (2013) New horizons in the

- pathogenesis, assessment and management of delirium. *Age Ageing* 42(6): 667–674. <https://doi.org/10.1093/ageing/aft148>
- Madrid-Navarro CJ, Sanchez-Galvez R, Martinez-Nicolas A, Marina R, Garcia JA, Madrid JA, Rol MA (2015) Disruption of circadian rhythms and delirium, sleep impairment and sepsis in critically ill patients. Potential therapeutic implications for increased light-dark contrast and melatonin therapy in an ICU environment. *Curr Pharm Des* 21(24): 3453–3468. <https://doi.org/10.2174/1381612821666150706105602>
- McKenna HT, Reiss IK, Martin DS (2017) The significance of circadian rhythms and dysrhythmias in critical illness. *J Intensive Care Soc* 18(2): 121–129. <https://doi.org/10.1177/11751143717692603>
- National Institute for Health and Care Excellence (2010) Delirium: prevention, diagnosis and management. Clinical guideline CG103. www.nice.org.uk/guidance/cg103 (accessed 25 January 2018)
- Neufeld KJ, Yue J, Robinson TN, Inouye SK, Needham DM (2016) Antipsychotic medication for prevention and treatment of delirium in hospitalized adults: a systematic review and meta-analysis. *J Am Geriatr Soc* 64(4): 705–714. <https://doi.org/10.1111/jgs.14076>
- Nguyen HM, Pon D (2016) Off-label use of dexmedetomidine for the treatment of delirium in the intensive care unit. *P T* 41(10): 642–643.
- Oh ES, Fong TG, Hshieh TT, Inouye SK (2017) Delirium in older persons: advances in diagnosis and treatment. *JAMA* 318(12): 1161–1174. <https://doi.org/10.1001/jama.2017.12067>
- Page V, Gough K (2010) Management of delirium in the intensive care unit. *Br J Hosp Med (Lond)* 71(7): 372–376. <https://doi.org/10.12968/hmed.2010.71.7.48994>
- Patel J, Baldwin J, Bunting P, Laha S (2014) The effect of a multicomponent multidisciplinary bundle of interventions on sleep and delirium in medical and surgical intensive care patients. *Anaesthesia* 69(6): 540–549. <https://doi.org/10.1111/anae.12638>
- Pisani MA, Araujo KL, Van Ness PH, Zhang Y, Ely EW, Inouye SK (2006) A research algorithm to improve detection of delirium in the intensive care unit. *Crit Care* 10(4): R121. DOI:10.1186/cc5027
- Potter J, George J, Guideline Development Group (2006) The prevention, diagnosis and management of delirium in older people: concise guidelines. *Clin Med (Lond)* 6(3): 303–308. <https://doi.org/10.7861/clinmedicine.6-3-303>
- Reade MC, Finfer S (2014) Sedation and delirium in the intensive care unit. *N Engl J Med* 370(5): 444–454. <https://doi.org/10.1056/NEJMra1208705>
- Reade MC, Eastwood GM, Bellomo R et al; DahlIA Investigators; Australian and New Zealand Intensive Care Society Clinical Trials Group (2016) Effect of dexmedetomidine added to standard care on ventilator-free time in patients with agitated delirium: a randomized clinical trial. *JAMA* 315(14): 1460–1468. <https://doi.org/10.1001/jama.2016.2707>
- Riker RR, Shehabi Y, Bokesch PM et al; SEDCOM (Safety and Efficacy of Dexmedetomidine Compared With Midazolam) Study Group (2009) Dexmedetomidine vs midazolam for sedation of critically ill patients: a randomized trial. *JAMA* 301(5): 489–499. <https://doi.org/10.1001/jama.2009.56>
- Robinson TN, Raeburn CD, Tran ZV, Brenner LA, Moss M (2011) Motor subtypes of postoperative delirium in older adults. *Arch Surg* 146(3): 295–300. <https://doi.org/10.1001/archsurg.2011.14>
- Ryan DJ, O'Regan NA, Caoimh RÓ et al (2013) Delirium in an adult acute hospital population: predictors, prevalence and detection. *BMJ Open* 3(1). pii: e001772. <https://doi.org/10.1136/bmjopen-2012-001772>
- Van Rompaey B, Elseviers MM, Schuurmans MJ, Shorridge-Baggett LM, Truijzen S, Bossaert L (2009) Risk factors for delirium in intensive care patients: a prospective cohort study. *Crit Care* 13(3): R77. <https://doi.org/10.1186/cc7892>
- Voyer P, Richard S, McCusker J et al (2012) Detection of delirium and its symptoms by nurses working in a long term care facility. *J Am Med Dir Assoc* 13(3): 264–271. <https://doi.org/10.1016/j.jamda.2010.11.002>
- Wang W, Li HL, Wang DX et al (2012) Haloperidol prophylaxis decreases delirium incidence in elderly patients after noncardiac surgery: a randomized controlled trial*. *Crit Care Med* 40(3): 731–739. <https://doi.org/10.1097/CCM.0b013e3182376e4f>

Organised by

BRITISH JOURNAL OF
HOSPITAL MEDICINE BRITISH JOURNAL OF
NEUROSCIENCE NURSING**20th ANNIVERSARY**

Parkinson's 2018

etc Victoria, 1 Drummond Gate, London
Monday 21st May 2018

To book your place:

☎ Call us on +44(0)20 7501 6796

🌐 www.mahealthcareevents.co.uk/parkinsons201825% EARLY BIRD RATE
UNTIL 9TH MARCH 2018