

# Osteosarcopenia

**O**steoporosis and sarcopenia are two chronic musculoskeletal conditions that can have devastating impacts both on individuals and wider society. With an ageing population, both conditions are likely to become increasingly prevalent in future, increasing the incidence of fragility fractures, and leading to greater morbidity, mortality and socioeconomic costs (Edwards et al, 2015).

Around one in two women and one in five men over 50 years old will sustain an osteoporotic fracture throughout their remaining lifetime, with more than 8.9 million osteoporotic fractures sustained annually worldwide (Hernlund et al, 2013). In 2010, the direct cost of osteoporotic fractures in EU countries was estimated at a staggering €35.4 billion, with hip fractures representing 54% of the cost. In the UK alone, around 79 000 hip fractures occurred in 2010 (Hernlund et al, 2013), with up to one in three patients dying within 1 year (Klop et al, 2014).

Osteoporosis, characterized by low bone mass and micro-architectural deterioration of bone tissue (Edwards et al, 2015), and sarcopenia, the loss of muscle mass, strength and function (Fuggle et al, 2017), are increasingly recognized as a 'hazardous duet' in the pathogenesis of fragility fractures, with the sarcopenic propensity for falls acting synergistically with the osteoporotic vulnerability of bones to increase fracture risk (Crepaldi and Maggi, 2005). As such, sarco-osteopenia, now more commonly known as osteosarcopenia, has been proposed as a new term to represent a frail subset of the elderly population with concomitant osteoporosis and sarcopenia (Binkley and Buehring, 2009). These patients are at significantly higher risk of falls, fractures and institutionalization (Huo et al, 2015), and have a significantly higher mortality rate than osteoporotic or sarcopenic patients alone (Yoo et al, 2018).

An increasing body of evidence demonstrates considerable overlap in the pathophysiology of osteoporosis and sarcopenia, bringing with it exciting possibilities to treat the conditions together. Bone and muscle not only interact mechanically, known as the 'mechanostat hypothesis' (Frost, 2003), but also communicate with each other biochemically via complex paracrine and endocrine mechanisms in order to maintain muscle and bone homeostasis (Girgis et al, 2014). This has facilitated development of new therapeutic agents which exploit this bone-muscle crosstalk (Girgis et al, 2014), which will be discussed later in this review.

With our ageing population and the enormous socioeconomic impact of fragility fractures, it is imperative that clinicians are aware of osteosarcopenia and the ways

## ABSTRACT

Osteosarcopenia is a newly described syndrome that describes the co-existence of osteoporosis and sarcopenia, two chronic musculoskeletal conditions associated with ageing. Osteoporosis, a condition of low bone mass and micro-architectural deterioration of bone, and sarcopenia, the loss of muscle mass, strength and function, often co-exist in a frail subset of the elderly population, leading to significantly worsened outcomes than seen in either condition alone. These include a greater risk of falls, fractures and institutionalization, and significant socioeconomic costs. With our ageing population, osteosarcopenia is a public health concern that will become increasingly relevant in the future. Its aetiology is multifactorial, with mechanical, biochemical, genetic and lifestyle factors all contributing to involution of the 'bone-muscle unit'. Improved understanding of the interactions between muscle and bone could facilitate the development of new therapeutic agents which target muscle and bone as one. Together with existing pharmacological, nutritional and exercise-based therapies, this should enable a more holistic approach to osteosarcopenia in the future.

in which it can be prevented and treated. This review summarizes the latest developments in osteosarcopenia, with a focus on epidemiology, diagnosis, pathophysiology and management.

## Epidemiology

### Osteoporosis

Osteoporosis affects hundreds of millions of people worldwide; in the EU alone, around 27.6 million had osteoporosis in 2010, representing a prevalence of 6.6% and 22.1% respectively in men and women over 50 years of age. In the UK, around 536 000 fragility fractures occur each year, with hip (79 000), forearm (69 000) and spine (66 000) being the most frequent sites (Hernlund et al, 2013).

**Dr James Paintin**, Academic FY2 Doctor, MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton SO16 6YD

**Professor Cyrus Cooper**, Director, MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton, Professor of Epidemiology, Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences, University of Oxford, Oxford, and Professor of Rheumatology and Honorary Consultant Rheumatologist, Southampton General Hospital, Southampton

**Professor Elaine Dennison**, Professor of Musculoskeletal Epidemiology, MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton and Honorary Consultant in Rheumatology, Southampton General Hospital, Southampton

Correspondence to: Dr J Paintin ([jamespaintin@hotmail.co.uk](mailto:jamespaintin@hotmail.co.uk))

Hip fractures represent the most devastating consequence of osteoporosis, associated with significant morbidity and mortality. In the UK, permanent disability affects 50% of hip fracture patients (Chan et al, 2016), and UK data suggest a 1-year mortality rate after hip fracture of 34.4% and 27.7% respectively in men and women  $\geq 85$  years of age (Klop et al, 2014).

Osteoporotic fractures carry a significant economic burden because of the costs of treatment and long-term care. In the UK, osteoporotic fractures cost the health service £4.4 billion in 2010 (Compston et al, 2017). The economic burden of osteoporotic fractures is likely to rise significantly in the future, with costs predicted to swell by 25% in the EU between 2010 and 2025 (Hernlund et al, 2013).

## Sarcopenia

Sarcopenia is also generally a condition of ageing: above 50 years of age, muscle mass is lost at a rate of 1–2% per year, with strength falling more precipitously at 1.5–3% per year (von Haehling et al, 2012). Owing to a lack of consensus regarding the definition of sarcopenia, the prevalence rate varies widely; however, it is estimated at 5–13% for adults aged 60–70 years and 11–50% for adults  $> 80$  years old (von Haehling et al, 2012). Sarcopenia currently affects  $> 50$  million worldwide, which is expected to rise to  $> 200$  million over the next 40 years (Cruz-Jentoft et al, 2010).

Sarcopenia contributes to frailty, disability, premature mortality, psychosocial problems and worsened quality of life (Janssen et al, 2004; Fuggle et al, 2017). It carries a significant economic burden, with direct health-care costs attributed to sarcopenia estimated at \$18.5 billion in the USA in 2000 (Janssen et al, 2004). Another major consequence of sarcopenia is that it predisposes many elderly patients to falls, increasing the risk of fragility fractures (Edwards et al, 2015).

## Osteosarcopenia

Epidemiological measures of osteosarcopenia are fairly limited because of the recent origin of the term. However, one large study of 680 elderly adults with a history of falls found an osteosarcopenia prevalence of 37%, with these patients having a higher frequency of comorbidities, impaired mobility and depression (Huo et al, 2015).

Osteosarcopenia is also associated with significantly increased mortality: one recent study of 324 elderly Korean patients with hip fracture found a 1-year mortality rate of 15.1% in the osteosarcopenic patients, more than that of osteoporotic (5.1%) or sarcopenic (10.3%) patients alone (Yoo et al, 2018). Another study of 316 Chinese adults  $\geq 65$  years old found that 10.4% of men and 15.1% of women were osteosarcopenic, with the odds of frailty being significantly higher in the osteosarcopenic patients compared to osteoporosis or sarcopenia alone (Wang et al, 2015).

## Diagnosis

The diagnosis of osteosarcopenia represents co-existence of both osteoporosis and sarcopenia, as explained below.

## Osteoporosis

Osteoporosis is defined as a ‘systemic skeletal disease characterised by low bone mass and micro-architectural deterioration of bone tissue, with a consequent increase in bone fragility and susceptibility to fracture’ (Consensus Development Conference, 1993). In 1994, the World Health Organization developed specific criteria to define osteoporosis in postmenopausal women, based on bone mineral density (World Health Organization, 1994) (Table 1).

The reference technology to assess bone mineral density recommended by World Health Organization is dual energy X-ray absorptiometry applied to the femoral neck. The femoral neck was chosen as the reference site for diagnosis because of its higher predictive value of fracture risk (Compston et al, 2017); however, lumbar spine measurements are more effective at monitoring treatment-induced changes (Hernlund et al, 2013).

Although bone mineral density explains part of the risk of fragility fractures, many other risk factors operate independently of bone mineral density to increase fracture risk. These include low body mass index, previous osteoporotic fracture, parental history of hip fracture, smoking, steroids, rheumatoid arthritis and alcohol (Compston et al, 2017). Thus, fracture risk algorithms were developed that incorporate these risk factors, with or without bone mineral density, to optimize risk stratification (Binkley and Buehring, 2009).

The most famous of these is the Fracture Risk Assessment Tool (FRAX), which predicts an individual’s risk of major osteoporotic and hip fracture over the next 10 years. In the UK, the National Institute for Health and Care Excellence recommends screening all women and men over 65 and 75 years old respectively, assisted by either FRAX or QFracture (another fracture risk algorithm

**Table 1. World Health Organization classification of osteopenia and osteoporosis**

Bone mineral density category	Definition
Normal	Bone mineral density no greater than 1 standard deviation below the reference mean (T-score $\geq -1$ )
Osteopenia	Bone mineral density between 1 and 2.5 standard deviation below the reference mean (T-score $< -1$ but $> -2.5$ )
Osteoporosis	Bone mineral density 2.5 standard deviation or more below the reference mean (T-score $\leq -2.5$ )
Severe osteoporosis	Bone mineral density 2.5 standard deviation or more below the reference mean (T-score $\leq -2.5$ ) plus one or more fragility fractures

*Bone mineral density is expressed in standard deviations relative to a sex-specific, young-adult mean (known as the T-score). From World Health Organization (1994).*

in the UK). The National Osteoporosis Guideline Group has created intervention thresholds for treatment based on FRAX estimates, allowing better targeting of treatment to those patients with the highest risk (Compston et al, 2017).

**Sarcopenia**

Sarcopenia describes the age-related loss of skeletal muscle mass, strength and function (Fuggle et al, 2017). In 2009, the European Working Group on Sarcopenia in Older People was the first to develop an operational definition of sarcopenia incorporating measures of these parameters (Cruz-Jentoft et al, 2010), with other working groups providing their own definitions since (Fielding et al, 2011; Chen et al, 2014; Studenski et al, 2014) (Table 2).

Dual energy X-ray absorptiometry is currently the modality of choice to assess muscle mass, as it is quantitative, inexpensive, minimizes radiation exposure, and is widely available with its current role in osteoporosis (Binkley and Buehring, 2009). Bioimpedance analysis is another technique which is significantly more portable than dual energy X-ray absorptiometry – useful in immobile, bedridden patients (Cruz-Jentoft et al, 2010). The most common means to assess muscle strength is grip strength using a hydraulic dynamometer, with the best of three attempts recorded (Fuggle et al, 2017). This technique is simple, requiring little training, and also correlates well with lower limb strength (Oliveira and Vaz, 2015). Finally, the most reliable measure of muscle performance in clinical practice is gait speed, with the European Working Group on Sarcopenia in Older People advising a cut-off for sarcopenia of ≤0.8 m/s (Cruz-Jentoft et al, 2010).

The heterogeneity in definitional approaches towards sarcopenia has limited comparability between studies, with differences in measured parameters, regions of interest and the absence of a standardized reference population all limiting progress (Binkley and Buehring, 2009). However, a significant breakthrough occurred

in 2016, with the inclusion of a separate ‘sarcopenia’ diagnosis in the International Classification of Disease (Anker et al, 2016).

**Pathophysiology**

The pathogenesis of osteosarcopenia is multifactorial, with an array of mechanical, biochemical, genetic and lifestyle factors all contributing to involution of the ‘bone–muscle unit’ (Girgis et al, 2014).

**Mechanical**

The traditional view of the prominent mechanical interactions between muscle and bone is emphasized by the mechanostat hypothesis. This theory states that muscle imposes mechanical forces on bone, with a certain threshold dictating whether bone is formed or resorbed (Frost, 2003). An increase in muscle mass leads to stretching of collagen fibres and periosteum, leading to stimulation of bone growth (Kaji, 2014). Both osteoporosis and sarcopenia can result from reduced physical activity observed in ageing (Daly, 2017), lending support to the role of mechanical loading in preserving the bone–muscle unit.

**Biochemical**

Although mechanical forces clearly have a significant role in the pathogenesis of osteosarcopenia, more recent research suggests that paracrine or endocrine crosstalk is also at play.

Important hormones mediating this crosstalk include growth hormone, insulin-like growth factor-1, gonadal sex hormones and vitamin D, with age-related decreases contributing to the development of osteosarcopenia (Girgis et al, 2014). Muscle and bone also secrete certain factors, known as myokines and osteokines respectively, which aid the communication between muscle and bone (Girgis et al, 2014). An extensively researched myokine is myostatin, a member of the transforming growth factor beta (TGF-β) superfamily which inhibits skeletal muscle growth but also has effects on bone and tendon (Kaji, 2014).

**Table 2. Cut-off thresholds for the diagnosis of sarcopenia used by different international groups**

Working group	Diagnostic criteria for sarcopenia		
	Muscle mass (dual energy X-ray absorptiometry)	Muscle strength	Physical performance
European Working Group on Sarcopenia in Older People	Appendicular lean mass/height <sup>2</sup> Several similar cut-offs based on different reference groups, e.g. men: ≤7.23 kg/m <sup>2</sup> , women: ≤5.67 kg/m <sup>2</sup>	Grip strength Men: <30 kg, women: <20 kg Also provide cut-offs based on patient's body mass index	Gait speed Both sexes: ≤0.8 m/s
International Working Group on Sarcopenia	Appendicular lean mass/height <sup>2</sup> Men: ≤7.23 kg/m <sup>2</sup> , women: ≤5.67 kg/m <sup>2</sup>	N/A	Gait speed Both sexes: <1 m/s
Foundation of the National Institute of Health	Appendicular lean mass/body mass index Men: <0.789, women: <0.512	Grip strength Men: <26 kg, women: <16 kg	N/A
Asian Working Group on Sarcopenia	Appendicular lean mass/height <sup>2</sup> Men: <7.0 kg/m <sup>2</sup> , women: <5.4 kg/m <sup>2</sup>	Grip strength Men: <26 kg, women: <18 kg	Gait speed Both sexes: <0.8 m/s

From Cruz-Jentoft et al (2010), Fielding et al (2011), Chen et al (2014), Studenski et al (2014).

### Exercise has many positive effects on the older population, with one meta-analysis finding exercise reduced overall fracture risk by 51% in adults over 45 years old.

Bone–muscle crosstalk is mediated by several important signalling pathways. For example, the canonical Wnt- $\beta$ -catenin signalling pathway controls osteoblastic activity and is also involved in muscle regeneration (Oliveira and Vaz, 2015); the growth hormone–insulin-like growth factor-1 axis is a key regulator of bone and muscle growth, mediating its effects via several signalling pathways. It is hoped that exploiting these pathways will facilitate the development of new therapeutic agents in the future (Girgis et al, 2014).

#### Genetic

Genetic polymorphisms of various genes also contribute to the pathogenesis of osteosarcopenia. As reviewed by Kaji (2014), these include androgen receptor, oestrogen receptor, catechol-O-methyltransferase, insulin-like growth factor-1, vitamin D receptor and low-density-lipoprotein receptor-related protein. Moreover, since muscle and bone cells are both derived from mesenchymal stem cells, they are both influenced by similar genetic factors. Risk factors for both osteoporosis and sarcopenia have a heritability in the range of 60–70% (Kaji, 2014).

#### Lifestyle factors

Physical activity diminishes with ageing, with up to 80% of waking hours spent sedentary in the elderly (Harvey et al, 2015), leading to a loss of mechanical loading and therefore bone and muscle loss (Frost, 2003). Caloric intake diminishes by around 25% between the ages of 40 and 70 years (Nieuwenhuizen et al, 2010), with reduced vitamin D levels and protein intake correlated with declining muscle strength (Marty et al, 2017). Vitamin D deficiency also leads to increased risk of falls through its multiple effects on muscle and bone (Kaji, 2014). Smoking and alcohol are significant risk factors for osteoporosis, with intake of  $\geq 3$  units of alcohol/day increasing fracture risk in a dose-dependent fashion (Compston et al, 2017).

#### Management

Treatments for osteosarcopenia include exercise, improved nutrition and pharmacotherapy.

#### Exercise

Exercise has many positive effects on the older population, with one meta-analysis finding exercise reduced overall fracture risk by 51% in adults over 45 years old (Kemmler et al, 2013). The optimal approach for osteosarcopenia may be targeted multi-modal programmes which incorporate traditional and high-velocity progressive resistance training, weight-bearing impact exercises and challenging balance and/or mobility activities (Daly, 2017).

Exercise programmes are even effective for frailer adults, with a meta-analysis of elderly residents in long-term care demonstrating a 29% reduced risk of falls in those that underwent combined resistance and balance training programmes (Silva et al, 2013). Regular exercise also reduces bone loss, particularly at the femoral neck (Girgis et al, 2014).

#### Nutrition

Nutritional approaches to osteosarcopenia focus on vitamin D, calcium and protein intake.

Vitamin D supplementation can have many beneficial effects, including increased muscle strength, decreased mortality and falls, and functional improvement (Oliveira and Vaz, 2015). The evidence base for supplementation is stronger in older, institutionalized adults and those who are initially deficient (Girgis et al, 2014; Daly, 2017). Guidelines generally advocate 800–2000 IU/day in older adults, aiming for a target serum 25(OH)D level of at least 50 nmol/litre (20 ng/ml) (Girgis et al, 2014). Adequate calcium intake is also advised in patients with osteosarcopenia, with a recommended daily intake of 700–1200 mg (Compston et al, 2017). Meta-analyses show combined supplementation of vitamin D and calcium is safe, and effectively reduces fracture risk in older individuals (Daly, 2017).

In the USA, 5–12% of older men and 20–24% of older women consume inadequate protein (defined as  $<0.66$  g/kg body weight per day) (Berner et al, 2013). Moreover, there is evidence to suggest a blunted anabolic response to protein in the elderly, further compounding the problem of insufficient intake (Daly, 2017). Thus, a higher protein intake of 1.0–1.2 g/kg/day is recommended in the elderly, with at least 20–25 g of high-quality protein with each meal and post-exercise (Rizzoli et al, 2014).

#### Pharmacotherapy

Most pharmacotherapies for osteosarcopenia target bone, although several new therapies are being developed which also target muscle (Girgis et al, 2014). In the UK, the National Osteoporosis Guideline Group offers intervention thresholds for treatment based on 10-year fracture probability derived from FRAX (Compston et al, 2017).

For osteoporosis, bisphosphonates such as alendronate, risedronate and zoledronate remain the first-line treatment. They work via signalling pathways which induce apoptosis of osteoclasts, thus reducing bone resorption, and increasing bone mineral density (Chan et al, 2016). Other major pharmacological therapies for osteoporosis include denosumab (receptor activator of nuclear factor kappa B (RANKL) inhibitor that inhibits osteoclastogenesis), selective oestrogen receptor modulators such as raloxifene (partial oestrogen agonist selective for bone), and anabolic agents such as teriparatide (a recombinant form of parathyroid hormone). The latter treatment is restricted to use in high-risk fracture patients who are refractory to other treatment, because of its high expense (Chan et al, 2016).

Several new therapies are being developed which target muscle in addition to bone. Selective androgen receptor modulators, such as andarine, have anabolic effects in muscle and bone, and because of their high tissue selectivity, limit the androgenic side effects associated with testosterone therapy (Girgis et al, 2014). Other promising agents centre on the activin-signalling pathway, and include myostatin-neutralising antibodies/propeptide, recombinant follistatin, follistatin derivatives and soluble activin receptors (Girgis et al, 2014).

Pathways that centrally regulate bone and muscle, such as growth hormone–insulin-like growth factor-1 and androgen signalling, can also be targeted. Possible therapeutic agents include recombinant growth hormone (which increases lean mass and lumbar bone mineral density, but has safety issues), growth hormone secretagogues (which increase growth hormone and insulin-like growth factor-1 levels through more physiological means), and testosterone therapy (which has positive effects on muscle mass, strength and bone mineral density, but is limited by androgenic side effects and concerns about cardiovascular events and prostate cancer) (Girgis et al, 2014).

In the future, improved understanding of bone–muscle crosstalk will bring further novel drugs and biomarkers for osteoporosis and sarcopenia (Kaji, 2014), bringing great therapeutic benefits to patients with osteosarcopenia.

## Conclusions

Osteosarcopenia is a condition of growing importance with significant negative sequelae on both patients and society alike. For individuals, major consequences include falls and fractures, increased morbidity, mortality and disability, and reduced quality of life. For society, osteosarcopenia brings a staggering socioeconomic burden. With population ageing, the number of individuals  $\geq 60$  years old worldwide is set to increase from around 600 million in 2000 to 2 billion by 2050 (Cruz-Jentoft et al, 2010); as such, the incidence of osteosarcopenia will increase dramatically over the coming decades.

However, there are reasons to be hopeful. A growing consensus of the definition of sarcopenia will facilitate better recruitment to studies, and allow optimal measurement of muscle health outcomes in therapeutic trials (Edwards et al, 2015). As our understanding of the bone–muscle unit increases, there should be development of new therapies which embrace both osteoporosis and sarcopenia by targeting pathways that affect both muscle and bone. Lifestyle interventions can further help, with clear positive benefits of exercise and improved nutrition on osteosarcopenia. All of this should enable a more holistic approach to osteosarcopenia in the future, with benefits to millions of patients worldwide. **BJHM**

*Conflict of interest: none.*

Anker SD, Morley JE, von Haehling S. Welcome to the ICD-10 code for sarcopenia. *J Cachexia Sarcopenia Muscle*. 2016 Dec;7(5):512–514. <https://doi.org/10.1002/jcsm.12147>

## KEY POINTS

- Osteosarcopenia is a new syndrome describing the co-existence of osteoporosis and sarcopenia.
- Osteoporosis is characterized by low bone mass and micro-architectural deterioration of bone tissue.
- Sarcopenia describes the loss of muscle mass, strength and function.
- Individuals with osteosarcopenia are at higher risk of falls, fractures and institutionalization, resulting in significant socioeconomic costs.
- An array of mechanical, biochemical, genetic and lifestyle factors contribute to the pathogenesis of osteosarcopenia.
- Treatment approaches include exercise, improved nutrition and pharmacotherapy.
- An increasing awareness of bone–muscle interactions should facilitate the development of novel pharmacological agents in the future.

- Berner LA, Becker G, Wise M, Doi J. Characterization of dietary protein among older adults in the United States: amount, animal sources, and meal patterns. *J Acad Nutr Diet*. 2013 Jun;113(6):809–815. <https://doi.org/10.1016/j.jand.2013.01.014>
- Binkley N, Buehring B. Beyond FRAX: its time to consider sarcopenia. *J Clin Densitom*. 2009 Oct;12(4):413–416. <https://doi.org/10.1016/j.jocd.2009.06.004>
- Chan CKY, Mason A, Cooper C, Dennison E. Novel advances in the treatment of osteoporosis. *Br Med Bull*. 2016 Sep;119(1):129–142. <https://doi.org/10.1093/bmb/ldw033>
- Chen LK, Liu LK, Woo J et al. Sarcopenia in Asia: Consensus Report of the Asian Working Group for Sarcopenia. *J Am Med Dir Assoc*. 2014 Feb;15(2):95–101. <https://doi.org/10.1016/j.jamda.2013.11.025>
- Compston J, Cooper A, Cooper C et al; National Osteoporosis Guideline Group (NOGG). UK clinical guideline for the prevention and treatment of osteoporosis. *Archives of Osteoporosis*. 2017 Dec;12(1):43. <https://doi.org/10.1007/s11657-017-0324-5>
- Consensus Development Conference. Consensus development conference: Diagnosis, prophylaxis, and treatment of osteoporosis. *Am J Med*. 1993 Jun;94(6):646–650. [https://doi.org/10.1016/0002-9343\(93\)90218-E](https://doi.org/10.1016/0002-9343(93)90218-E)
- Crepaldi G, Maggi S. Sarcopenia and osteoporosis: A hazardous duet. *J Endocrinol Invest*. 2005;28(10) Suppl:66–68.
- Cruz-Jentoft AJ, Baeyens JP, Bauer JM et al. Sarcopenia: European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia in Older People. *Age Ageing*. 2010 Apr;39(4):412–423. <https://doi.org/10.1093/ageing/afq034>
- Daly RM. Exercise and nutritional approaches to prevent frail bones, falls and fractures: an update. *Climacteric*. 2017 Mar 04;20(2):119–124. <https://doi.org/10.1080/13697137.2017.1286890>
- Edwards MH, Dennison EM, Aihie Sayer A, Fielding R, Cooper C. Osteoporosis and sarcopenia in older age. *Bone*. 2015 Nov;80:126–130. <https://doi.org/10.1016/j.bone.2015.04.016>
- Fielding RA, Vellas B, Evans WJ et al; International Working Group on Sarcopenia. Sarcopenia: an undiagnosed condition in older adults. Current consensus definition: prevalence, etiology, and consequences. International working group on sarcopenia. *J Am Med Dir Assoc*. 2011 May;12(4):249–256. <https://doi.org/10.1016/j.jamda.2011.01.003>
- Frost HM. Bones mechanostat: A 2003 update. *Anat Rec*. 2003 Dec;275A(2):1081–1101. <https://doi.org/10.1002/ar.a.10119>
- Fuggle N, Shaw S, Dennison E, Cooper C. Sarcopenia. *Best Pract Res Clin Rheumatol*. 2017 Apr;31(2):218–242. <https://doi.org/10.1016/j.berh.2017.11.007>
- Girgis CM, Mokbel N, DiGirolamo DJ. Therapies for musculoskeletal disease: can we treat two birds with one stone? *Curr Osteoporos Rep*. 2014 Jun;12(2):142–153. <https://doi.org/10.1007/s11914-014-0204-5>
- Harvey JA, Chastin SFM, Skelton DA. How sedentary are older

- people? A systematic review of the amount of sedentary behavior. *J Aging Phys Act.* 2015 Jul;23(3):471–487. <https://doi.org/10.1123/japa.2014-0164>
- Hernlund E, Svedbom A, Ivergård M et al. Osteoporosis in the European Union: medical management, epidemiology and economic burden. *Arch Osteoporos.* 2013 Dec;8(1-2):136. <https://doi.org/10.1007/s11657-013-0136-1>
- Huo YR, Suriyaarachchi P, Gomez F et al. Phenotype of osteosarcopenia in older individuals with a history of falling. *J Am Med Dir Assoc.* 2015 Apr;16(4):290–295. <https://doi.org/10.1016/j.jamda.2014.10.018>
- Janssen I, Shepard DS, Katzmarzyk PT, Roubenoff R. The healthcare costs of sarcopenia in the United States. *J Am Geriatr Soc.* 2004 Jan;52(1):80–85. <https://doi.org/10.1111/j.1532-5415.2004.52014.x>
- Kaji H. Interaction between muscle and bone. *J Bone Metab.* 2014;21(1):29–40. <https://doi.org/10.11005/jbm.2014.21.1.29>
- Kemmler W, Häberle L, von Stengel S. Effects of exercise on fracture reduction in older adults. *Osteoporos Int.* 2013 Jul;24(7):1937–1950. <https://doi.org/10.1007/s00198-012-2248-7>
- Klop C, Welsing PMJ, Cooper C et al. Mortality in British hip fracture patients, 2000–2010: A population-based retrospective cohort study. *Bone.* 2014 Sep;66:171–177. <https://doi.org/10.1016/j.bone.2014.06.011>
- Marty E, Liu Y, Samuel A, Or O, Lane J. A review of sarcopenia: enhancing awareness of an increasingly prevalent disease. *Bone.* 2017 Dec;105:276–286. <https://doi.org/10.1016/j.bone.2017.09.008>
- Nieuwenhuizen WF, Weenen H, Rigby P, Hetherington MM. Older adults and patients in need of nutritional support: review of current treatment options and factors influencing nutritional intake. *Clin Nutr.* 2010 Apr;29(2):160–169. <https://doi.org/10.1016/j.clnu.2009.09.003>
- Oliveira A, Vaz C. The role of sarcopenia in the risk of osteoporotic hip fracture. *Clin Rheumatol.* 2015 Oct;34(10):1673–1680. <https://doi.org/10.1007/s10067-015-2943-9>
- Rizzoli R, Stevenson JC, Bauer JM et al; ESCEO Task Force. The role of dietary protein and vitamin D in maintaining musculoskeletal health in postmenopausal women: A consensus statement from the European Society for Clinical and Economic Aspects of Osteoporosis and Osteoarthritis (ESCEO). *Maturitas.* 2014 Sep;79(1):122–132. <https://doi.org/10.1016/j.maturitas.2014.07.005>
- Silva RB, Eslick GD, Duque G. Exercise for falls and fracture prevention in long term care facilities: a systematic review and meta-analysis. *J Am Med Dir Assoc.* 2013 Sep;14(9):685–689.e2. <https://doi.org/10.1016/j.jamda.2013.05.015>
- Studenski SA, Peters KW, Alley DE et al. The FNIH sarcopenia project: rationale, study description, conference recommendations, and final estimates. *J Gerontol A Biol Sci Med Sci.* 2014 May;69(5):547–558. <https://doi.org/10.1093/geron/69.5.547>
- von Haehling S, Morley JE, Anker SD. From muscle wasting to sarcopenia and myopenia: update 2012. *J Cachexia Sarcopenia Muscle.* 2012 Dec;3(4):213–217. <https://doi.org/10.1007/s13539-012-0089-z>
- Wang YJ, Wang Y, Zhan JK et al. Sarco-osteoporosis: prevalence and association with frailty in Chinese Community-dwelling older adults. *Int J Endocrinol.* 2015;2015:1–8. <https://doi.org/10.1155/2015/482940>
- World Health Organization. Assessment of fracture risk and its application to screening for postmenopausal osteoporosis. Report of a WHO Study Group. *World Health Organ Tech Rep Ser.* 1994;843:1–129.
- Yoo JI, Kim H, Ha YC, Kwon HB, Koo KH. Osteosarcopenia in patients with hip fracture is related with high mortality. *J Korean Med Sci.* 2018;33(4):e27. <https://doi.org/10.3346/jkms.2018.33.e27>

Organised by  
BRITISH JOURNAL OF  
**HOSPITAL  
MEDICINE**

6th national conference

# Osteoporosis 2018

Queen Elizabeth II Centre, London  
18th June 2018

Highlights will include:

- Developments in imaging practice in the UK **Dr Karen Knapp, Exeter**
- Incorporating fall prevention into osteoporosis management **Sally Ann Belward, Solent**
- Calcium and vitamin D for osteoporosis management **Dr Harish Datta, Newcastle**
- Case Studies: Investigating secondary causes of osteoporosis **Professor David Reid, Aberdeen**
- Case Studies: Osteoporosis management in men **Dr Stephen Tuck, Newcastle**



Follow us on Twitter: @MAHealthEvents  
Tweet about the conference: #osteoporosis18

To book your place:

Call us on +44(0)20 7501 6761

[www.mahealthcareevents.co.uk/osteoporosis2018](http://www.mahealthcareevents.co.uk/osteoporosis2018)

