

titration with a nonacidic quercetin solution, only one equivalent point (pK_2) would be attainable.

Thermodynamic parameters for the dissociation of quercetin were calculated from the slopes of the straight lines $\log K_d^0 = f\left(\frac{1}{T}\right)$: $\Delta H = 72.5$ kJ/mol, $\Delta S = 0.17$ kJ/K mol

and $\Delta G = 20.4$ kJ/mol, for the first dissociation constant, and $\Delta H = 86.3$ kJ/mol, $\Delta S = 0.11$ kJ/Kmol and $\Delta G = 54.1$ kJ/mol, for the second dissociation constant of quercetin.

Experimental

1. Apparatus

A pH-meter (pHM-82 Radiometer Copenhagen), accuracy of ± 0.001 pH, equipped with the combined electrode (No. CW.733 Serial No. 35162, Russel) was used. The temperature was controlled within ± 0.2 K by a circulating water thermostat (Serie U, MLW Freital, Germany).

2. Reagents

Quercetin; absolute ethanol, NaOH, NaNO_3 , by Merck (Darmstadt, Germany); all of *p.a.* grade. The solution of NaOH (carbonate free) was standardized by potentiometric titration against A.R. potassium hydrogen phthalate (dried 1^h at 120°C). The solution of HCl, added to the quercetin solution before titration, was standardized by potentiometric titration with 0.514 M NaOH.

3. General procedure

Titration of 0.0025 M quercetin solution with 0.514 M NaOH were carried out in a 51 ml glass vessel closed by a cover with five holes. Through the holes on the lid, the thermometer, the combined electrode, and the nitrogen inlet tubes were inserted. One of the holes was closed with a glass stopper, while the titrant was successively introduced through the remaining hole. The measurements were performed at three temperatures (28 , 34 and 39°C) and at three values of ionic strength (0.0024 , 0.034 and 0.062 M). The ionic strength was adjusted by 2 M NaNO_3 . Each potentiometric titration, with about 180 experimental points, was done in triplicate.

Acknowledgement: We gratefully acknowledge the partial financial support in grant no. 1458 of the Ministry for Science, Technology and Development of the Republic of Serbia.

References:

- Zheng, W.; Wang, Y. S.: *J. Agric. Food Chem.* **49**, 5165 (2001)
- Nevckar, E. M.; Nazarenko, V. A.: *Z. Analit. himii* **9**, 1699 (1972)
- Escandar, M. G.; Sala, L. F.: *Can. J. Chem.* **69**, 1994 (1991)
- Sauerwald, N.; Schwenk, M.; Polster, J.; Bengsch, E.: *Z. Naturforsch.* **53b**, 315 (1998)
- Rossotti, H.: *The study of Ionic Equilibria*, Longman, London 1978.
- Rossotti, F. J. C.; Rossotti, H.: *The Determination of Stability Constants*, McGraw-Hill, New York 1961

Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Comenius University, Bratislava, Slovak Republic

Thermodynamic study of local anesthetics based on heptacainium chloride derivatives

Study of local anesthetics. Part 163*

F. ANDRIAMAINTY, J. ČIŽMÁRIK

Received January 23, 2003, accepted January 27, 2003

Dr. Fils Andriamainty, PhD., Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Comenius University, Odbojárov 10, 832 32 Bratislava, Slovak Republic

Pharmazie **58**: 440–441 (2003)

We have previously studied the critical micellar concentration (CMC) of the derivatives of piperidinoethylesters of 2-alkoxyphenylcarbamic acid. We found that in a homologous series of alkyloxy substituents, the CMC decreases with the number of the carbon atoms up to the heptyloxy ($n = 7$) and then increases [1].

The aim of the present paper was to study the thermodynamic parameters of local anesthetics of heptacainium chloride derivatives in the medium of distilled water, using UV spectrophotometry at a temperature range $T = 294$ – 318 K. Based on our results (Table 1), the thermodynamic magnitudes were calculated ($-\Delta H^\circ$, $-\Delta G^\circ$, $-\Delta S^\circ$) according to the "phase separation (PS_1 and PS_2)" model [2]. The equation $\text{CMC} = f(T) = A + BT + CT^2$ represents the dependence of the critical micellar concentration upon temperature at $\text{pH} \approx 4.5$ – 5 , where (A , B , C) = constants of the second degree polynomial and T = absolute temperature. Gibbs energy change can be estimated according to the equation;

$$\Delta G^\circ = \gamma RT \ln (\text{CMC}) \quad (1)$$

where R = gas constant and γ = degree of counterion binding (if $\gamma = 1$ the anti-ions are completely ionized, if $\gamma = 2$ all the anti-ions are bound to micelles). The enthalpy of micellization is defined by the equation

$$\Delta H^\circ = -\gamma RT^2 [\partial \ln (\text{CMC}) / \partial T] \quad (2)$$

and the entropy contribution of micellization can be calculated as follows:

$$\Delta S^\circ = (\Delta H^\circ - \Delta G^\circ) / T \quad (3)$$

ΔG° , ΔH° , ΔS° values are listed in Table 2.

Table 1: CMC (mol/l) values in relation to temperature T (K)

T (K)	CMC (mol/l)			
	$n = 5$	$n = 7$	$n = 8$	$n = 9$
294	3.990×10^{-4}	3.151×10^{-4}	3.914×10^{-4}	4.763×10^{-4}
298	4.016×10^{-4}	3.199×10^{-4}	4.122×10^{-4}	4.841×10^{-4}
303	4.145×10^{-4}	3.212×10^{-4}	4.198×10^{-4}	4.906×10^{-4}
308	4.311×10^{-4}	3.258×10^{-4}	4.288×10^{-4}	5.029×10^{-4}
313	4.399×10^{-4}	3.332×10^{-4}	4.419×10^{-4}	5.084×10^{-4}
318	4.424×10^{-4}	3.545×10^{-4}	4.539×10^{-4}	5.243×10^{-4}

Table 2: Thermodynamic parameters of local anesthetics of heptacainium chloride derivates in distilled water medium and temperature range $T = 294\text{--}318\text{ K}$

Substance	T (K)	ΔG° (kJ/mol)		ΔH° (kJ/mol)		ΔS° (kJ/mol)	
		PS ₁	PS ₂	PS ₁	PS ₂	PS ₁	PS ₂
13 (n = 5)	294	−19.14	−38.28	−3.10	−6.20	16.04	32.08
	298	−19.36	−38.71	−3.58	−7.16	15.78	31.56
	303	−19.62	−39.23	−4.20	−8.41	15.41	30.82
	308	−19.87	−39.73	−4.87	−9.74	15.00	30.00
	313	−20.10	−40.21	−5.57	−11.13	14.54	29.07
	318	−20.33	−40.66	−6.30	−12.61	14.03	28.05
19 (n = 7)	294	−19.69	−39.38	0.91	1.82	20.60	41.20
	298	−19.96	−39.92	−0.44	−0.88	19.52	39.04
	303	−20.27	−40.55	−2.23	−4.46	18.04	36.09
	308	−20.56	−41.11	−4.14	−8.28	16.42	32.83
	313	−20.81	−41.61	−6.17	−12.34	14.63	29.27
	318	−21.02	−42.05	−8.33	−16.66	12.70	25.39
22 (n = 8)	294	−19.08	−38.17	−2.02	−4.04	17.06	34.12
	298	−19.31	−38.62	−2.51	−5.03	16.80	33.60
	303	−19.59	−39.18	−3.16	−6.33	16.42	32.85
	308	−19.85	−39.71	−3.85	−7.70	16.00	32.00
	313	−20.11	−40.21	−4.58	−9.16	15.53	31.05
	318	−20.35	−40.70	−5.35	−10.70	15.00	29.99
25 (n = 9)	294	−18.69	−37.39	−2.14	−4.29	16.55	33.10
	298	−18.92	−37.84	−2.42	−4.83	16.50	33.01
	303	−19.19	−38.38	−2.77	−5.54	16.42	32.84
	308	−19.46	−38.92	−3.15	−6.29	16.31	32.63
	313	−19.72	−39.44	−3.54	−7.08	16.18	32.36
	318	−19.98	−39.95	−3.96	−7.92	16.02	32.04

Based on the results presented it can be generalized that

- ΔG° values are negative and decline slightly with temperature.
- Depression of standard molar enthalpy ΔH° is more significant at more negative values. This means that the micellization process becomes more exothermic with increasing temperature.
- ΔS° values are positive and decline with increasing temperature.
- It was noticed that the dependence of ΔG° on n is paraboloid and for all temperature intervals has a minimum at ($n = 7$).

These conclusions correspond with the results, which had been reached for the basic substance heptacainium chloride [3].

Experimental

The derivatives of piperidinoethyl esters of 2-alkoxyphenylcarbamic acid (substances 13 where $R = C_5H_{11}$, 19 where $R = C_7H_{15}$, 22 where $R = C_8H_{17}$, and 25 where $R = C_9H_{19}$) were synthesised as described earlier [4].

Distilled water was used to prepare stock solutions (10^{-3} mol/l). From the appropriate stock solutions various concentrations of diluted solutions were prepared with $pH \approx 4.5\text{--}5.0$ at temperature range $T = 294\text{--}318\text{ K}$. pH was measured with a pH meter (Portamess 943 pH , Elektronische Messgeräte GmbH Co., Berlin) and the temperature was controlled by a Thermostat (Veb ML W Prüfgeräte-Werk Medingen/Sity/Freital (BRD). The critical micellar concentration was determined using an HP 8452 A Diode Array spectrophotometer (Hewlett Packard, BRD) [5].

*Part 162: Pharmazie **58**, 288 (2003)

This study was supported by the VEGA grant 1/8213/2001

References

- 1 Andriamainty, F.; Čižmárik, J.: Pharmazie **58**, 288 (2003)
- 2 Evans, D. F.; Wightmann, P. J.: Coll. Inter. Sci. **16**, 484 (1961)

3 Andriamainty, F.; Čižmárik, J.: Pharmazie **54**, 629–630 (1999)

4 Čižmárik, J.; Borovanský, A.: Chem. Pap. **29**, 119 (1975)

5 Ščukin, E. D.; Percov, A. V.; Amelinová, E. A.: Acad. Praha, 1. Vyd., 303 (1990)