

Protein Research Department, Genetic Engineering and Biotechnology Research Institute, Mubarak City for Scientific Research and Technology Applications/Pharmaceutics Department, Faculty of Pharmacy, King Saud University, Kingdom of Saudi Arabia

The philosophy behind exo/endo/existing antioxidants and our built-in oxidant and antioxidant system

A.A. AMARA

Received February 28, 2010, accepted May 28, 2010

Asst. Prof. Amro Abd Al Fattah Amara, Protein Research Department, GEBRI, MuCSAT/Pharmaceutics Department, Faculty of Pharmacy, KSU, KSA.
amroamara@web.de

Pharmazie 65: 711–719 (2010)

doi: 10.1691/ph.2010.0557

Free radicals (FRs) and oxidants (Os) have different vital activities in our bodies. Biological redox reactions are the major source for FRs-Os. In normal cases, they are under the control of our body's existing antioxidant (antiO) system. If their amounts exceeds a certain limit, they start to attack and damage various macromolecules and cells. In the case of DNA damage, dangerous diseases such as cancer "the human killer" can be elevated. The expression and regulation of the genes coding for the endogenous (endo) antiO enzymes are important for our health. Genetic polymorphism between different populations reflects the differences between individuals and populations abilities to produce antiOs. This review will discuss general aspects about different types of FRs-Os and antiOs as well as their direct and indirect interaction with genetic materials. The benefit of the antiOs, which can do a lot for our health, will be highlighted.

"Variation is a unique criterion in the biological system." "A certain level of balance is required in all aspects of the life including food." When you start to read this review you will find something different! Is it the style? Or it may be something else! Each of us has his fingerprint, and our genes are different too. This is the life. The author.

1. Introduction

1.1. Free radicals and oxidants, our enemies or friends?

Harman (1956) was the first to describe the damage caused by Free radicals (FRs) and oxidants (Os) to life processes. He correlated the damage in tissues and cells, especially during ageing processes, with the activities of FRs-Os.

FRs-Os in the bio-system gain their importance from their ability to damage key biological sites (Pacher et al. 2007; Sies 1997). However, they are involved in many essential and vital processes. They prevent diseases by assisting the immune system during phagocytosis process, mediating cell signaling, and playing an essential role in apoptosis as a part of normal aerobic and xenobiotic metabolism (Bahorum et al. 2006; Valko et al. 2004, 2007; Droge 2002; Seifried et al. 2007). FRs-Os are formed from different molecules via the breakage of their chemical bonds. Each fragment keeps one electron, by cleavage of a radical to give another radical and via redox reactions (Bahorum et al. 2006; Valko et al. 2004, 2007; Droge 2002; Seifried et al. 2007).

1.2. Enzymatic and non-enzymatic FRs-Os formation

FRs have at least one unpaired electron and are capable of independent existence (Ghafourifar and Cadenas 2004). They are high-energy particles that ricochet wildly and damage cells.

FRs include hydroxyl (OH^\bullet), superoxide ($\text{O}_2^{\bullet-}$), nitric oxide (NO^\bullet), nitrogen dioxide (NO_2^\bullet), peroxy (ROO^\bullet) and lipid peroxy (LOO^\bullet). Os such as hydrogen peroxide (H_2O_2), ozone (O_3), singlet oxygen ($^1\text{O}_2$), hypochlorous acid (HOCl), nitrous acid (HNO_2), peroxy nitrite (ONOO^-), dinitrogen trioxide (N_2O_3), and lipid peroxide (LOOH) are not FRs but they are unstable molecules (Pacher et al. 2007; Sies 1997; Bahorum et al. 2006; Valko et al. 2004; Willcox et al. 2004; Genestra 2007). They cause damage to our biological system. When this damage exceeds the existing antiOs system [the endogenous (endo) antiOs plus the exogenous (exo) antiOs supplemented through diet] as well as the system for DNA regeneration and correction, different health problems begin to appear and vital organs will be affected (Bahorum et al. 2006; Valko et al. 2004, 2007; Droge 2002; Seifried et al. 2007). Those dangerous FRs-Os could appear as a result of excessive exercise, stress, alcohol, cigarettes smoking, air pollution, food etc. (Bahorum et al. 2006; Valko et al. 2004, 2007; Droge 2002; Seifried et al. 2007; Ghafourifar and Cadenas 2005). When the sources of the FRs-Os are irremovable, exo-antiOs should be supplemented through diet or drugs (Ghafourifar and Cadenas 2005). The endo/exo FRs-Os sources could be controlled or reduced by different strategies. The deficiency in one or more of the antiOs gene(s), FRs-Os controlling systems, essential pathway(s), as well as loss of precursor(s) or mutation(s), will result in an increase in FRs-Os. The amount of antiOs and their related precursors in our bodies could be optimized through diets. Food is the most promising candidate as a safe source of antiOs. "From *vivo* to *vivo* is usually better than from *chemo* to *vivo*". However, the simplest solution will be to stop all the sources of FRs-Os as causative agents. Adequate amounts of exo-antiOs will compensate for any deficiency in the endo-antiOs. Key genes are responsible for both FRs-Os and antiOs production and control (Valko et al. 2007).

AntiO genes are the major players for maintaining our health (Valko et al. 2007).

The balance between beneficial and harmful effects of free radicals is achieved by mechanism(s) called “redox regulation”. The process of “redox regulation” protects living organisms from various types of FRs-O stresses and maintains “redox homeostasis” by controlling the redox status *in vivo* and during human diseases (Droge 2002; Valko et al. 2007). FRs-Os are generated from end sources such as mitochondria, respiratory chain, immune system activity, inflammation, mental stress, excessive exercise, ischemia, infections, cancer, change in the intestine microflora, ageing, genetic susceptibility, phagocytosis, and Cytochrome enzymes 450 (CYP450) (Valko et al. 2004, 2007; Droge 2002; Seifried et al. 2007; Ghafourifar and Cadenas 2004). FRs-Os can also be generated from exo-sources such as air and water pollution, cigarette smoke, alcohol, heavy or transition metals (Cd, Hg, Pb, Fe, As), certain drugs such as ciclosporine, tacrolimus, gentamycin and bleomycin, industrial solvents, cooking (smoked meat, used oil, fat) and radiation (Pacher et al. 2007; Bahorun et al. 2006; Valko et al. 2004; Genestra 2007; Davis et al. 2009; de Oliveira et al. 2009; Halliwell 2007; Young 2001; Valko et al. 2005; Parthasarathy 1999). After their penetration into the body by different routes, these exo-compounds are decomposed or metabolized into FRs-Os (Valko et al. 2004, 2007; Droge 2002; Seifried et al. 2007; Ghafourifar and Cadebas 2005). Diet, lifestyle, and the exposure levels are important factors in determining the capacity of an organism to mount a protective response (Valko et al. 2004; Droge 2002). Different diseases have been associated with FRs-Os such as cancer, arterosclerosis, pulmonary disease, cardiovascular diseases, neurodegenerative diseases, allergies, metabolic diseases, aging, insulin resistance, Down’s syndrome, familial amyotrophic lateral sclerosis (ALS), transplantation complications, and many other conditions (Pacher et al. 2007; Bahorun et al. 2006; Valko et al. 2007; Droge 2002; Young 2001). It is important to highlight that any new disease will be an additional source of FRs-Os! Chemoprevention by nutrients, CYP450 inhibitors, and antiO enzyme inducers, nutraceutical food, good life style, etc., have been successfully used to inhibit the formation of reactive species (Wender et al. 1981).

1.3. Our food and the problems

In both plants and animals, energy is extracted from food molecules by a process of gradual oxidation or controlled burning. Mitochondria harness the energy from the oxidation of food molecules, such as sugars, to produce adenosine triphosphate (ATP), the basic chemical fuel that powers most of the cell’s activities (Valko et al. 2007; Droge 2002; Davis et al. 2009; de Oliveria et al. 2009; Raha and Robinson 2000). The mitochondria as a major source for FRs-O generation shows that we have a built-in and a continuous source of FRs-Os. Unfortunately, food is the major source for both antiOs and FRs-Os. Food is transformed to energy and FRs-Os, which results in the inverse correlation between metabolic rate and the ageing process (Valko et al. 2007; Nakabeppu et al. 2006; Stadtman 1992; Raha and Robinson 2000; Davies 1995; Chui and Carol 2008; McCann 1997; Schrauzer and Peter 2009).

When the existing-antiO system starts to work, it needs energy, which is derived from different metabolic pathways. Some of these intermediate pathways are the source of FRs-Os, too. Continuous activation of the endo-antiO system is an extra load on genetic material, which could cause DNA damage. The DNA damage could be due to guanine residues resulting in 8-oxo-deoxyguanosine (8-oxo-dG). These induce G:C to T:A transversions that have been observed in oncogenes and tumor

suppressor genes (eg p53) known to have significant roles in carcinogenesis. The mutant generation rate in the presence of high levels of FRs-Os is greater than that produced in normal cases (Valko et al. 2005; Droge 2002; Davies 1995).

1.4. CYP450 one coin with two faces

CYP450s is another built-in FRs-O generator. The CYP450 enzymes convert organic chemicals to a more hydrophilic metabolite, while the concomitant reduction of the other oxygen atom forms water molecules. As some of most potent precarcinogens are converted into their ultimate carcinogenic form by the CYP450 system, these electrophilic metabolites can exert their biological effect by covalent interaction with cellular macromolecules, with the critical target most likely being DNA to yield DNA adducts. However, the action of CYP450 enzymes are not usually hazardous. These enzymes also play a central role in the detoxification of foreign compounds by increasing their water solubility and facilitating their excretion. The case of CYP450 is a good example of FRs-Os/antiOs possible interaction with the exo/endo effective candidates. CYP450 is able to detoxify large amounts of toxic material and exotic chemicals; it is also able to convert some pre-carcinogenic compounds to carcinogenic ones (Awney et al. 1997). We should not forget that although our biological system is intelligent, it is not intelligent enough to cover all our mistakes.

1.5. AntiOs, the hidden soliders

AntiOs are substances that slow down or prevent oxidation and reactions promoted by FRs-Os (Sies 1997; Bahroun et al. 2006; Valko et al. 2007; Seifried et al. 2007; Willcox et al. 2004; de Oliveira et al. 2009). In our bodies, there are different types of endo antiO enzymes such as superoxide dismutase (SOD), which was the first antiO enzyme to be discovered (McCord and Fridovich 1969; Wang 2009). Catalase (CAT), glutathione peroxidase (GPx) (both of which reduce peroxide to water), glutathione S-transferases, hemeoxygenase-1 (HO-1), thiol-specific antiO enzyme and macrophage stress protein all play a central role in protection against FRs-Os (Siow et al. 1995; Yim et al. 1994; Sachdev and Davies 2008; Vertuani et al. 2004; Chaudiere and Ferrari-Iliou 1999; Sies 1993). As a network, glutathione and vitamin C can act in concert to alleviate a variety of O stresses. Vitamin C is able to regenerate tocopherol in the lipid phase (Jindal 2008; Joyner 2009). β -Carotene and tocopherol act synergistically against lipid peroxidation. Polyphenols, such as flavonoids, provide antiO protection, which is enhanced by vitamin C. AntiO vitamins play major roles in the protection against FOs-Os and different related diseases (Jindal 2008; Platz 2009; Rayman et al. 2009; Schrauzer 2009; Shaw and Wan-Hsuan 2009).

Exogenous antiOs will be of great benefit to the body. They will save cellular energy and genetic materials, and will improve our health significantly. In the presence of a continuous extra dosage of FRs-Os, antiOs could give reverse results. Studies, which have been conducted on cigarette smokers, proved that β -carotenes could induce cancer rather than prevent it (De Luca and Ross, 2009).

This shows clearly who the players in the FRs-Os/antiOs struggle are: the FRs-Os and exo/endo/existing antiOs. Macromolecules under attack are also included. When attacked, they cause changes in their own cell system. The time of exposure to FRs-Os is another important factor. However, the most important part is the genetic material, which cannot be easily recovered. Other candidates, such as nutrients, environment, mental disorder, stress, different physiological and pathological disorders,

etc., are involved. The antiOs/FRs-Os struggle is individually based and should be studied case-by-case (Ortega 2006).

1.6. Is our endo-antiOs system able to stand-alone?

In fact, our endogenous antiO system is diet dependent. Most of our endo-antiO systems' platforms are derived from our diet (Traber and Packer 1995).

Other candidates are involved in protecting us from FRs-Os. Melanin is increased in the skin as a response to exposure to ultraviolet light (Cutler 1984). Human antiO defenses are effective but they are not infallible, and oxidative damage to key biological sites occurs, accumulates with age, and contributes to senescence and age-related diseases (Benzie 2003; Ames et al. 1993; Beckman and Ames 1998; Halliwell and Gutteridge 1999; Vasdev and Gill 2006).

The level of glutathione, the major cellular antiO in the body, declines with ageing. Glutathione is a protein, which is easily digested and cannot be effectively supplemented orally. Food, which contains high amounts of cysteine, will be better than that which does not contain cysteine. The cell membrane transports cysteine ten times faster than cystine. This shows how food and its constituents are efficacious in support of the antiO system (Vasdev and Gill 2006). Niacin is required for synthesis of nicotinamide adenine dinucleotide hydrogen (NADH)/NADPH and sulfur (methionine and cysteine) for synthesis of GSH (Jacob 1995).

Scurvy is a clear example about our body's demand for antiOs (Valko et al. 2005, 2006, 2007; Davies 1995).

2. Classification of human antiOs

While the major classification for the antiOs is the endo/exo/existing system, other classifications for the biological antiOs will be of great importance. There are four other major classifications for antiOs.

2.1. Enzymes and low molecular weight antiOs (LMWA)

Biological antiOs may be classified into two principal groups; enzymes such as CAT, Prx and SOD or as LMWA, such as tocopherols, ascorbic acid and glutathione. Metal-containing SODs, including Cu, Zn-SOD (SOD1), Mn-SOD (SOD2), and extracellular Cu, Zn-SOD (SOD3). SOD provides the first line of defense against oxidative stress and catalyzes the dismutation of superoxide anion to molecular oxygen and hydrogen peroxide. SOD1 is mainly expressed in astrocytes and neurons, whereas SOD2 is predominantly localized in neurons (Jacob 1995; McCord and Fridovich 1969; McCord 1976; Marklund 1982; Johnson and Giulivi 2005). Catalase is an intracellular antiO enzyme that catalyzes the conversion of H₂O₂ into water and molecular oxygen (Dringen et al. 2005).

The family of Prx's consists of six thio-specific antiO proteins that are involved in the enzymatic degradation of H₂O₂, organic hydroperoxides, and peroxynitrite (Rhee et al. 2005).

Prx's are abundantly expressed in the cytosol; however, several isoforms can also be found in mitochondria, peroxisomes, nuclei, and membranes (Hofmann et al. 2002).

The glutathione system is one of the most important antiO systems for most tissue types *in vivo*. It has been demonstrated in astrocytes that when hydrogen peroxide clearance by catalase is inhibited, the glutathione system almost completely compensates for this (Liddell et al. 2006 a,b). Neuronal cells contain high concentrations of the tripeptide glutathione (GSH) and display high GPx and glutathione reductase activity (Rhee et al. 2005). Inducible hemoxygenase-1 (HO-1) and constitu-

tive HO-2 belong to the family of heat shock proteins and protect brain cells from oxidative stress. NADH quinone oxidoreductase (NQO1) is mainly expressed in astrocytes and brain endothelial cells and, together with NQO2, prevents the generation of FRs-Os (Siegel and Ross 2000; van Muiswinkel et al. 2004; van Horssen et al. 2006).

Micronutrients play an important role in the antiO defense system such as vitamin C (ascorbic acid) which is an effective quencher of many oxyradicals in the body and a protector against scurvy. Vitamin E (tocopherol) provides antiO protection in the body, especially the protection of the unsaturated fatty acids of the cell membrane. Carotenoids in fruits and vegetables are also believed to provide antiO protection to lipid rich tissues; glucose, pyruvate, bilirubin, carnosine, and the methyl rich compounds carnitine, choline, betaine, and methionine are involved. A variety of minor plant constituents such as flavonoids, limonoids and procyanins exhibit antiO activity (Valko et al. 2005, 2007; Droge 2002; Davies 1995; Halliwell and Gutteridge 1999; Lauterburg et al. 1984).

2.2. AntiO classification based on action

AntiOs have three main modes of actions include preventing, chain-breaking/scavenging and chelating mechanisms. In the case of preventing the production *ab initio* of FRs-Os and their destruction, catalase is an example. Chain-breaking/scavenging the radicals examples are vitamin E and C, as well as enzyme systems to control FRs-Os (Hammadeh et al. 2008). Carnosine as a chelating agent is an efficient singlet-oxygen scavenger, quenching singlet oxygen more effectively than histidine. Carnosine, anserine, and histidine have a protective effect against (γ -irradiation, which gives rise to oxidative DNA damage, protects against oxidative stress, and suggests that it and related compounds may serve as natural antiOs in skeletal muscle and brain (Gulewitsch and Amiradzibi 1900).

2.3. AntiO classification based on solubility in water or lipids

Vitamin C is present in the body's aqueous compartments such as the blood plasma and cell cytosol. Vitamin E (tocopherol) is present in the body's lipid phases. Carotenoids in fruits and vegetables are also believed to provide antiO protection to lipid rich tissues (α -Lipoic acid (ALA)) has the ability to exist in both aqueous and lipid phase. ALA has a role in the increase in cellular GSH concentration. ALA is readily absorbed from the diet and is converted rapidly to its reduced form, dihydrolipoic acid (DHLA) (Smith et al. 2004). Both ALA and DHLA are powerful antiOs and exert their effects by scavenging free radicals, metal ion chelation and antiO recycling. DHLA as an antiO is stronger than ALA and can act synergistically with other antiOs such as glutathione, ascorbate and tocopherol (Jacob 1995; Navari-Izzo et al. 2002).

2.4. AntiO classification based on life span

Melatonin, once oxidized, cannot be reduced to its former state because it forms several stable products upon reacting with FRs-Os. Therefore, it has been referred to as a terminal (or suicidal) antiO (Tan et al. 2000). Ascorbic acid and tocopherol have long life spans. Ascorbic acid is able to regenerate tocopherol (Valko et al. 2005, 2007; Droge 2002; Davies 1995; Tan 2000). Melatonin reduces oxidative damage caused by ischemia in heart, liver, and kidney (Sewerynek et al. 1995; Reiter and Tan 2003; Sewerynek et al. 1996; Okatani et al. 2003; Sener et al. 2002).

3. Regulation and expression

Different studies proved that the endo-antiO system could respond to the presence of end/exo source of FRs-Os. Different individuals show different response based on their age, gender, genetics, and structural complexity of the organ, health condition, and nutritional status as well as different tissue specific responses. Human studies indicate that environmentally induced FRs elicit antiO responses in different tissues, which should have some link to the tissue gene specificity. The role of antiO proteins in their responses to FRs-Os was investigated using knockout animals (Kensler et al. 2007; Kobayashi and Yamamoto 2006; Itoh 2004). Various factors including cell stress lead to an increase in the expression levels of the antiO enzyme in tissues; however, coordinated expression does not necessarily occur in response to stimuli. As an example, NF-E2-related factor 2 (Nrf2) induces the expression of endo-antiOs enzymes. Nrf2 belongs to a subset of basic leucine-zipper (bZip) genes sharing a conserved structural domain, called the CNC domain. In mammals this CNC family is composed of four closely related proteins, p45-NFE2, Nrf1, Nrf2, and Nrf3, which function as heterodimeric transcription factors by pairing with other bZip proteins, including small Maf proteins (Kensler et al. 2007; Kobayashi and Yamamoto 2006; Itoh 2004).

Nrf2-antiO responsive element (ARE) activation induces the production of a battery of endogenous enzymes, such as SODs, CAT, GPx's, Prx's, NQOs and HOs (Harris 1992; Jaiswal 1994; Wasserman and Fahl 1997).

Transcriptional regulation appears to be the most common form of regulation over antiO enzyme expression (McCord 1976; Rhee et al. 2005; McCord and Edeas 2005; Mills 1957; Kim et al. 2007; Li and Jaiswal 1992; Long and Jaiswal 2000; Iskander et al. 2006; Jaiswal 2000; Ishii et al. 2000; Wagener et al. 2003; McCord 1976; Maier and Chan 2002; Harris 1992; Nanji et al. 1995; Wilson and Johnson 2000).

4. Medicinal food

Humans are one of the few vertebrates that cannot Synthesise vitamin C because the gene for the flavoenzyme, L-gulonolactone oxidase is not transcribed and is highly mutated (Chatterjee 1973).

The importance of foods which have high amounts of active compounds such as those used in modern medicine is often underestimated. There is an increasing need for more knowledge about the biological activities and/or chemical constituents of that nutraceutical food, especially for the discovery of new therapeutic drugs and novel chemical structures. In addition to this, nutraceutical food serves as an alternative medicine. Scientists are intensively investigating different plants for active compounds. Dubick (1986) reported that the medical use of herbs is deeply rooted in human history and folklore. He describes ginseng and garlic, which are plants having active compounds that are widely used in traditional medicine and food. Ginseng and garlic are well known for the existence of large amounts of antiOs (Dubick 1986; Cumming et al. 2000; Wesnes et al. 2000). Masella et al. (2004) reported that extra virgin olive oil biophenols inhibit cell-mediated oxidation of low-density lipoprotein (LDL) by increasing the mRNA transcription of glutathione-related enzymes

Awney et al. (1997) report a study on the effect of 12 different plant extracts on the aryl hydrocarbon hydroxylase (AHH) activity and H₂O₂ production by lindane induced mice hepatic microsomes. The different plant crude extracts which contain different antiOs show significant H₂O₂ and AHH inhibition, especially edible ones including *Sesamum indicum*, *Nigella sativa*, *Cinnamomum zeylanicum*, *Ficus carica*, *Curcuma longa* and *Zin-*

giber officinale (Awney et al. 1997). The same extracts are able to inhibit tumor formations using *Agrobacterium tumefaciens in vivo* potato disk antitumor bioassay which have a significant correlation to the 3PS (*in vivo*, murine leukemia) (Awney et al. 1997; Amara et al. 2008; Galsky and Wilsey 1980; 1981 a,b).

Diets containing antiOs are usually derived from vegetables and fruits including grape seeds, isothiocyanate from cruciferous vegetables, catechins from green tea, and curcumin from turmeric (Awney et al. 1997; Keum et al. 2004; Surh et al. 2005; Son et al. 2008).

Plant antiOs activate the Nrf2-mediated adaptive response (Surh et al. 2005; Son et al. 2008; Surh 2003; Kohle and Bock 2006). Studies show that consumption of fruits and vegetables containing antiOs provides protection from cancer (Surh 2003). An oversupply of antiOs results in a reducing intracellular environment, which keeps more Keap1 molecules in the reduced configuration. With less oxidized Keap1 molecules present, ubiquitination and degradation of Nrf2 increases. A decline in Nrf2 would lead to impairing the endo-antiOs (Kohle and Bock 2006; Bjelakovic et al. 2004; Bairati et al. 2006; Hercberg et al. 2007; Omenn 2007; Plummer et al. 2007; Lin et al. 2009). Taking low-dose prooxidants should be beneficial (Calabrese 2001). Low-doses of isothiocyanate and its related antiOs may activate the Nrf2 antiO system, providing protection (Calabrese 2001).

While "protection is usually better than treatment" is a common idiom and a correct scientific fact, there is a general developing theory among scientists concerning protection against and treatment for diseases using edible plants, food, or nutraceuticals. Herpeicious plants could have unique types of active constituents, but usually there is a risk of the presence of detectable or hidden harmful compounds (Awney et al. 1997). In contrast, edible plants prove historically that they have nearly no side effects (Amara et al. 2008; Galsky and Wilsey 1980; Galsky et al. 1981 a,b).

Active combination between the different types of nutraceutical foods in most cases leads to better treatment. To maintain health, a particular food diet should be used for each age/situation, especially if there is an expectation for the presence of particular or inherited diseases, or if there is an expected interaction between food and our DNA. In the case of nutraceutical foods, their composition of antiOs should be calculated (Valko et al. 2007; Droge 2002; Valko et al. 2005; Davies 1995).

5. FRs-Os/antiO/food interaction

There is convincing evidence that genetics have a crucial role in impairing the balance between AntiO/FRs-O systems. The genetic differences could affect the exo/endo/existing antiO and FRs-O systems and their interaction with nutrients and active compounds in food including mutations, impaired genes, methylated genes, single nucleotide polymorphism (SNP) and damaged genes. Homozygotic twins having identical alleles at corresponding chromosomal loci are considered genetically identical. The genetic differences will be clearer between individuals other than twins. However many exo/endo factors are not inherited and could affect our genes, such as the environment, food, and imprinting processes during fetal development which can cause methylation. Other differences across entire regions of an individual's genome result from various nutrient imbalances and deficiencies. Genetic variation between individuals is the rule not the exception (Marklund 1982; Rhee et al. 2005; McCord and Edeas 2005; Mills 1957; Kim et al. 2007; Li and Jaiswal 1992; Long and Jaiswal 2000; Iskander et al. 2006; Jaiswal 2000; Ishii et al. 2000; Wagener et al. 2003; McCord 1976; Maier and Chan 2002).

Minor changes in our genetic material could cause dangerous diseases. In most cases those changes are static. There are different mechanisms which can cause a genetic change in a person who is born normal. Cancer is an example of the interaction between food (which is either an inducer or a protector) and the genetic materials (Valko et al. 2004; Droge 2002; Chui and Carol 2008). "Cancer is a detectable internal modification that happens in our gene(s), However many other forms are undetectable or neglectable"

6. Our genes are susceptible

Is there is a real interaction between food and our DNA? In other words, what influence has food on our DNA? There is increasing evidence that nutrients have a direct effect on our genomic DNA, especially during the infant stage. To simplify this, what could be the affect of nutrients on our DNA. Examples from lower eukaryotes and prokaryotes should be considered. Studies on the human genome are rather complicated and in many cases discontinuous in regard to many aspects including time, environment, lifestyle, nutrients sources, tradition, etc. (Valko et al. 2006, 2007; Droge 2002; Davies 1995).

6.1. Learning from prokaryotes

Perhaps one interesting example of prokaryotes is *E. coli* cell XL1-Red (endA1, gyrA96, thi-1, hsdR17, supE44, relA1, lac mutD5, mutS, mutT, Tn10(Tc^r)) Stratagene® (Bachmann 1987; Glickman and Radman 1980).

Strain XL1-Red is an *E. coli* strain deficient in three of the primary DNA repair pathways. They are: *mutS* (error-prone mismatch repair), *mutD* (deficient in 3'-to 5'- exonuclease of DNA polymerase III) and *mutT* (unable to hydrolyse 8-oxodGTP). This strain grows extremely slowly in rich media (e.g. LB media), having a doubling time of ~90-120 min (Glickman and Radman 1980; Scheuermann et al. 1983).

This deficiency enables an irreversible DNA mistake causing mutations. If a specific gene comes under stress during replication or during the processes leading to a particular protein in absence of the repair mechanism, these conditions will lead to mutations. As an example Amara et al. (2002) established *in vivo* a random mutagenesis protocol for mutant screening, *phaC_{Ap}* synthase was the target gene.

Why does normal *E. coli* cultivated under the same conditions not induce such mutations? Something is different. In our case, we know the main difference between normal *E. coli* and *E. coli* XL1 Red. *E. coli* XL1 Red is a sick cell with impaired DNA repairing genes, which enable mutagenesis (Glickman and Radman 1980; Scheuermann et al. 1983; Amara et al. 2002).

Human cells under similar conditions could give the same results. When there is a deficiency, in particular essential genetic materials such as DNA repairing genes or those genes responsible for expressing essential enzymes including antiO enzymes, the problem will be exponential. The nutrients used for cultivating *E. coli* XL1 Red will start its growth, replication and the DNA mutagenesis process. The nutrients in our food could do the same job and will follow the same steps. If we have a problem in one or more of the essential genes, which could be in interaction or under the influence of a particular nutrient, this gene will be under pressure. When an external or a mutagenic factor such as FRs-O exists, more damage to the target gene(s) could occur.

When AntiOs are preventing or decreasing the amount of FOs-Os in our bodies, they are doing a vital activity (Albanes 2009; Bandera et al. 2009; Block 2009 a,b; Chen et al. 2009).

Eliminating the stress source(s) will give better results for healthy individuals, and this will always be best for those who have genetic problems. On other hand, normal *E. coli* and *E. coli* XL1-Red are examples of the genetic polymorphism, which is a real fact in the bio-system especially those that reproduce sexually or have inherited disease(s).

6.2. A real example

DNA damage and base change can affect gene transcription, mRNA stability, or protein function (McCann 1997; Dringen et al. 2005; Cha et al. 2004).

Detrimental amino acid changes and altered gene regulation have been the focus of the most attention so far. Genetic variation in oxidative stress-related genes could reveal the degree of variability that is tolerated in healthy life (e.g., acatalasemia). Furthermore, the effect of antiOs on expression levels needs to be determined. In the eukaryotic system, the results will be of course more complicated. Cancer is a clear example of above normal DNA/Protein interactions. Different proteins are working as antiOs to control the FRs-Os from damaging cell macromolecules. Among the three isoforms of SOD (SOD1, SOD2 and SOD3), SOD2 polymorphisms have largely been implicated with cancer risk. The most commonly studied polymorphism of Mn-SOD is Val 16 Ala on mitochondrial target sequence. Other examples like Pro 197 Leu SNP of the GPx1 (GPx) gene have been associated with FRs-O related diseases (Ambrosone et al. 1999; Grasbon-Frodl et al. 1999; Hu and Diamond 2003). CAT SNP is responsible for Japanese acatalasemia (Hirono et al. 1995). The CAT C-262T SNP is the most widely studied to date; it has been associated with a decrease in CAT activity and an increased or decreased risk of developing FRs-Os (Nadif et al. 2005). Other SNPs located within this region include G-844A, which has been associated with hypertension (Jiang et al. 2001; Gavalas et al. 2006).

These SNP regions can influence either nutritional status or the renutrition process because renutrition becomes progressively more difficult with age (Nadif et al. 2005).

The success in using the new molecular biology tools such as genomics, proteomics and metabolomics (Young 2002; Fay and German 2008; Fiehn 2001) will map the interactions between genes, nutrients and environment and in this way will highlight their downstream effects on human health (Young 2002; Fabre et al. 2008).

6.3. Thank God our erythrocytes contain no DNA!

Erythrocytes transport large amounts of oxygen over their lifespan resulting in oxidative stress. Red cells have potent antiO protection consisting of enzymatic and nonenzymatic pathways that modify highly active FRs-Os into substantially less reactive intermediates (Çimen 2008).

DNA and RNA are the major sensitive macromolecules in the cell. The absence of DNA in our red blood cells enables maximum expected protection against FRs-Os. Three major hereditary conditions in the erythrocytes can cause major cell disorders including sickle cell anemia (SCA), thalassemia, and glucose-6-phosphate dehydrogenase (G6PD) deficiency (Comporti et al. 2002; Scott et al. 1993; Kakhlon and Cabantchik 2002; Scott et al. 1991; Kirkman and Gaetani 2007). However, increased oxidative damage to red cells is commonly associated with all three disorders and have been attributed to the accelerated red cell destruction in all cases. Food which contains antiOs is recommended for general health improvement for those patients (Valko et al. 2005, 2007; Droge 2002; Davies 1995).

7. Nutrient personalization

Currently there is increased interest in nutrient personalization. Do the differences between humans in response to their diet components enable the building of a genetically based diet? The increasing data from genomic, proteomic and transcriptome research prove this argument. Research on gender revealed that there are different effects of antioxidants on cancer risk (Higdon and Frei 2005; Waters et al. 2004). Providing nutritional foods for helping individuals recovering from diseases is a clear opportunity to develop parallel approaches along with personalized medicine. The different observations derived from different ethnic groups indicate that humans have long known about the interaction between food and health. Scurvy is an example. Linking the genome, proteome, and transcriptome with nutrients and with each other will match our aim of establishing the nutrient personalization science. Building database(s) about genetic polymorphisms and their interaction with food aiming to solve the existing gene/food based problems or protecting against expected diseases through supplying specific nutrient(s) is in demand (Marklund 1982; Rhee et al. 2005; McCord and Edeas 2005; Mills 1957; Kim 2007; Li and Jaiswal 1992; Long and Jaiswal 2000; Iskander et al. 2006; Jaiswal 2000; Ishii et al. 2000; Wagener et al. 2003; McCord 1976; Maier and Chan 2002).

8. Conclusion

FRs-Os are not always enemies but are an essential part of our biological system. They are involved in many vital and positive processes. They are controlled with the aid of our endo-antiO system as well as the exo-antiOs, which we gain during our normal daily diets. The antiOs are groups of compounds produced mainly by the biological system and are used to remove / control the FRs-Os. They can do this individually or work as a network. Because they are produced by the biological system, most of them have references in our genetic materials.

AntiOs can be classified according to their source, nature, solubility, life span and their mode of action. In special cases, our existing antiOs do not satisfy our demand and cannot remove the excess of FRs-Os, which are generated from different exo and indo sources. Therefore, we need external sources of antiOs, preferably gained from our diet. Foods that have an extra amount of antiOs usually are called nutraceutical foods. Nutraceutical foods supply us with adequate amounts of antiOs. They can, in most cases, reduce the amount of FRs-Os and satisfy our bodies' demands.

The study of individuals and population genetic polymorphisms will be a challenging task for scientists because they have to optimize their knowledge about the role of genomes and proteomes in our health status. Those who have inheritable diseases need to gain exo-antiOs to recover from deficiencies. The overcontinuous expression and induction of FRs-O or antiO systems could be a source of self-damage. Exogenous sources of antiOs will not only satisfy our need and cover any kind of deficiency but also will enable the cells to be more relaxed and to direct their power to other vital processes. AntiOs should be used wisely because free radicals also have essential roles in our life. FRs-O levels should always be under control to promote our best health. Personalizing food will show an increasing interest in *ab initio* protection based on the genetic polymorphisms between different individuals. We are not equal genetically. A case-by-case healthy nutrition regime is a key factor in our healthy future.

"With hundred of friends, one has problems! Reducing the number will be better! Treat FRs-Os as friends and not as enemies, but reduce their amounts." The author.

References

- Albanes D (2009) Vitamin supplements and cancer prevention: where do randomized controlled trials stand? *J Natl Cancer Inst* 101: 2–4.
- Amara AA, El-masry MH, Bagdady HH (2008) Plant crude extracts could be the solution: extracts showing *in vivo* antitumorigenic activity. *Pak J Pharm Sci* 21: 159–171.
- Amara AA, Steinbüchel A, Rehm BHA (2002) *In vivo* evolution of the *Aeromonas punctata* polyhydroxyalkanoate (PHA) synthase: Isolation and characterization of modified PHA synthases with enhanced activity. *Appl Microbiol Biotechnol* 59: 477–482.
- Ambrosone C, Freudenheim J, Thompson PA, Bowman E, Vena JE, Marshall JR, Graham S, Laughlin R, Nemoto T, Shields PG (1999) Manganese super-oxide dismutase (MnSOD) genetic polymorphisms, dietary antioxidants, and risk of breast cancer. *Cancer Res* 59: 602–606.
- Ames N, Shigenaga MK, Hagan TM (1993) Oxidants, antioxidants and the degenerative diseases of aging. *Proc Natl Acad Sci USA* 90: 7915–7922.
- Awney HA, Amara AA, El-Masry MH, Bagdady HH (1997) Effect of 12 plant extract on hepatic microsomal pento(a) pyrene hydroxylation and hydrogen peroxide production in mice treated with lendane. *J Environm Nutrition Interact* V(1):P 129–142.
- Bachmann BJ. (1987) Linkage map of *Escherichia coli* K12. In: F. C. Neidhardt (ed), *Escherichia coli* and *Salmonella typhimurium*, second edition. Vol. 2, ASM, Washington DC 807–876.
- Bahorun T, Soobrattee MA, Luximon-Ramma V, Aruoma OI (2006) Free radicals and antioxidant in cardiovascular health and disease. *Internet J Med Update* 1: 1–17.
- Bairati I, Meyer F, Jobin E, Gélinais M, Fortin A, Nabid A, Brochet F, Têtu B (2006) Antioxidant vitamins supplementation and mortality: a randomized trial in head and neck cancer patients. *Int J Cancer* 119: 2221–2224.
- Bandera EVG, Dina M, Moore DF, McCullough ML, Kushi LH (2009) Antioxidant vitamins and the risk of endometrial cancer: a dose-response meta-analysis. *Cancer Causes Control* 20: 699–711.
- Beckman KB, Ames BN (1998) The free radical theory of ageing matures. *Physiol Rev* 78: 547–581.
- Benzie IFF (2003) Evolution of dietary antioxidants *Comparat Biochem Physi* 136: 113–126.
- Bjelakovic G, Nikolova D, Simonetti RG, Gluud C (2004) Antioxidant supplements for prevention of gastrointestinal cancers: a systematic review and meta-analysis. *Lancet* 364: 1219–1228.
- Block KI (2009a) Antioxidants: Selected out? *Integr Cancer Ther* 8: 5–8.
- Block KI (2009b) Antioxidant supplementation and cancer prevention. *JAMA* 301:1877–1878.
- Bunn HF (1991) Pathophysiology of the anemias. In: Wilson JD, Braunwald E, Isselbacher KJ (eds) *Harrison's Principle of Internal Medicine*. New York: McGraw-Hill Inc 1514–8.
- Calabrese EJ (2001) Overcompensation stimulation: a mechanism for hormetic effects. *Crit Rev Toxicol*. 31: 425–470.
- Cha MK, Hong SK, Lee DS, Kim IH (2004) *Vibrio cholerae* thiol peroxidase-glutaredoxin fusion is a 2-Cys TSA/AhpC subfamily acting as a lipid hydroperoxide reductase. *J Biol Chem* 279: 11035–11041.
- Chatterjee IB (1973) Evolution and the biosynthesis of ascorbic acid. *Science* 182: 1271–1272.
- Chaudière J, Ferrari-Iliou R (1999) Intracellular antioxidants: from chemical to biochemical mechanisms. *Food Chem Toxicol* 37 (9–10): 949–962.
- Chen CSS, Jeremy A, Wells PG (2009) Reduced tumorigenesis in p53 knockout mice exposed in utero to low-dose vitamin E. *Cancer* 115: 1563–1575.
- Chui MHG, Carol E. (2008) Antioxidant vitamins reduce acute meal-induced memory deficits in adults with type 2 diabetes. *Nutr Res* 28: 423–429.
- Çimen MYB (2008) Free radical metabolism in human erythrocytes. *Clin Chim Acta* 390: 1–11.
- Comporti M, Signorini C, Buonocore G, Ciccoli L (2002) Iron release, oxidative stress and erythrocyte ageing. *Free Radic Biol Med* 32:568–576.
- Cumming RG, Mitchell P, Wayne S (2000) Diet and cataract. The blue mountains eye study. *Ophthalmology* 107: 450–456.
- Cutler RG (1984) Urate and ascorbate: their possible roles as antioxidants in determining longevity of mammalian species. *Arch Gerontol Geriat* 3: 321–348.
- Davies K (1995) Oxidative stress: the paradox of aerobic life. *Biochem Soc Symp* 61: 1–31.
- Davis JMM, Angela E, Carmichael, MD, Davis B (2009) Quercetin increases brain and muscle mitochondrial biogenesis and exercise tolerance. *Am J Physiol Regul Integr Comp Physiol* 296: 1071–1077.

- De Luca LM, Ross SA (2009) Beta-caroten increases lung cancer incidence in cigarette smoker. *Nutrition Rev* 54: 178–180.
- de Oliveira MRO, Soares MW, Behr GA, Hoffa MLM, da Rocha RF, Moreira JCF (2009) Evaluation of the effects of vitamin A supplementation on adult rat substantia nigra and striatum redox and bioenergetic states: mitochondrial impairment, increased 3-nitrotyrosine and alpha-synuclein, but decreased D2 receptor contents. *Prog Neuropsychopharmacol Biol Psychiatry* 33: 353–362.
- Deneke SM, Fanburg BL (1989) Regulation of cellular glutathione. *Am J Physiol* 257: L163–L173.
- Dringen R, Pawlowski PG, Hirrlinger J (2005) Peroxide detoxification by brain cells. *J Neurosci Res* 79: 157–165.
- Droge W (2002) Free radicals in the physiological control of cell function. *Review. Physiol Rev* 82: 47–95.
- Dubick AM (1986) Historical perspectives on the use of herbal preparations to promote health. *J Natr* 116: 1348–1354.
- Fabre EE, Raynaud-Simon A, Golmard J-L, Hebert M, Dulcire X, Succari M, Myara J, Durand D, Nivet-Antoine V (2008) Gene polymorphisms of oxidative stress enzymes: prediction of elderly renutrition. *Am J Clin Nutr* 87: 1504–1512.
- Fay LB, German BJ (2008) Personalizing foods: is genotype necessary? *Current Opin Biotechnol* 19: 121–128.
- Fiehn O (2001) Combining genomics, metabolome analysis, and biochemical modelling to understand metabolic networks. *Comp Func Genom* 2: 155–168.
- Galsky AG, Kozimor R, Piotrowski D, Powell RG (1981a) The crown-gall potato disk bioassay as a primary screen for compounds with antitumor activity. *J Natl Cancer Inst* 67: 689–692.
- Galsky AG, Scheppeler JA, Cranford MS (1981b) Crown-Gall tumors possess tumor-specific antigenic sites on their cell walls. *Plant Physiol* 67: 1195–1197.
- Galsky AG, Wilsey JP (1980) Crown Gall Tumor Disc Bioassay: A possible aid in the detection of compounds with antitumor activity. *Plant Physiol* 65: 184–185.
- Gann PH (2009) Randomized trials of antioxidant supplementation for cancer prevention: first bias, now chance—next, cause. *JAMA* 301: 102–103.
- Gavalas N, Akhtar S, Gawkrödger DJ, Watson PF, Weetman AP, Kemp EH (2006) Analysis of allelic variants in the catalase gene in patients with the skin depigmenting disorder vitiligo. *BBRC* 345: 1586–1591.
- Genestra M (2007) Oxy radicals, redox-sensitive signalling cascades and antiO₂. *Review. Cell Signal* 19: 1807–1819.
- Ghafourifar P, Cadenas E (2005) Mitochondrial nitric oxide synthase. *Trends Pharmacol Sci* 26: 190–195.
- Glickman, BW, Radman M (1980) *Escherichia coli* mutator mutants deficient in methylation-instructed DNA Mismatch correction. *Proc Natl Acad Sci USA* 77: 1063–1067.
- Grasbon-Frodl E, Kosel S, Riess O, Muller U, Mehraein P, Graeber MB (1999) Analysis of mitochondrial targeting sequence and coding region polymorphisms of the manganese superoxide dismutase gene in German Parkinson disease patients. *Biochem Biophys Res Commun* 255: 749–752.
- Gulewitsch W, Amiradzibi S (1900) über das Carnosin, eine neue organische Base des Fleischextraktes. *Ber Deutsh Chem Ges* 33: 1902–1903.
- Halliwell B (2007) Biochemistry of oxidative stress. *Biochem Soc Trans* 35: 1147–1150.
- Halliwell B, Gutteridge JMC (1999) *Free Radicals in Biology and Medicine*. third ed. Clarendon Press, Oxford, UK.
- Hammadeh ME, Al Hasani S, Rosenbaum P, Schmidt W, Fischer HC (2008) Reactive oxygen species, total antioxidant concentration of seminal plasma and their effect on sperm parameters and outcome of IVF/ICSI patients. *Arch Gynecol Obstet* 277: 515–526.
- Harman D (1956) Aging - A theory based on free-radical and radiation chemistry. *J Gerontol* 11: 298–300.
- Harris ED (1992) Regulation of antioxidant enzymes. *FASEB J* 6: 2675–2683.
- Hercberg S, Ezzedine K, Guinot C, Preziosi P, Galan P, Bertrais S, Estaquio C, Briançon S, Favier A, Latreille J, Malvy D (2007) Antioxidant supplementation increases the risk of skin cancers in women but not in men. *J Nutr* 137: 2098–2105.
- Higdon JV, Frei B (2005) Is there a gender difference in the effect of antioxidants on cancer risk? *Brit J Nutr* 94: 139–140.
- Hirono A, Sasaya-Hamada F, Kanno H, Fujii H, Yoshida T, Miwa S (1995) Novel human catalase mutation (358 T→del) causing Japanese-type acatalasemia. *Blood Cells Mol Dis* 21: 232–234.
- Hofmann B, Hecht HJ, Flohe L (2002) Peroxiredoxins. *Biol Chem* 383: 347–364.
- Hu Y, Diamond A (2003) Role of glutathione peroxidase 1 in breast cancer: loss of heterozygosity and allelic differences in the response to selenium. *Cancer Res* 63: 3347–3351.
- Ishii T, Itoh K, Takahashi S, Sato H, Yanagawa T, Katoh Y, Bannai S, Yamamoto M (2000) Transcription factor Nrf2 coordinately regulates a group of oxidative stress-inducible genes in macrophages. *J Biol Chem* 275: 16023–16029.
- Iskander K, Li J, Han S, Zheng B, Jaiswal AK (2006) NQO1 and NQO2 regulation of humoral immunity and autoimmunity. *J Biol Chem* 281: 30917–30924.
- Itoh K, Tong KI, Yamamoto M (2004) Molecular mechanism activating Nrf2–Keap1 pathway in regulation of adaptive response to electrophiles. *Free Radic Biol Med* 36: 1208–1213.
- Jacob RA (1995) The integrated antioxidant system. *Nutrition Res* 15(5):755–771.
- Jaiswal AK (1994) Antioxidant response element. *Biochem Pharmacol* 48: 439–444.
- Jaiswal AK (2000) Regulation of genes encoding NAD(P)H:quinone oxidoreductases. *Free Radic Biol Med* 29: 254–262.
- Jiang Z, Akey JM, Shi J, Xiong M, Wang Y, Shen Y, Xu X, Chen H, Wu H, Xiao J, Lu D, Huang W, Jin L (2001) A polymorphism in the promoter region of catalase is associated with blood pressure levels. *Hum Genet* 109: 95–98.
- Jindal SK (2008) Does vitamin E supplementation help in COPD? *Indian J Med Res* 128: 686–687.
- Johnson F, Giulivi C (2005) Superoxide dismutases and their impact upon human health. *Mol Aspects Med* 26: 340–352.
- Joyner MJ (2009) Keeping the juices flowing with age: vitamin C and exercise hyperaemia. *J Physiol* 587: 2423.
- Kakhlon O, Cabantchik ZI (2002) The labile iron pool: Characterization, measurement, and participation in cellular processes. *Free Radic Biol Med* 33: 1037–1046.
- Kensler TW, Wakabayashi N, Biswal S (2007) Cell survival responses to environmental stresses via the Keap1–Nrf2–ARE pathway. *Annu Rev Pharmacol Toxicol* 47: 89–116.
- Keum YS, Jeong WS, Kong AN (2004) Chemoprevention by isothiocyanates and their underlying molecular signaling mechanisms. *Mutat Res* 555: 191–202.
- Kim YJ, Ahn JY, Liang P (2007) Human *prx1* gene is a target of Nrf2 and is up-regulated by hypoxia/reoxygenation: implication to tumor biology. *Cancer Res* 67: 546–554.
- Kirkman HN, Gaetani GF (2007) Mammalian catalase: a venerable enzyme with new mysteries. *Trends Biochem Sci* 32: 44–50.
- Kobayashi M, Yamamoto M (2006) Nrf2–Keap1 regulation of cellular defense mechanisms against electrophiles and reactive oxygen species. *Adv. Enzyme Regul* 46: 113–140.
- Kohle C, Bock KW (2006) Activation of coupled Ah receptor and Nrf2 gene batteries by dietary phytochemicals in relation to chemoprevention. *Biochem Pharmacol* 72: 795–805.
- Lauterburg BH, Adams JD, Mitchell JR (1984) Hepatic glutathione homeostasis in the rat: efflux accounts for glutathione turnover. *Hepatology* 4: 586–590.
- Li Y, Jaiswal AK (1992) Regulation of human NAD(P)H:quinone oxidoreductase gene: role of AP1 binding site contained within human antioxidant response element. *J Biol Chem* 267: 15097–15104.
- Liddell JR, Dringen R, Crack PJ, Robinson SR (2006) Glutathione peroxidase 1 and a high cellular glutathione concentration are essential for effective organic hydroperoxide detoxification in astrocytes. *Glia* 54: 873–879.
- Liddell JR, Hoepken HH, Crack PJ, Robinson SR, Dringen R (2006) Glutathione peroxidase 1 and glutathione are required to protect mouse astrocytes from iron-mediated hydrogen peroxide toxicity. *J Neurosci Res* 84: 578–586.
- Lin J, Cook NR, Albert C, Zaharris E, Gaziano JM, Van Denburgh M, Buring JE, Manson JE (2009) Vitamins C and E and beta carotene supplementation and cancer risk: a randomized controlled trial *J Natl Cancer Inst* 101: 14–23.
- Long DJ, Jaiswal AK (2000) NRH:quinone oxidoreductase2 (NQO2). *Chem Biol Interact* 129: 99–112.
- Maier CM, Chan PH (2002) Role of superoxide dismutases in oxidative damage and neurodegenerative disorders. *Neuroscientist* 8: 323–334.
- Marklund SL (1982) Human copper-containing superoxide dismutase of high molecular weight. *Proc Natl Acad Sci USA* 79: 7634–7638.

- Masella R, Vari R, D'Archivio M, Benedetto RD, Matarrese P, Malorni W, Scazzocchio B, Giovannini C (2004) Extra virgin olive oil biophenols inhibit cell-mediated oxidation of LDL by increasing the mRNA transcription of glutathione-related enzymes. *Nutr* 134: 785–791.
- McCann SM (1997) The nitric oxide hypothesis of brain aging. *Exp Gerontol* 32: 431–440.
- McCord JM (1976) Iron- and manganese-containing superoxide dismutases: structure, distribution, and evolutionary relationships. *Adv Exp Med Biol* 74: 540–550.
- McCord JM, Edeas MA (2005) SOD, oxidative stress and human pathologies: a brief history and a future vision. *Biomed Pharmacother* 59: 139–142.
- McCord JM, Fridovich I (1969) Superoxide dismutase an enzymic function for erythrocyte Hemocuprein. *J Biol Chem* 244: 6049–6055.
- McCord JM, Fridovich I (1969) Superoxide dismutase: an enzymic function for erythrocyte (hemocuprein). *J Biol Chem* 244: 6049–6055.
- Mills GC (1957) Hemoglobin catabolism. I. Glutathione peroxidase, an erythrocyte enzyme which protects hemoglobin from oxidative breakdown. *J Biol Chem* 229: 189–197.
- Nadif R, Mintz M, Jedlicka A, Bertrand JP, Kleeburger SR, Kauffmann F (2005) Association of CAT polymorphisms with catalase activity and exposure to environmental oxidative stimuli. *Free Radic Res* 39: 1345–1350.
- Nakabeppu Y, Sakumi K, Sakamoto K, Tsuchimoto D, Tsuzuki T, Nakatsu Y (2006) Mutagenesis and carcinogenesis caused by the oxidation of nucleic acids. *Biol Chem* 387: 373–379.
- Nanji AA, Griniuvienė B, Sarzadeh SMH, Levitsky S, McCully JD (1995) Effect of type of dietary fat and ethanol on antioxidant enzyme mRNA induction in rat liver. *J Lipid Res* 36: 736–744.
- Navari-Izzo F, Quartacci MF, Sgherri C (2002) Lipoic acid: a unique antioxidant in the detoxification of activated oxygen species. *Plant Physiol Biochem* 40: 463–470.
- Okatani Y, Wakatsuki A, Reiter RJ, Enzan H, Miyahara Y (2003) Protective effect of melatonin against mitochondrial injury induced by ischemia and reperfusion of rat liver. *Eur J Pharmacol* 469: 145–152.
- Omenn GS (2007) Chemoprevention of lung cancers: lessons from CARET, the beta-carotene and retinol efficacy trial, and prospects for the future. *Eur J Cancer Prev* 16: 184–191.
- Ortega RM (2006) Importance of functional foods in the Mediterranean diet. *Public Health Nutr* 9 (8A): 1136–1140.
- Pacher P, Beckman JS, Liaudet L (2007) Nitric oxide and peroxynitrite in health and disease. *Physiol Rev* 87: 315–424.
- Parthasarathy S, Santanam N, Ramachandran S, Meilhac O (1999) Oxidants and antioxidant in atherosclerosis: an appraisal. *J Lipid Res* 40: 2143–2157.
- Platz EA (2009) Selenium, genetic variation, and prostate cancer risk: epidemiology reflects back on selenium and vitamin E cancer prevention trial. *J Clin Oncol* 27: 3569–3572.
- Plummer M, Vivas J, Lopez G, Bravo JC, Peraza S, Carillo E, Cano E, Castro D, Andrade O, Sánchez V, Garcia R, Buiatti E, Aebischer C, Franceschi S, Oliver W, Muñoz N (2007) Chemoprevention of precancerous gastric lesions with antioxidant vitamin supplementation: a randomized trial in a high-risk population. *J Natl Cancer Inst* 99: 137–146.
- Raha S, Robinson B (2000) Mitochondria, oxygen free radicals, disease and aging. *Trends Biochem Sci* 25: 502–508.
- Rayman MPC, Gerald F, Waters, DJ (2009) Selenium and vitamin E supplementation for cancer prevention. *JAMA* 301: 1876–1877.
- Reiter RJ, Tan DX (2003) Melatonin: a novel protective agent against oxidative injury of the ischemic/reperfused heart. *Cardiovasc Res* 58: 10–19.
- Rhee SG, Chae HZ, Kim K (2005) Peroxiredoxins: a historical overview and speculative preview of novel mechanisms and emerging concepts in cell signaling. *Free Radic Biol Med* 38: 1543–1552.
- Sachdev S, Davies KJ (2008) Production, detection, and adaptive responses to free radicals in exercise. *Free Radic Biol Med* 44: 215–223.
- Scheuermann R, Tam S, Burgers PM, Lu C, Echols H (1983) Identification of the epsilon-subunit of Escherichia coli DNA polymerase III holoenzyme as the dnaQ gene product: a fidelity subunit for DNA replication. *Proc Natl Acad Sci USA* 80: 7085–7089.
- Schrauzer GN (2009) RE: Lessons from the selenium and vitamin E cancer prevention trial. *Crit Rev Biotechnol* 29: 81.
- Schrauzer GNS, Peter F (2009) Selenium in human and animal nutrition: resolved and unresolved issues. A partly historical treatise in commemoration of the fiftieth anniversary of the discovery of the biological essentiality of selenium, dedicated to the memory of Klaus Schwarz (1914–1978) on the occasion of the thirtieth anniversary of his death. *Crit Rev Biotechnol* 29: 2–9.
- Scott MD, van den Berg JJ, Repka T, Rouser-Fessard P, Hebbel RP, Beuzard Y, Lubin BH (1993) Effect of excess alpha-hemoglobin chains on cellular and membrane oxidation in model beta-thalassemic erythrocytes. *J Clin Invest* 91: 1706–1712.
- Scott MD, Zuo L, Lubin BH, Chiu DT (1991) NADPH, not glutathione, status modulates oxidant sensitivity in normal and glucose-6-phosphate dehydrogenase-deficient erythrocytes. *Blood* 77: 2059–2064.
- Seifried HE, Anderson DE, Fisher EI, Milner JA (2007) A review of the interaction among dietary antioxidants and reactive oxygen species. *J Nutritional Biochem* 18: 567–579.
- Sener G, Sehirlı AO, Keyer-Uysal M, Arbak S, Ersoy Y, Yegen BC (2002) The protective effect of melatonin on renal ischemia-reperfusion injury in the rat. *J Pineal Res* 32: 120–126.
- Sewerynek E, Melchiorri D, Chen L, Reiter RJ (1995) Melatonin reduces both basal and bacterial lipopolysaccharide-induced lipid peroxidation *in vitro*. *Free Rad Biol Med* 19: 903–909.
- Sewerynek E, Reiter RJ, Melchiorri D, Ortiz GG, Levinski A (1996) Oxidative damage in the liver induced by ischemia-reperfusion: protection by melatonin. *Hepato-gastroenterology* 43: 898–905.
- Shaw H-MC, Wan-Hsuan (2009) Effects of pregnenolone-16alpha-carbonitrile on vitamin E status and protein levels of antioxidant enzymes in male rats fed a vitamin E-supplemented diet. *Redox Rep* 14: 61–68.
- Siegel D, Ross D (2000) Immunodetection of NAD(P)H:quinone oxidoreductase 1 (NQO1) in human tissues. *Free Radic Biol Med* 29: 246–253.
- Sies H (1997) Oxidative stress: oxidants and antioxidants. *Exp Physiol* 82: 291–295.
- Sies H (1993) Strategies of antioxidant defense. *Eur J Biochem* 215: 213–219.
- Siow RCM, Ishii T, Sato H, Taketani S, Leake DS, Sweiry JH, Pearson JD, Bannai S, Mann GE (1995) Induction of the antioxidant stress proteins heme oxygenase-1 and MSP23 by stress agents and oxidized LDL in cultured vascular smooth muscle cells. *FEBS Lett* 368: 239–242.
- Smith AR, Shenvi SV, Widlansky M, Suh JH, Hagen TM (2004) Lipoic acid as a potential therapy for chronic diseases associated with oxidative stress. *Curr Med Chem* 11: 1135–1146.
- Son TG, Camandola S, Mattson MP (2008) Hormetic dietary phytochemicals. *Neuromolecular Med* 10: 236–246.
- Stadtman E (1992) Protein oxidation and aging. *Science* 257: 1220–1224.
- Surh YJ (2003) Cancer chemoprevention with dietary phytochemicals. *Nat Rev Cancer* 3: 768–780.
- Surh YJ, Kundu JK, Na HK, Lee JS (2005) Redox-sensitive transcription factors as prime targets for chemoprevention with anti-inflammatory and antioxidative phytochemicals. *J Nutr* 135: 2993S–3001S.
- Tan DX, Manchester LC, Reiter RJ, Qi WB, Karbownik M, Calvo JR (2000) Significance of melatonin in antioxidative defense system: reactions and products. *Biol Signals Recept* 9: 137–159.
- Traber MG, Packer L (1995) Vitamin E: Beyond antioxidant function. *Am J Clin Nutr* 62: 1501–1509.
- Valko M, Izakovic M, Mazur M, Rhodes CJ, Telser J (2004) Role of oxygen radicals in DNA damage and cancer incidence. *Mol Cell Biochem* 266: 37–56.
- Valko M, Leibfritz D, Moncol J (2007) Free radicals and antioxidants in normal physiological functions. *Int J Biochem Cell Biol* 39: 44–84.
- Valko M, Leibfritz D, Moncola J, Cronin MT, Mazur M, Telser J (2007) Free radicals and antioxidant in normal physiological functions and human disease. Review. *Int J Biochem Cell Biol* 39: 44–84.
- Valko M, Morris H, Cronin MTD (2005) Metals, toxicity and oxidative stress. *Curr Med Chem* 12: 1161–1208.
- Valko M, Rhodes CJ, Moncol J, Izakovic M, Mazur M (2006) Free radicals, metals and antioxidant in oxidative stress-induced cancer. Mini-review. *Chem Biol Interact* 160: 1–40.
- van Horsen J, Schreiber G, Bö L, Montagne L, Drukarch B, van Muiswinkel FL, de Vries HE (2006) NAD(P)H:quinone oxidoreductase 1 expression in multiple sclerosis lesions. *Free Radic Biol Med* 41: 311–317.
- van Muiswinkel FL, de Vos RA, Bol JG, Andringa G, Jansen SEN, Ross D, Siegel D, Drukarch B (2004) Expression of NAD(P)H:quinone oxidoreductase in the normal and Parkinsonian substantia nigra. *Neurobiol Aging* 25: 1253–1262.
- Vasdev SV, Gill PK, Singal PK (2006) Singa Beneficial effect of low ethanol intake on the cardiovascular system: possible biochemical mechanisms. *Vasc Health Risk Manag* 2: 263–276.
- Vertuani S, Angusti A, Manfredini S (2004) The antioxidants and pro-antioxidants network. *Curr Pharm Des* 10: 1677–1694.

- Volpe EP (1993) Blood and circulation. In: Dubuque WC (ed) *Biology and Human Concerns* Brown Publishers 253–265.
- Wagener FA, Volk HD, Willis D (2003) Different faces of the heme–heme oxygenase system in inflammation. *Pharmacol Rev* 55: 551–571.
- Wang Y (2009) Differential effects of sodium selenite and nano-Se on growth performance, tissue se distribution, and glutathione peroxidase activity of avian broiler. *Biol Trace Elem Res* 128: 184–190.
- Wasserman WW, Fahl WE (1997) Functional antioxidant responsive elements. *Proc Natl Acad Sci USA* 94: 5361–5366.
- Waters DJ, Chiang EC, Cooley DM, Morris JS (2004) Making sense of sex and supplements: differences in the anticarcinogenic effects of selenium in man and women. *Mutant Res* 551: 91–107.
- Wender DF, Thulin GE, Smith GJ, Warshaw JB (1981) Vitamin E affects lung biochemical and morphologic response to hyperoxia in the newborn rabbit. *Pediatr Res* 15:262–268.
- Wesnes KA, Ward T, McGinty A, Petrini O (2000) The memory enhancing effects of a Ginkgo biloba/Panax ginseng combination in healthy middle-aged volunteers. *Psychopharmacology* 152: 351.
- Willcox JK, Ash SL, Catignani GL (2004) Antioxidant and prevention of chronic disease. Review. *Crit Rev Food Sci Nutr* 44: 275–295.
- Wilson DO, Johnson P (2000) Exercise modulates antioxidant enzyme gene expression in rat myocardium and liver. *J Appl Physiol* 88: 1791–1796.
- Yim MB, Chae HZ, Rhee SG, Chock PB, Stadtman ER (1994) On the protective mechanism of the thiol- specific antioxidant enzyme against the oxidative damage of biomacromolecules. *J Biol Chem* 269: 1621–1626.
- Young IS, Woodside Jr (2001) Antioxidant in health and disease. *J Clin Pathol* 54: 176–186.
- Young VR (2002) Human nutrient requirements: the challenge of the post-genome era. *J Nutr* 132: 621–629.