

## Effects of Chinese herb medicine *Radix Scrophulariae* on ventricular remodeling

WEI-LIANG GU, CHANG-XUN CHEN, QI WU, JIAN LÜ, YING LIU, SHI-JIE ZHANG

Received March 4, 2010, accepted May 7, 2010

Chang-Xun Chen, Professor, Department of Pharmacology, Shanghai University of Traditional Chinese Medicine, 1200 Cai Lun Road, Shanghai 201203, P.R. of China  
cxchen6@126.com

Pharmazie 65: 770–775 (2010)

doi: 10.1691/ph.2010.0561

The effects and mechanism of the extract of *Radix Scrophulariae* (ERS), a traditional Chinese herb, on experimental ventricular remodeling in rats was studied. Rats were separated randomly into 5 groups: sham, model, captopril ( $40 \text{ mg} \cdot \text{kg}^{-1}$ ) and ERS (8,  $16 \text{ g} \cdot \text{kg}^{-1}$ ). The experimental ventricular remodeling was induced with ligating the left anterior descending branch of the coronary artery of the rats. The sham group was conducted the same procedure without ligation. After 4 weeks treatment with intragastric administration of the corresponding drugs, the left ventricular weight index (LVWI) and heart weight index (HWI) were determined. The concentrations of angiotensin II (Ang II) and hydroxyproline (Hyp) in myocardium were detected. Myocardium tissue was stained with HE and picric acid/Sirius red for cardiocyte cross-section area and collagen content measurements. Real-time RT-PCR was used to detect the gene expressions of  $\text{AT}_1\text{R}$ ,  $\text{TNF-}\alpha$  and  $\text{TGF-}\beta_1$  mRNA. ERS could significantly reduce the LVWI, HWI, decrease the content of Ang II, Hyp, diminish cardiocyte cross-section area and ameliorate collagen deposition. In addition, ERS could down regulate the gene expressions of  $\text{AT}_1\text{R}$ ,  $\text{TNF-}\alpha$  and  $\text{TGF-}\beta_1$  mRNA in myocardium. ERS has beneficial effect against ventricular remodeling. The mechanism may be related to decreasing the level of Ang II and cardiac fibrosis, modulating some gene expressions associated with cardiac hypertrophy.

### 1. Introduction

*Radix Scrophulariae*, the dried root of *Scrophularia ningpoensis* Hemsl., is a traditional Chinese herb recorded in a classic source (Shen Nong Ben Cao Jing). This means that the herb has been used for more than two thousand years. In TCM theory, *Radix Scrophulariae* has the effects of clearing away heat from the blood, nourish yin, purging fire, and relieving toxins, which increase patients' ability to keep the balance of yin and yang and protect them from invasion by internal or external pathogenic influences. Indications include: consumption of yin caused by febrile disease, deep-red tongue and polydipsia, maculae caused by virulent pathogen, constipation due to impairment of body fluid, hectic fever and phthical cough, conjunctivitis, sore throat, carbuncles, sores and other pyogenic skin infections (Jialin 2007).

Myocardial infarction (MI) is one of the leading causes of death throughout the world. Left ventricular (LV) remodeling after MI involves cardiocyte hypertrophy, chamber dilation, and interstitial fibrosis. Alterations in cardiocytes and collagen matrix lead to ventricular enlargement, contractile dysfunction and heart failure (HF). Hence, LV remodeling is regarded as a major risk factor of heart disease.

In the present study, we investigated whether long-term treatment with ERS could prevent the progression of LV remodeling after MI in rats and revealed the underline mechanism of its action to a certain extent.

### 2. Investigations and results

#### 2.1. Effects on left verticals weight index and heart weight Index

The degree of cardiac hypertrophy was assessed by the increase of left ventricular weight index (LVWI) and heart weight index (HWI). As shown in Table 1, LVWI and HWI were increased significantly in the model group compared with those in the sham-operated group ( $p < 0.01$ ). ERS treatment significantly decreased LVWI and HWI. The reference drug captopril also significantly reduced LVWI and HWI.

#### 2.2. Effects on angiotensin II and hydroxyproline


LV remodeling is associated with the renin-angiotensin-aldosterone system. As shown in Table 1, angiotensin II (Ang II) and hydroxyproline (Hyp) concentrations were significantly higher in the model than in the sham-operated rats ( $p < 0.01$ ), but they were attenuated in ERS and captopril groups ( $p < 0.01$ ).

#### 2.3. Effects on cardiocyte cross-section area

The ventricles were fixed with 10% formaldehyde, embedded in paraffin and sectioned. The sections were stained with HE and photographed under  $400 \times$  magnification, as shown in Fig. 1. HE staining showed about a 2-fold increase in the cardiocyte

**Table 1: Effect of ERS on cardiac weight indexes, myocardium Ang II and Hyp concentrations in rats with ventricular remodeling induced by coronary artery ligation ( $\bar{x} \pm s$ , n = 11)**

Group	Dose	LVWI (mg/g)	HWI (mg/g)	Ang II pg/mg-prot	Hyp ( $\mu$ g/mg-prot)
Sham	--	2.34 $\pm$ 0.14**	2.98 $\pm$ 0.17*	278.21 $\pm$ 19.94**	0.957 $\pm$ 0.168**
Model	--	2.53 $\pm$ 0.16	3.18 $\pm$ 0.23	348.34 $\pm$ 18.95	1.162 $\pm$ 0.090
Captopril	40 mg/kg	2.28 $\pm$ 0.17**	2.87 $\pm$ 0.19**	315.83 $\pm$ 23.91**	0.696 $\pm$ 0.349**
ERS-L	8 g/kg	2.44 $\pm$ 0.14	3.08 $\pm$ 0.15	319.97 $\pm$ 27.20*	0.745 $\pm$ 0.209**
ERS-H	16 g/kg	2.38 $\pm$ 0.12*	2.97 $\pm$ 0.11*	307.84 $\pm$ 25.49**	0.569 $\pm$ 0.301**

\* $P < 0.05$ , \*\* $P < 0.01$  vs model groupFig. 1: Effect of ERS on cardiocyte cross-section area in rats with ventricular remodeling induced by coronary artery ligation. A: sham-operated group; B: model group; C: captopril group; D: ERS 8 g·kg<sup>-1</sup> group; E: ERS 16 g·kg<sup>-1</sup> group

cross-section area in model rats compared with that in the sham-operated rats. Cardiocyte cross-section area was smaller in ERS and captopril treated groups than in the model group ( $p < 0.01$ ), as shown in Table 2.

## 2.4. Effects on collagen deposition

### 2.4.1. Perivascular collagen area and collagen volume fraction

The ventricular sections were stained with Sirius red in aqueous saturated picric acid for examination of perivascular and interstitial fibrosis in myocardium. As shown in Figs. 2, 3 and Table 2, the perivascular collagen area (PVCA) and collagen volume fraction (CVF) were markedly increased in the model group, compared to that in the control group. The treatment with ERS or captopril significantly decreased PVCA and CVF. ( $p < 0.01$ )

### 2.4.2. Effects on collagen distributions of type I and III

The contents of subtype I and III collagen in myocardium were determined using a microscope with polarimetric filter analysis

at the same time. As shown in Figs. 4, 5 and Table 2, collagen distributions of type I and III were significantly increased in model rats compared with that in the sham-operated rats. However, those were significantly decreased in ERS or captopril treated groups ( $p < 0.01$ ).

## 2.5. AT<sub>1</sub>R, TNF- $\alpha$ and TGF- $\beta$ 1 mRNA expression

To further evaluate the effect of ERS on ventricular remodeling, we investigated some gene expressions associated with cardiac hypertrophy such as AT<sub>1</sub>R, TNF- $\alpha$  and TGF- $\beta$ 1. As shown in Table 3, the AT<sub>1</sub>R, TNF- $\alpha$  and TGF- $\beta$ 1 mRNA expression were increased significantly in the LV of model rats compared to that of sham-operated rats. ERS remarkably reduced these gene expression levels ( $p < 0.01$ ). Captopril also significantly reduced these gene expressions ( $p < 0.01$ ).

$$\text{relative expression of mRNA} = 2^{-\Delta\text{ct}} \times 100\%$$

$$\Delta\text{ct} = \text{CT}(\text{objective gene}) - \text{CT}(\text{GAPDH})$$

**Table 2: Effects of ERS on cardiocyte cross-section area and collagen deposition in rats with ventricular remodeling induced by coronary artery ligation ( $\bar{x} \pm s$ , n = 8)**

Group	Dose	Cardiocyte cross-section area (pixel)	PVCA	CVF (%)	Collagen I (%)	Collagen III (%)
Sham	--	6340.0 $\pm$ 531.1**	4.49 $\pm$ 1.50**	2.95 $\pm$ 1.37**	0.25 $\pm$ 0.20**	0.15 $\pm$ 0.09**
Model	--	12957.6 $\pm$ 523.9	9.31 $\pm$ 4.57	5.97 $\pm$ 2.44	0.56 $\pm$ 0.38	0.27 $\pm$ 0.14
Captopril	40 mg/kg	7497.4 $\pm$ 430.4**	3.55 $\pm$ 1.65**	2.99 $\pm$ 0.81**	0.26 $\pm$ 0.16**	0.16 $\pm$ 0.10*
ERS-L	8 g/kg	8124.4 $\pm$ 243.5**	4.17 $\pm$ 2.00**	3.25 $\pm$ 0.81**	0.28 $\pm$ 0.15**	0.15 $\pm$ 0.09*
ERS-H	16 g/kg	8785.6 $\pm$ 296.3**	3.35 $\pm$ 1.02**	3.03 $\pm$ 0.96**	0.27 $\pm$ 0.12**	0.15 $\pm$ 0.07**

\* $P < 0.05$ , \*\* $P < 0.01$  vs model group




Fig. 2: Effect of ERS on myocardial perivascular collagen area in rats with ventricular remodeling induced by coronary artery ligation. A: sham-operated group; B: model group; C: captopril group; D: ERS 8 g·kg<sup>-1</sup> group; E: ERS 16 g·kg<sup>-1</sup> group




Fig. 3: Effect of ERS on myocardial collagen volume fraction in rats with ventricular remodeling induced by coronary artery ligation. A: sham-operated group; B: model group; C: captopril group; D: ERS 8 g·kg<sup>-1</sup> group; E: ERS 16 g·kg<sup>-1</sup> group




Fig. 4: Effect of ERS on myocardial perivascular collagen distributions of type I and III in rats with ventricular remodeling induced by coronary artery ligation. A: sham operated group; B: model group; C: captopril group; D: ERS 8 g·kg<sup>-1</sup> group; E: ERS 16 g·kg<sup>-1</sup> group

### 3. Discussion

In the last 20 years, the role of MI and the subsequent alteration in ventricular architecture of the infarcted and non-infarcted myocardium have become increasingly associated with a phenomenon known as ventricular remodeling. This process consists of left ventricular wall thinning in the infarction area, ventricular chamber dilatation, and compensatory hypertrophy of the noninfarcted portion of the myocardium

(Paul, 1995). Because collagen deposition occurs early after infarction and tends to make the shape deformation permanent (Jugdutt and Amy 1986), it would seem logical to begin therapy before collagen deposition as soon as possible. In this study, the therapy was begun next day after ligation.

Left ventricular remodeling marked by inflammation, interstitial fibrosis, cardiocyte hypertrophy, systolic and diastolic dysfunction (Tenhunen et al. 2006).




Fig. 5: Effect of ERS on myocardial collagen distributions of type I and III in rats with ventricular remodeling induced by coronary artery ligation. A: sham-operated group; B: model group; C: captopril group; D: ERS 8 g·kg<sup>-1</sup> group; E: ERS 16 g·kg<sup>-1</sup> group

**Table 3: Effects of ERS on AT<sub>1</sub>R, TNF- $\alpha$  and TGF- $\beta$ 1 mRNA expressions in rats with ventricular remodeling induced by coronary artery ligation. ( $\bar{x} \pm s$ , n = 5)**

Group	Dose	AT <sub>1</sub> R mRNA	TNF- $\alpha$ mRNA	TGF- $\beta$ 1 mRNA
Sham	--	1.49 $\pm$ 0.89**	3.14 $\pm$ 2.00**	1.51 $\pm$ 0.86**
Model	--	36.54 $\pm$ 9.82	33.48 $\pm$ 9.74	40.05 $\pm$ 18.99
Captopril	40 mg/kg	4.38 $\pm$ 4.91**	6.86 $\pm$ 6.04**	2.94 $\pm$ 1.33**
ERS-L	8 g/kg	13.90 $\pm$ 8.83**	9.84 $\pm$ 3.17**	6.56 $\pm$ 2.99**
ERS-H	16 g/kg	3.70 $\pm$ 0.48**	5.60 $\pm$ 2.74**	3.01 $\pm$ 2.07**

\* $P < 0.05$ , \*\* $P < 0.01$  vs model group

The present study demonstrated that administration of ERS could significantly reduce the LVWI, HWI; decrease the content of Ang II, Hyp; diminish cardiocyte cross-section area and ameliorate collagen deposition. In addition, ERS could down regulate the gene expressions of AT<sub>1</sub>R, TNF- $\alpha$  and TGF- $\beta$ 1 mRNA in myocardium. The mechanism may be versatile and complex.

There is considerable evidence that the renin-angiotensin-aldosterone system is activated after acute myocardial infarction and that activation of this system has important physiological and long-term morphological consequences.

In the study, the degree of cardiac hypertrophy was assessed by the increase of LVWI, HWI and the transverse area of cardiocytes. Cardiocyte hypertrophy was demonstrable microscopically, with an up to 100% increase in cell transverse area. The result is similar to the report (Anversa et al. 1985). Treatment for four weeks with low and high dose of ERS prevented cardiac hypertrophy.

There is now evidence that LV hypertrophy is associated with the induction of gene expression for angiotensin converting enzyme (ACE) and increased local synthesis of Ang II within the ventricular myocardium (Schunkert et al. 1990). Inhibition of ACE is beneficial for suppressing ventricular remodeling (Schieffer et al. 1994). ERS significantly attenuated the increase of Ang II concentration, it may be one of the mechanisms of its action against cardiac hypertrophy.

Many reports underscored the importance of the interaction between cytokines and the RAS in ventricular remodeling in the progression of CHF (Frolkis et al. 2001; Gurlek et al. 2001; Tsutamoto et al. 2000). The hearts that have suffered a MI have an increase in the expression of all the pro-inflammatory cytokines. TNF- $\alpha$  is among the most important inflammatory cytokines (Testa et al. 1996; Irwin et al. 1999). Ang II and TNF- $\alpha$  can potentiate the effects of each other, resulting in a vicious cycle towards CHF. Ang II secreted from cytoplasmic granules

and the corresponding hypertrophic response is mediated by AT<sub>1</sub> receptors (AT<sub>1</sub>R). In the present study, treatment with ERS significantly decreased the elevated gene expression of AT<sub>1</sub>R alongside a decrease in TNF- $\alpha$  gene expression in the heart.

Ang II and the cytokine TGF- $\beta$ 1 are connected in the pathogenesis of cardiac remodeling. Ang II can up-regulate TGF- $\beta$ 1 expression via activation of AT<sub>1</sub>R in cardiocyte and fibroblast (Diniz et al. 2007). TGF- $\beta$ 1 also can mediate Ang II induced structural ventricular remodeling in an autocrine/paracrine manner (Gray et al. 1998). TGF- $\beta$ 1 increases early in the infarct zone, stimulating macrophage and fibroblast chemotaxis and fibroblast proliferation (Desmouliere et al. 1993). Proliferation of fibroblasts during cardiac hypertrophy is a major mechanism contributing to the development of fibrosis, and fibroblasts accumulate in fibrotic and scarred areas (Deblois et al. 2005). In the present study, we observed that ERS inhibited myocardial fibrosis in the non-infarcted myocardium. Like captopril (van Krimpen et al. 1991), ERS given early during healing postinfarction but not later, inhibited collagen deposition in the non-infarcted myocardium as well.

Additionally, the major components of cardiac extracellular matrix (ECM) proteins are collagens, of which collagen type I composes approximately 85% (Heeneman et al. 2003). In particular, the accumulation of collagen type I which is characterized by tensile strength (Weber 1989), may contribute to myocardial stiffness by limiting the motion of cardiocytes, and may promote arrhythmias by electrical isolation of adjacent cardiocytes. Type III collagen, which is deposited during healing, has little tensile strength. The type I/III collagen ratio might increase between 21 and 90 days after MI in rats (Jugdutt et al. 1996). Therefore, the suppressive effect of ERS on collagen type I and III observed in the present study may provide a molecular substrate for the beneficial effects on LV structure and function, and the antifibrotic effects appeared to be attributable to the suppressive effect on the TNF- $\alpha$  secretion.

ACEI blocks local angiotensin II and TGF- $\beta_1$ , the latter being a potent modulator of the extracellular matrix and a stimulator of collagen synthesis (Sadoshima and Izumo 1993). So ACEI can reduce the fibrosis and inhibit collagen deposition that accompanies LV remodeling in a variety of experimental models (Brilla et al. 1990). ACEI's inhibition of Ang II and TGF- $\beta_1$  might be especially important mediators of antagonizing myocardial hypertrophy and fibrosis.

ERS has beneficial effects against ventricular remodeling. The mechanism may be related to decreasing the level of Ang II and cardiac fibrosis, modulating AT<sub>1</sub>R, TNF- $\alpha$  and TGF- $\beta_1$  mRNA expressions associated with cardiac hypertrophy.

The effects of ERS are similar to those of ACEI, which are well known to prevent ventricular remodeling after MI. These results suggest that administration of a sufficient dose of ERS may have the same cardioprotective effects as ACEI in this animal model. However, the mechanism of ERS in detail and the effective components of ERS remain to be further explored.

## 4. Experimental

### 4.1. Herb extract and drug

*Radix Scrophulariae* was purchased from Yanghetang Medical Material Company (Shanghai, China), authenticated by Dr. Zhao, Department of Pharmacognosy, Shanghai University of TCM. It was decocted with distilled water, and the water soluble extract yield was obtained for the experiment (60% yield of prepared extracts in terms of starting crude material). Captopril tablets (batch number: 070704) were purchased from Shanghai Hengshan Pharmaceutical Co. Ltd., China. It was dissolved in distilled water.

### 4.2. Animals

Male Sprague Dawley rats weighing 220–250 g were obtained from the Experimental Animal Center, Chinese Academy of Sciences (Shanghai, China). The rats were maintained on 12-h light/12-h dark cycle in the room at a temperature of  $23 \pm 1^\circ\text{C}$ , humidity of  $40 \pm 5\%$ . These rats received humane care and had free access to a standard diet and drinking water. The animal experiment was carried out according to the Guide for Care and Use of Laboratory Animals, published by US guidelines (NIH publication #85-23, revised in 1985).

### 4.3. Animal model and protocol

Rats were anesthetized by intraperitoneal administration of pentobarbital (40 mg kg<sup>-1</sup>), intubated and artificially ventilated. After left thoracotomy and pericardiotomy, the left coronary artery was ligated approximately 2 mm from its origin with a 4-0 silk suture. Then the thorax was closed immediately. The survival rate of the surgery was about 60%. After coronary ligation, all rats were randomly assigned to following 4 groups: model, captopril (40 mg kg<sup>-1</sup>), ERS (8, 16 g kg<sup>-1</sup>). The sham-operated rats underwent the same operation without ligation. After the surgery, the rats of all groups were given antibiotic (penicillin,  $1 \times 10^4$  U·kg<sup>-1</sup>, i.m.) and allowed to recover.

The rats were orally administrated with the corresponding extract or drug at the above described doses once a day. And distilled water was administered in the same manner for the sham-operated and model control groups. Treatment started from the next day after operation and continued for four weeks.

### 4.4. LVWI and HWI

Four weeks after the surgery the rats were weighed and then sacrificed by decapitation. The hearts were taken out and the left ventricle was separated from the atria, aorta and adipose tissue. The heart weight and left ventricle weight were measured, then, LVWI and HWI were estimated by calculating HW-to-BW and LVW-to-BW ratio.

The LV was divided by a cross section in the middle. The upper half part was fixed with 10% formaldehyde for pathological sectioning. The lower half part was further dissected into 3 parts and then quickly frozen in liquid nitrogen, and preserved at  $-70^\circ\text{C}$  for biochemical and gene expression analysis.

### 4.5. Angiotensin II and hydroxyproline determination

Tissue homogenates (2%) were made for determination of Ang II and Hyp concentrations. The homogenized tissue was centrifuged ( $4^\circ\text{C}$ ,

1780 g, 15 min.) and the supernatant liquid was collected for measurement. Ang II concentrations were quantified by radioimmunoassay with an iodine [<sup>125</sup>I] Ang II radioimmunoassay kit (Beijing North Institute of Biological Technology, Beijing, China). Hyp concentrations were detected by ultraviolet spectrophotometry with the Hyp kit (Nanjing Jiancheng Institute of Bioengineering, Nanjing, China). Protein concentrations of myocardial homogenates were assayed with the Coomassie Brilliant Blue Kit (Nanjing Jiancheng Institute of Bioengineering, Nanjing, China). Ang II and Hyp concentrations were corrected for protein concentrations.

### 4.6. Histological analysis

The ventricles were fixed with 10% formaldehyde, embedded in paraffin, sectioned at 5  $\mu\text{m}$  in thickness, stained with hematoxylin and eosin (HE) for measurement of cardiocyte cross section size, and stained with Sirius red in aqueous saturated picric acid for examination of perivascular and interstitial fibrosis in myocardium. Each sample slice was photographed with the digital camera (Olympus BX51, Japan) connected to microscope (400  $\times$  magnification) and computer. They were analyzed with image-Pro Plus 6.0 analyzing software (Media Cybernetics, U.S.A.).

Three fields per sample slice were randomly selected and 20 myocardial cells per-field were chosen to calculate cell size. Another 3 fields were randomly selected for calculating the ratio of picric Sirius red stained fibrosis area to total myocardial area. The perivascular fibrosis was determined as the ratio of the area of fibrosis surrounding the vessel wall to the total vessel area (Shinzato et al. 2007). The contents of subtype I and III collagen in myocardium were determined by using microscope with polarimetric filter analysis at the same time.

### 4.7. Real-time RT-PCR determination

Total RNA was extracted from the tissues by using the TRIzol reagent (Invitrogen) according to the manufacturer's instructions. RNA yields and purity were assessed by spectrophotometric analysis. Total RNA (1  $\mu\text{g}$ ) from each well was subjected to reverse transcription with random hexamer primers, deoxynucleoside triphosphates (dNTPs), and Maloney murine leukemia virus (M-MLV) reverse transcriptase in a total reaction volume of 20  $\mu\text{l}$ .

The real-time RT-PCR reactions (50  $\mu\text{l}$ ) consisted of 32.5  $\mu\text{l}$  SYBR Green Mix, 1  $\mu\text{l}$  mixed primers, 2  $\mu\text{l}$  cDNA and 14.5  $\mu\text{l}$  double-distilled water. A typical protocol included incubation at  $50^\circ\text{C}$  for 2 min. and taq activation at  $95^\circ\text{C}$  for 10 min. followed by 40 cycles with  $95^\circ\text{C}$  denaturation for 20 s,  $55^\circ\text{C}$  annealing for 30 s, and  $72^\circ\text{C}$  extension for 30 s. The sequences of primer were as follows: AT<sub>1</sub>R mRNA sense: 5'-GCACACTGGCAATG-TAATGC-3', AT<sub>1</sub>R mRNA anti-sense: 5'-GTTGAACAGAACAAAGTGA-CC-3'; TNF- $\alpha$  mRNA sense: 5'-TGACTTCTCCTGGTATGAAATGG-3', TNF- $\alpha$  mRNA anti-sense: 5'-TGACTTCTCCTGGTATGAAATGG-3'; TGF- $\beta_1$  mRNA sense: 5'-GCTGCTGACCCCACTGAT-3', TGF- $\beta_1$  mRNA anti-sense: 5'-TGCCGGACAACCTCCAGTGA-3'; GAPDH sense: 5'-CCGAGGGCCCACTAAAGG-3', GAPDH anti-sense: 5'-GCTG-TTGAAGTCACAGGAGACAA-3'.

Real-time RT-PCR was carried out with 7300 Sequence Detection System and data analyzed by ABI Prism 7300 SDS Software.

All values obtained with the AT<sub>1</sub>R, TNF- $\alpha$  or TGF- $\beta_1$  primers were normalized to the values obtained with the GAPDH primers. The results were expressed as the relative integrated intensity.

### 4.8. Statistics

All data were expressed as the mean  $\pm$  SD. Statistical analysis was performed by one-way analysis of variance (ANOVA) for multiple comparisons, followed by Dunnett's test to evaluate the difference between two groups through the software of SPSS 13.0 for Windows (SPSS Inc, Chicago, IL, USA).

Acknowledgement: This study was supported by a grant from the National Natural Science Foundation of China (No. 30572379).

## References

- Anversa P, Beghi C, Kikkawa Y, Olivetti G (1985) Myocardial response to infarction in the rat. Morphometric measurement of infarct size and myocyte cellular hypertrophy. *Am J Pathol* 118: 484–492.
- Brilla CG, Pick R, Tan LB, Janicki JS, Weber KT (1990) Remodeling of the rat right and left ventricles in experimental hypertension. *Circ Res* 67: 1355–1364.
- Delouis D, Tea BS, Beaudry D, Hamet P (2005). Regulation of therapeutic apoptosis: a potential target in controlling hypertensive organ damage. *Can J Physiol Pharmacol* 83: 29–41.

- Desmouliere A, Geinoz A, Gabbiani F, Gabbiani G (1993) Transforming growth factor-beta 1 induces alpha-smooth muscle actin expression in granulation tissue myofibroblasts and in quiescent and growing cultured fibroblasts. *J Cell Biol* 122:103–111.
- Diniz GP, Carneiro-Ramos MS, Barreto-Chaves ML (2007) Angiotensin type 1 (AT1) and type 2 (AT2) receptors mediate the increase in TGF-beta1 in thyroid hormone-induced cardiac hypertrophy. *Pflugers Arch* 454: 75–81.
- Frolkis I, Gurevitch J, Yuhas Y, Iaina A, Wollman Y, Chernichovski T, Paz Y, Matsa M, Pevni D, Kramer A, Shapira I, Mohr R (2001) Interaction between paracrine tumor necrosis factor-alpha and paracrine angiotensin II during myocardial ischemia. *J Am Coll Cardiol* 37: 316–322.
- Gray MO, Long CS, Kalinyak JE, Li HT, Karliner JS (1998) Angiotensin II stimulates cardiac myocyte hypertrophy via paracrine release of TGF-beta 1 and endothelin-1 from fibroblasts. *Cardiovasc Res* 40: 352–363.
- Gurlek A, Kilickap M, Dincer I, Dandachi R, Tutkak H, Oral D (2001) Effect of losartan on circulating TNFalpha levels and left ventricular systolic performance in patients with heart failure. *J Cardiovasc Risk* 8: 279–282.
- Heeneman S, Cleutjens JP, Faber BC, Creemers EE, van Suylen RJ, Lutgens E, Cleutjens KB, Daemen MJ (2003) The dynamic extracellular matrix: intervention strategies during heart failure and atherosclerosis. *J Pathol* 200: 516–525.
- Irwin MW, Mak S, Mann DL, Qu R, Penninger JM, Yan A, Dawood F, Wen WH, Shou Z, Liu P (1999) Tissue expression and immunolocalization of tumor necrosis factor-alpha in postinfarction dysfunctional myocardium. *Circulation* 99: 1492–1498.
- Jialin Teng, Xingguang Li, Yimin Le (2007) *Chinese Materia Medica* 9: 344.
- Jugdutt BI, Amy RW (1986) Healing after myocardial infarction in the dog: changes in infarct hydroxyproline and topography. *J Am Coll Cardiol* 7: 91–102.
- Jugdutt BI, Joljart MJ, Khan MI (1996) Rate of collagen deposition during healing and ventricular remodeling after myocardial infarction in rat and dog models. *Circulation* 94: 94–101.
- Paul S (1995) The pathophysiologic process of ventricular remodeling: from infarct to failure. *Crit Care Nurs Q* 18: 7–21.
- Sadoshima J, Izumo S (1993) Molecular characterization of angiotensin II –induced hypertrophy of cardiac myocytes and hyperplasia of cardiac fibroblasts. Critical role of the AT1 receptor subtype. *Circ Res* 73: 413–423.
- Schieffer B, Wirger A, Meybrunn M, Seitz S, Holtz J, Riede UN, Drexler H (1994) Comparative effects of chronic angiotensin-converting enzyme inhibition and angiotensin II type 1 receptor blockade on cardiac remodeling after myocardial infarction in the rat. *Circulation* 89: 2273–2282.
- Schunkert H, Dzau VJ, Tahn SS, Hirsch AT, Apstein CS, Lorell BH (1990) Increased rat cardiac angiotensin converting enzyme activity and mRNA expression in pressure overload left ventricular hypertrophy. *J Clin Invest* 86: 1913–1920.
- Shinzato T, Ohya Y, Nakamoto M, Ishida A, Takishita S (2007) Beneficial effects of pioglitazone on left ventricular hypertrophy in genetically hypertensive rats. *Hypertens Res* 30: 863–873.
- Tenhunen O, Rysa J, Ilves M, Soini Y, Rushkoaho H, Leskinen H (2006) Identification of cell cycle regulatory and antiinflammatory genes as predominant targets of p38 mitogenactivated protein kinase in the heart. *Circ Res* 99: 485–493.
- Testa M, Yeh M, Lee P, Fanelli R, Loperfido F, Berman JW, LeJemtel TH (1996) Circulating levels of cytokines and their endogenous modulators in patients with mild to severe congestive heart failure due to coronary artery disease or hypertension. *J Am Coll Cardiol* 28: 964–971.
- Tsutamoto T, Wada A, Maeda K, Mabuchi N, Hayashi M, Tsutsui T, Ohnishi M, Sawaki M, Fujii M, Matsumoto T, Kinoshita M (2000) Angiotensin II type 1 receptor antagonist decreases plasma levels of tumor necrosis factor alpha, interleukin-6 and soluble adhesion molecules in patients with chronic heart failure. *J Am Coll Cardiol* 35: 714–721.
- van Krimpen C, Schoemaker RG, Cleutjens JP, Smits JF, Struyker-Boudier HA, Bosman FT, Daemen MJ (1991) Angiotensin I converting enzyme inhibitors and cardiac remodeling. *Basic Res Cardiol* 86: 149–155.
- Weber KT (1989) Cardiac interstitium in health and disease: the fibrillar collagen network. *J Am Coll Cardiol* 13: 1637–1652.