

Department of Orthopaedics and Traumatology¹, Longhua Hospital, Shanghai University of Traditional Chinese Medicine; Institute of Spine², Shanghai University of Traditional Chinese Medicine; Institute of Integrated Traditional Chinese Medicine and Western Medicine³, Huashan Hospital, Fudan University, Shanghai, China

The osteogenetic effect of astragaloside IV with centrifugating pressure on the OCT-1 cells

QIN BIAN^{1,2,3,*}, JIAN-HUA HUANG^{3,*}, QIAN-QIAN LIANG^{1,2}, BING SHU^{1,2}, WEI HOU^{1,2}, HAO XU^{1,2}, YONG-JIAN ZHAO^{1,2}, SHENG LU^{1,2}, QI SHI^{1,2}, YONG-JUN WANG^{1,2}

Received July 29, 2010, accepted August 2, 2010

Prof. Yong-jun Wang, Institute of Spine, Shanghai University of Traditional Chinese Medicine, Shanghai, 200032, China

yjwang88@126.com

*Both authors contributed equally to this work

Pharmazie 66: 63–68 (2011)

doi: 10.1691/ph.2011.0219

Astragaloside IV (ASI), a pure compound derived from *Radix Astragali*, is commonly used in degenerative bone diseases such as osteoporosis. Our previous study identified *in vivo* the osteogenetic effect of *Fu Fang Qi She Pills* (FFQSP), a Chinese herbal formula containing *Radix Astragali* from which ASI was extracted. In this study, we investigated the osteogenetic effects of ASI under the conditions of centrifugating pressure on OCT-1 cells. These preosteoblasts were grown in 3D-culture, and treated with ASI at 50 $\mu\text{mol/l}$ with centrifugation at 200 rpm, 500 rpm for 3 and 5 days. Morphocytological examination, morphometry of alkaline phosphatases (ALP) staining was performed. Expression of type I collagen (Col I) was detected by immunocytochemistry assays. ALP, Col1a2, Osteocalcin (OC), and runt-related transcription factor-2 (Runx2) mRNA expression were determined via real-time PCR. The results showed ASI plus 500 rpm for 3 days and ASI plus 200 rpm for 5 days significantly induced osteogenesis related protein and gene expression. We concluded that ASI would promote osteogenesis when cells of preosteoblast OCT-1 were subjected to proper centrifugating pressure and a pertinent period of time.

1. Introduction

Astragaloside IV (ASI) is one of the active components of *Radix Astragali* and usually used as standard substance in quality evaluation since its high correlation to various pharmacological actions of the herb. For example, it was effective in improving endothelial dysfunction, cardiomyocyte damage, memory function and delaying the aging, which was somewhat related to its antioxidant activity (Hu et al. 2008; Li et al. 2007; Qiu et al. 2010; Qu et al. 2009; Xu et al. 2008; Zhang 2007a etc.; Zhang 2010c etc.; Zhang 2006d etc.; Zhang 2006e etc.; Zhao et al. 2009). On the other hand, ASI prevented hepatic fibrosis, chronic experimental asthma, experimental Parkinson's disease, diabetic complications such as peripheral neuropathy since it acted as an aldose-reductase inhibitor (Chan et al. 2009; Chen et al. 2008; Du et al. 2008; Liu 2009a etc; Lv et al. 2010; Motomura et al. 2009; Yu et al. 2006; Zhu et al. 2008). It also elicited the activity of anticancer, anti hepatitis B and Coxsackie virus B3 (CVB3) (Qi et al. 2010; Wang 2009a etc.; Zhang 2006f etc.). Recently, it was reported that AST could promote bone mesenchymal stem cells (MSCs) proliferation and differentiation in rats, partly by increasing stem cell factor (SCF) mRNA and ALP excretion (Liu 2006b etc.; Tan et al. 2010).

However, the effects of ASI on osteogenesis, especially combined with mechanical pressure, an important factor of promoting bone formation, remain largely unknown. In our previous study, *Fu Fang Qi She Pills* (FFQSP) a Chinese herbal compound composed of *Radix Astragali* obviously promoted

lumbar vertebral bone formation of rats in upright posture (Bian et al. 2010). The content of ASI in FFQSP was within the range of 5.175~36.225 μg determined by HPLC-ELSD (Zhang 2008b etc.). In the present study, we focus on the effect of ASI on osteogenesis *in vitro* associated with overloading imposed by centrifugation, which mimics mechanical environment in upstanding posture *in vivo* (Wong and Rabie 2008).

2. Investigations, results and discussion

MTT is a common method for observing cellular proliferation. However, in our study, cells were cultured in a solid phase, which brought difficulties in MTT detection. So, we used frozen section and H&E staining method for morphological examination. It showed that cells were able to anchor in the agarose gel matrix which is hyaline and like branches of trees; little cells were located on the inner section of the matrix. In the 200 rpm groups and 500 rpm groups, more cells could be seen. Cells tended to be bigger when treated with ASI, but no significant changes in cellular shapes were seen. These results suggested that the preosteoblasts tended to differentiate into osteoblasts. Three ASI groups showed an increasing amount of cells after centrifugation for 3 d and 5 d, and the extra cellular matrix was augmented. These changes might acclimatize the cells in hypergravity condition.

Type I collagen (Col I) is expressed both in the committed preosteoblast and the mature osteoblast (Robling et al.




Fig. 1: H&E staining for OCT-1 cells cultured in agarose gel at 1% concentration. Three ASI groups showed an increasing amount of cells after centrifugation for 3d and 5d, and the extra cellular matrix was augmented ($\times 200$)

2006). It is reported that type I collagen was enhanced when osteoblasts were exposed to pressure (Nagatomi et al. 2003). In our study, the immunofluorescence study demonstrated OCT-1 cells expressed increased Col I protein when treated with ASI after centrifugation for 3 days. However, there were no significant changes between 200 rpm group and 200 rpm + ASI group. ASI, with or without centrifugation by 200 rpm for 5 days significantly promoted Col I expression while ASI with 500 rpm centrifugation slightly inhibited it. The result is consistent with the changes in Col I mRNA level. It indicated that ASI could

facilitate the expression of Col I protein and mRNA, especially with the assistant of centrifugation at 500 rpm. Alkaline phosphatase (ALP) expressed by mature osteoblasts, is a key enzyme in the mineralization process (Robling et al. 2006). Although ALP activity is an early indicator of bone formation and is transient, making it a less sensitive marker of osteogenesis, especially in response to mechanical stimuli (Lian and Stein 1995). Osteocalcin (OC) is one of abundant bone extracellular matrix proteins and molecular markers of osteoblast differentiation (Robling et al. 2006; Candelieri et al. 2001). We




Fig. 2: ALP staining for 3D-cultured OCT-1 cells and Morphometry of ALP staining. OCT-1 cells were cultured in centrifugal condition at 0 rpm, 200 rpm, and 500 rpm with or without ASI for 3d and 5d. Brown sediments indicated the positive staining ($\times 400$). Histogram shows the morphometry of ALP staining ($n = 6$). *: $p < 0.05$, **: $p < 0.01$ ASI+centrifugation vs centrifugation at the same centrifugal force




Fig. 3: Immunofluorescence study of Coll. OCT-1 cells expressed increased Coll protein when treated with ASI after centrifugation for 3 days, especially in ASI+500 rpm group. ASI, with or without centrifugation by 200 rpm for 5 days significantly promoted Coll expression while ASI with 500 rpm centrifugation slightly inhibited it ($\times 100$)

found both ALP and OC mRNA up-regulated after three days. In 5 d groups, ASI raised both ALP and OC mRNA level at 0 rpm and 200 rpm. On the contrast, 500 rpm centrifugation inhibited the osteogenetic effect of ASI. The morphometry of ALP staining showed the accordant tendency with the changes in mRNA expression.

After centrifugation for three days, the up-regulation of ALP, Col1a2 and OC genes suggested an adaptation of ASI to OCT-1 cells in centrifugation condition, and 500 rpm+ASI showed the most effective in osteogenesis on this phase. Nevertheless, 500 rpm+ASI attenuated the osteogenetic effect on the fifth day by down-regulating the related gene mRNAs. It was supposed that ASI started to produce an effect on the third day, and this

effect was enhanced by the support of high dose of pressure (500 rpm). However, this osteogenetic effect of ASI and centrifugation might become fatigued and over its threshold value at later period, leading to loss the action due to overconsumption.

The transcription factors runt-related transcription factor-2(Runx2) is considered a "master gene" that plays a critical role in the formation of the skeleton (Komori et al. 1997). In detail, expression of Runx2 is required to push the proliferating precursor cells toward the osteoblast lineage and away from other lineages also yielded by the mesenchymal stem cell. Further differentiation of the preosteoblast into a mature, bone forming osteoblast phenotype requires the expression of Runx2 (Ducy et al. 1997). Our results showed that Runx2 mRNA increased only

Table: Sequences of primers used in the Real-time PCR

| Genes | Forward primer | Reverse primer | Product length(bp) |
|------------|-------------------------|--------------------------|--------------------|
| ALP | TGACCTTCTCTCCTCCATCC | CTTCCTGGGAGTCTCATCCT | 183 |
| Beta-actin | GGAGATTACTGCCCTGGCTCCTA | GACTCATCGTACTCCTGCTTGCTG | 150 |
| Col 1a2 | TCCTGGCAATCGTGGTTCAA | ACCAGCTGGGCCAACATTTTC | 133 |
| OC | TGCCCTCCTGGTTCATTCT | TTCTGTTCCCTCCCTGCTGT | 119 |
| Runx2 | CCATAACGGTCTTCACAAATCCT | TCTGTCTGTGCCTTCTTGGTTC | 99 |




Fig. 4: The mRNA expression of ALP, Col1a2, OC and Runx2. The columns represent the mean \pm SE of three independent experiments. *: $p < 0.05$, **: $p < 0.01$ ASI+centrifugation vs centrifugation at the same centrifugal force

after centrifugation at 200 rpm, but there was no significant difference at 500 rpm, which indicated a magnitude- but not a time-dependent induction by centrifugating pressure. However, ASI induced increased Runx2 gene expression merely in 5d 500 rpm group. These variations were different from those of other three

genes, hinting that the upstream of Runx2 includes several other molecules besides ALP and Col I molecules (Alliston et al. 2001). And, the targets and effects were manifold of ASI with centrifugating pressure on osteogenesis, which caused discordant changes in Runx2 molecule and other three molecules.

We concluded that ASI would promote osteogenesis combined with proper centrifuging pressure and pertinent period of time in preosteoblast OCT-1.

3. Experimental

3.1. Materials

Astragaloside Ir (ASI) was purchased from Shanghai Pharmaceutical Company Ltd, China. The preosteoblast strain OCT-1 originates from the skull of osteocalcin driven element SV-40 transgenic mice (OC-Tag), presented by Prof. Di Chen (University of Rochester, U.S.). High-glucose Dulbecco's modified Eagle's medium (DMEM) (Gibico, US), fetal bovine serum (FBS) (HyClone, Logan, UT), antibiotic/antimycotic (Gibico, US), trypsin-EDTA (Gibico, US), agarose (Biowest, France), RNAPrep pure Cell Kit (Tiangen Biotech Co., LTD, Beijing, China), Histostain-Plus Kit (Jingmei Biotech, Shenzhen, China), ALP staining kits (Jiancheng Biotech Institute, Nanjing, China), and Collagen I antibody (Abcam Ltd., Cambridge, UK).

3.2. Preparation of ASI

The molecular weight of ASI is 784.98 g/l. 50 nmol/l ASI was obtained by 39.24 mg ASI powders and 1 ml DMSO. Then, the above fluid was diluted into 100 μ mol/l using double concentration of DMEM containing 4% FBS.

3.3. Grouping

There were two groups based on different durations: 3 days and 5 days. Each group was divided into six subgroups according to different centrifugal speed: 0 rpm, 200 rpm, and 500 rpm with or without ASI.

3.4. 3D culture

Cells were cultured to confluence at 8×10^6 mononuclear cells per cm^2 in medium consisting of high DMEM, 2% fetal bovine serum, 1% antibiotic/antimycotic, and then detached with 0.05% trypsin/1 mM EDTA. An equal volume of cells were suspended at 4×10^7 cells per millilitre in double concentration of high DMEM, fetal bovine serum and antibiotic/antimycotic, and then combined with the same volume of agarose solution, which were assembled by autoclaving 2% agarose in ultrapure water and then cooling the solution to 40 °C. The final concentration was 2×10^7 cells per millilitre in 3D culture system of 1% agarose and 50 μ mol/l ASI, 2% FBS.

3.5. Calculation of centrifugal speed

It has been reported that human's L₄-L₅ disc produced a pressure of 0.1 MPa when lying prone. More value has been measured as followed: relaxed standing, 0.5 MPa; and lifting a 20-kg weight with round flexed back, 2.3 MPa. From the first formula: $P = mg/S$ (mechanism pressure), $P = \rho gh$ (hydraulic pressure), it can be derived that regardless of what pressure the system is at, pressure is related to gravity "g". According to the above formula, we supposed that $G_{\text{lying}} = P/\rho h = 0.1/\rho h = 1g$, $G_{\text{relaxed standing}} = P/\rho h = 0.5/\rho h = 5g$, and $G_{\text{flexed back}} = P/\rho h = 2.3/\rho h = 23g$. The value shows that the two standing postures need to overcome a force of gravity of 5g and 23g respectively. For the second formula as follows, we put 5g and 23g as an RCF into the calculation. according to Eq. (1).

$$\text{RPM} = \sqrt{\frac{\text{RCF}}{1.118 \times 10^5 \times (\text{Radius})}}$$

RCF : centrifugal force(g)

RPM : rotation speed per minute

Radius : length of shaft radius

(1)

The highest rotation speed of the centrifuge (Allegra™ X-22R, Beckman, USA) was 10000 rpm and the maximum centrifugal force was 10528 g. According to the second formula, the radius was worked out as 9.4168 cm. We put the value into the second formula as follows: $\sqrt{1/(1.118 \times 10^{-5})} \times 9.4168 = 97$ rpm; $\sqrt{5/(1.118 \times 10^{-5})} \times 9.4168 = 218$ rpm; $\sqrt{23/(1.118 \times 10^{-5})} \times 9.4168 = 467$ rpm. The values of 5g or 23g may be obtained using the centrifuge at 218 rpm or 467 rpm. In fact, the centrifuge can work at 200 rpm and 500 rpm limited by its accuracy. Furthermore, 5g and 23g correspond to physiological level of five ustrain and 23 ustrain respectively (Hughes-Fulford 2003). The 200 rpm groups and the 500 rpm groups were centrifuged for three hours once a day under room conditions for one day or five days. The control groups were located in the same environment,

but without centrifugation. All the samples were put back into an incubator (37 °C, 5% CO₂) immediately after centrifugation or after three hours standing under room conditions.

3.6. H&E Staining

Cells were fixed in 4% paraformaldehyde for 20 min, and then washed with PBS. Frozen sections were stained with hematoxylin and eosin. A morphometric study was performed by an image auto-analysis system (Olympus BX50; Japan).

3.7. Morphometry of ALP

The substrates were dissolved by dimethylformamide. The buffer and solid violet B were then added and mixed to form the fluid. The fixed slides were put into the fluid, and cultured at 37 °C for 45 min. Positive staining of brown sediments could be seen and counted by image auto-analysis system (Olympus BX50; Japan) after counterstaining with hematoxylin.

3.8. Immunofluorescence analysis

Sections were deparaffinised and incubated in 3% H₂O₂ to block endogenous peroxidase activity for 15 min, washed with PBS for 2 min, followed by a digestion with protease K for 10 min. Sections were blocked with 5% BSA in which the secondary antibody was produced, and then washed with PBS three times for 2 min. Specific primary rabbit polyclonal antibodies against type I collagen (1:200) were incubated overnight at 4 °C. After extensive washing with PBS three times for 2 min. A fluorescence microscope was used for observation and a photograph was taken within one hour to prevent fluorescent from decreasing.

3.9. Real-time PCR detection

Cells were directly processed following RNA prep pure Cell Kit protocol. One microgram of total RNA was reversely transcribed using the Advantage RT-for-PCR kit (Qiagen, Valencia, CA) following the manufacturer's protocol. Freshly transcribed cDNA (1 μ l) was used for quantitative real-time PCR using SYBR Green (Bio-Bad, Hercules, CA) to monitor DNA synthesis using specific primers (Table) designed by TaKaRa Biotechnology Co. Ltd. The PCR was carried out in RotorGene real-time DNA amplification system (Corbett Research, Sydney, Australia) using the following cycling protocol: a 95 °C denaturation step for 15 min followed by 45 cycles of 95 °C denaturation (20 s), 56–62 °C annealing (30 s), and 72 °C extension (30 s). Gene expression was normalized to the housekeeping gene β -actin. PCR products were subjected to a melting curve analysis, and the data were analyzed and quantified with the RotorGene6.0 analysis software.

3.10. Statistical analysis

The data is expressed as mean \pm SE and statistical significance was calculated using a Student t test and analysis of variance by SPSS software (SPSS Inc, Chicago, USA). The significance level was defined as $p < 0.05$.

Acknowledgments: We thank Professor Di Chen, School of Medicine and Dentistry, Department of Orthopaedics, University of Rochester, USA for providing OCT-1 cells, and Professor Yi-ping Li, department of cytokine biology, the Forsyth Institute & Harvard School of dental medicine, Massachusetts, USA for manuscript revision. This work was supported by National Basic Research Program in China (973 Plan, 2010CB530400), Youth Health research project, the National Science Fund for Distinguished Young Scholars (30625043), the International Cooperation Programs of National Natural Science Foundation of China (30710103904), the National Natural Foundation Committee (NSFC) of China key grant (30930111), the Project of National Natural Science Foundation of China (30701118).

References

- Alliston T, Choy L, Ducey P, Karsenty G, Derynck R (2001) TGF- β -induced repression of CBFA1 by Smad3 decreases cbfa1 and osteocalcin expression and inhibits osteoblast differentiation. [EMBO J 20: 2254–2272.](#)
- Bian Q, Liang QQ, Hou W, Lu S, Zhao YJ, Wang YJ (2010) Observation on the effect of Fu Fang Qi She Pian on upright posture rats' vertebral bone formation. *Chin J Spin Spine Cord* 20: 311–316.
- Candelieri GA, Liu F, Aubin JE (2001) Individual osteoblasts in the developing calvaria express different gene repertoires. [Bone 28: 351–361.](#)

- Chan WS, Durairajan SS, Lu JH, Wang Y, Xie LX, Kum WF, Koo I, Yung KK, Li M (2009) Neuroprotective effects of Astragaloside IV in 6-hydroxydopamine-treated primary nigral cell culture. Neurochem Int 55: 414–422.
- Chen J, Gui D, Chen Y, Mou L, Liu Y, Huang J (2008) Astragaloside IV improves high glucose-induced podocyte adhesion dysfunction via α 3 β 1 integrin upregulation and integrin-linked kinase inhibition. Biochem Pharmacol 76: 796–804.
- Du Q, Chen Z, Zhou LF, Zhang Q, Huang M, Yin KS (2008) Inhibitory effects of astragaloside IV on ovalbumin-induced chronic experimental asthma. Can J Physiol Pharmacol 86: 449–457.
- Ducy P, Zhang R, Geoffroy V, Ridall AL, Karsenty G (1997) *Osf2/Cbfa1*: a transcriptional activator of osteoblast differentiation. Cell 89:747–754.
- Hu JY, Han J, Chu ZG, Song HP, Zhang DX, Zhang Q, Huang YS (2008) Astragaloside IV attenuates hypoxia-induced cardiomyocyte damage in rats by upregulating superoxide dismutase-1 levels. Clin Exp Pharmacol Physiol 36: 351–357.
- Hughes-Fulford M (2003) Function of the cytoskeleton in gravisensing during spaceflight. Adv Space Res 32: 1585–1593.
- Komori T, Yagi H, Nomura S, Yamaguchi A, Sasaki K, Shimizu Y, Bronson RT, Gao YH, Inada M, Sato M, Okamoto R, Kitamura Y, Yoshiki S, Kishimoto T (1997) Targeted disruption of *Cbfa1* results in a complete lack of bone formation owing to maturational arrest of osteoblasts. Cell 89: 755–764.
- Li WZ, Li WP, Yin YY (2007) Effects of AST and ASI on metabolism of free radical in senescent rats treated by HC. Zhongguo Zhong Yao Za Zhi 32: 2539–2542.
- Lian JB, Stein GS (2001) Development of the osteoblast phenotype: molecular mechanisms mediating osteoblast growth and differentiation. Iowa Orthop J 15: 118–140.
- Liu H, Wei W, Sun WY, Li X (2009) Protective effects of astragaloside IV on porcine-serum-induced hepatic fibrosis in rats and *in vitro* effects on hepatic stellate cells. J Ethnopharmacol 122: 502–508.
- Liu HJ, Wang XP, Lin J, Yu YC, Jiang XQ, Zhang XL, Zhou ZT, Zhang WD (2006) The effect of icariin and astragalosid I on the proliferation and differentiation of bone marrow stromal cells. Zhong Yao Cai 29: 1062–1065.
- Lv L, Wu SY, Wang GF, Zhang JJ, Pang JX, Liu ZQ, Xu W, Wu SG, Rao JJ (2010) Effect of astragaloside IV on hepatic glucose-regulating enzymes in diabetic mice induced by a high-fat diet and streptozotocin. Phytother Res 24: 219–224.
- Motomura K, Fujiwara Y, Kiyota N, Tsurushima K, Takeya M, Nohara T, Nagai R, Ikeda T (2009) Astragalosides isolated from the root of *Astragalus radix* inhibit the formation of advanced glycation end products. J Agric Food Chem 57: 7666–7672.
- Nagatomi J, Arulanandam BP, Metzger DW, Meunier A, Bizios R (2003) Cyclic pressure affects osteoblast functions pertinent to osteogenesis. Ann Biomed Eng 31: 917–923.
- Qi H, Wei L, Han Y, Zhang Q, Lau A. S, Rong J (2010) Proteomic characterization of the cellular response to chemopreventive triterpenoid astragaloside IV in human hepatocellular carcinoma cell line HepG2. Int J Oncol 36: 725–735.
- Qiu LH, Xie XJ, Zhang BQ (2010) Astragaloside IV improves homocysteine-induced acute phase endothelial dysfunction via antioxidant. Biol Pharm Bull 33: 641–646.
- Qu YZ, Li M, Zhao YL, Zhao ZW, Wei XY, Liu JP, Gao L, Gao GD (2009) Astragaloside IV attenuates cerebral ischemia-reperfusion-induced increase in permeability of the blood-brain barrier in rats. Eur J Pharmacol 606: 137–141.
- Robling AG, Castillo AB, Turner CH (2006) Biomechanical and molecular regulation of bone remodeling. Annu Rev Biomed Eng 8: 455–498 (2006).
- Tan YF, Yin XC, Xiong YJ, Wang Y (2010) Stem cell factor secretion by bone mesenchymal stem cells stimulated with astragaloside IV. Zhongguo Dang Dai Er Ke Za Zhi 12: 290–292.
- Wang S, Li J, Huang H, Gao W, Zhuang C, Li B, Zhou P, Kong D (2009) Anti-hepatitis B virus activities of astragaloside IV isolated from *radix Astragali*. Biol Pharm Bull 32: 132–135.
- Wong RW, Rabie B (2008) Chinese red yeast rice (*Monascus purpureus*-fermented rice) promotes bone formation. Chin Med 3: 4.
- Xu XL, Chen XJ, Ji H, Li P, Bian YY, Yang D, Xu JD, Bian ZP, Zhang JN (2008) Astragaloside IV improved intracellular calcium handling in hypoxia-reoxygenated cardiomyocytes via the sarcoplasmic reticulum Ca-ATPase. Pharmacology 81: 325–332.
- Yu J, Zhang Y, Sun S, Shen J, Qiu J, Yin X, Yin H, Jiang S (2006) Inhibitory effects of astragaloside IV on diabetic peripheral neuropathy in rats. Can J Physiol Pharmacol 84: 579–587.
- Zhang C, Wang XH, Zhong MF, Liu RH, Li HL, Zhang WD, Chen H (2007) Mechanisms underlying vasorelaxant action of astragaloside IV in isolated rat aortic rings. Clin Exp Pharmacol Physiol 34: 387–392.
- Zhang JQ, Liu XH, Zhang N, Liu M (2008) Studies on the quality control of compound Qishe Tablets. Lishizhen Med Mater Med Res 19: 977–978.
- Zhang W, Li W, Li W, Sun X, Zhou S, Xu X (2010) Study on preventative and curative effects of astragaloside (AST) on mice memory impairment and expression of amyloid precursor protein and beta secretase mRNA induced by dexamethasone. Zhongguo Zhong Yao Za Zhi 35: 642–646.
- Zhang WD, Chen H, Zhang C, Liu RH, Li HL, Chen HZ (2006) Astragaloside IV from *Astragalus membranaceus* shows cardioprotection during myocardial ischemia *in vivo* and *in vitro*. Planta Med 72: 4–8.
- Zhang WD, Zhang C, Wang XH, Gao PJ, Zhu DL, Chen H, Liu RH, Li HL (2006) Astragaloside IV dilates aortic vessels from normal and spontaneously hypertensive rats through endothelium-dependent and endothelium-independent ways. Planta Med 72: 621–626.
- Zhang Y, Zhu H, Huang C, Cui X, Gao Y, Huang Y, Gong W, Zhao Y, Guo S (2006) Astragaloside IV exerts antiviral effects against coxsackievirus B3 by upregulating interferon-gamma. J Cardiovasc Pharmacol 47: 190–195.
- Zhao Z, Wang W, Wang F, Zhao K, Han Y, Xu W, Tang L (2009) Effects of Astragaloside IV on heart failure in rats. Chin Med 4: 6.
- Zhu SQ, Qi L, Rui YF, Li RX, He XP, Xie ZP (2008) Astragaloside IV inhibits spontaneous synaptic transmission and synchronized Ca²⁺ oscillations on hippocampal neurons. Acta Pharmacol Sin 29: 57–64 (2008).