

DNA STRAND EXCHANGE PROTEINS: A BIOCHEMICAL AND PHYSICAL COMPARISON

Piero R. Bianco ¹, Robert B. Tracy ^{1,2,3} and Stephen C. Kowalczykowski ^{1,2}

¹Sections of Microbiology and of Molecular and Cellular Biology, ²Microbiology Graduate Group, University of California, Davis, CA 95616, ³ Norris Comprehensive Cancer Center, USC School of Medicine, 1441 Eastlake Ave., Los Angeles, CA 90033

Received 5/24/98 Accepted 6/11/98

TABLE OF CONTENTS

1. Abstract	
2. Introduction	
3. Biochemical models for homologous recombination	
3.1. Introduction	
3.1.1. <i>Escherichia coli</i>	
3.1.2. Bacteriophage T4	
3.1.3. <i>S. cerevisiae</i>	
3.2. Initiation	
3.2.1. <i>Escherichia coli</i>	
3.2.2. Bacteriophage T4	
3.2.3. <i>S. cerevisiae</i>	
3.3. Homologous pairing and DNA strand exchange	
3.3.1. <i>Escherichia coli</i>	
3.3.2. Bacteriophage T4	
3.3.3. <i>S. cerevisiae</i>	
3.4. DNA heteroduplex extension	
3.4.1. <i>Escherichia coli</i>	
3.4.2. Bacteriophage T4	
3.4.3. <i>S. cerevisiae</i>	
3.5. Resolution	
3.5.1. <i>Escherichia coli</i>	
3.5.2. Bacteriophage T4	
3.5.3. <i>S. cerevisiae</i>	
4. Mechanism of DNA strand exchange	
4.1. Presynapsis (figure 3, step 1)	
4.1.1. <i>Escherichia coli</i>	
4.1.2. Bacteriophage T4	
4.1.3. <i>S. cerevisiae</i>	
4.2. Synapsis (figure 3, steps 2, 3)	
4.2.1. <i>Escherichia coli</i>	
4.2.2. Bacteriophage T4	
4.2.3. <i>S. cerevisiae</i>	
4.3. Branch migration (figure 3, step 4)	
4.3.1. <i>Escherichia coli</i>	
4.3.2. Bacteriophage T4	
4.3.3. <i>S. cerevisiae</i>	
4.4. ATP hydrolysis in DNA strand exchange	
4.4.1. <i>Escherichia coli</i>	
4.4.2. Bacteriophage T4	
4.4.3. <i>S. cerevisiae</i>	
5. Structural comparison	
5.1. Filament structure	
5.1.1. <i>Escherichia coli</i>	
5.1.2. Bacteriophage T4	
5.1.3. <i>S. cerevisiae</i>	
5.2. Primary amino acid sequence and crystal structure	
5.2.1. <i>Escherichia coli</i>	
5.2.2. Bacteriophage T4	
5.2.3. <i>S. cerevisiae</i>	
6. Perspective	
7. Acknowledgments	
8. References	

Table 1A. Proteins Involved in Genetic Recombination

General Function	ORGANISM		
	<i>E. coli</i>	Bacteriophage T4	<i>S. cerevisiae</i>
Initiation	RecBCD, RecQ	gp46, gp47, gp41, gp59	Mre11, Rad50, Xrs2, Spo11
DNA strand exchange	RecA	UvsX	Rad51
ssDNA-binding	SSB	gp32	RPA
Accessory	RecF, RecO, RecR	UvsY	Rad52, Rad55, Rad57, Rad59
Branch migration	RuvA, RuvB RecG	gp41, gp59 UvsW Dda	Rad54
Holliday Junction cleavage	RuvC	gp49	-
Other proteins	DNA topoisomerase I and II	DNA topoisomerases	DNA topoisomerases
	DNA ligase	DNA ligase	DNA ligases
	DNA polymerase I	DNA polymerases	DNA polymerases

1. ABSTRACT

Homologous genetic recombination is an essential biological process that involves the pairing and exchange of DNA between two homologous chromosomes or DNA molecules. It is of fundamental importance to the preservation of genomic integrity, the production of genetic diversity, and the proper segregation of chromosomes. In *Escherichia coli*, the RecA protein is essential to recombination, and biochemical analysis demonstrates that it is responsible for the crucial steps of homologous pairing and DNA strand exchange. The presence of RecA-like proteins, or their functional equivalents, in bacteriophage, other eubacteria, archaea, and eukaryotes, confirms that the mechanism of homologous pairing and DNA strand exchange is conserved throughout all forms of life. This review focuses on the biochemical and physical characteristics of DNA strand exchange proteins from three diverse organisms: RecA protein from *E. coli*, UvsX protein from bacteriophage T4, and Rad51 protein from *Saccharomyces cerevisiae*.

2. INTRODUCTION

General genetic recombination involves the exchange of homologous regions between two chromosomes or double-stranded DNA (dsDNA) molecules. The resulting, recombinant DNA contains genetic information originally present in each of the parental molecules. Genetic studies demonstrated that, in *Escherichia coli*, there are several recombination pathways (1-4), with many proteins involved in the process (tables 1A-B) (5-7). One of these, the RecA protein, is essential to the primary pathways of homologous genetic recombination. The function of RecA protein is conserved from bacteriophage to humans and, thus, the focus of this review is to compare two other RecA protein homologues, the UvsX protein of bacteriophage T4, and the Rad51 protein of *Saccharomyces cerevisiae*, with the prototypic and the most well studied protein, the RecA protein of *E. coli*.

Clark and Margulies discovered the *recA* gene of *E. coli* by virtue of the strong effects of *recA* mutations on conjugal recombination (8). Subsequent to this discovery, the RecA protein was shown to be essential for homologous genetic recombination and for induction, following DNA damage, of both prophage and the SOS response (9-11). Included in the SOS response are LexA and phage repressor cleavage, amplification of RecA protein levels, induction of a

mutagenic mode of DNA repair, filamentation, prophage induction, and Weigle reactivation (10,12). Not surprisingly, mutations in *recA* are pleiotropic, affecting not only recombination, but also DNA repair, mutagenesis, and cell division.

In vitro, the RecA protein (M_r 37,842) promotes the ATP- and single-stranded DNA (ssDNA)-dependent cleavage of the LexA repressor (13), which results in induction of the SOS response. In addition, it performs two reactions important to homologous recombination: ATP-stimulated DNA strand annealing (renaturation) between complementary single strands of DNA (14-17), and ATP-dependent DNA strand invasion (exchange) between ssDNA and a homologous sequence within dsDNA (18-26). Over the last 18 years, RecA protein has been the subject of intense study (relevant activities are listed in table 2), and as a result, has become the prototypic DNA strand exchange protein to which all other DNA strand annealing and DNA strand exchange proteins are compared. Homologous proteins have been identified in a variety of other organisms including bacteriophage, UvsX protein of T4 phage (27), at least 60 other bacteria (28-30), all eukaryotes examined (31), and, finally, the Archaea (32,33). Thus, RecA protein homologues comprise a family of universally conserved, homologous recombination proteins.

As stated above, a member of this family is the T4 phage-encoded UvsX protein. Harm first identified *UvsX* as a gene that controls UV-sensitivity in bacteriophage T4 (34). Subsequent to this work, a number of studies revealed that mutations in *UvsX* are remarkably diverse in their effect. Mutants which are defective in *UvsX*, 1) are recombination deficient *in vivo* (34-37); 2) have decreased mutagenesis (38); 3) display discontinuous T4 DNA replication *in vivo* (37); and 4) are deficient in repair of DNA damage caused by various chemical and physical agents (39). In addition, lethal mutations in *gene 49* are suppressed by mutation of the *UvsX* gene (37,40). *Gene 49* encodes an endonuclease that resolves recombination intermediates (branched DNA) into linear duplex DNA which can then be packaged into the phage capsid (41,42). *UvsX* is not absolutely required for T4 growth, however, since T4 phage has at least one additional recombination pathway which does not require the involvement of *UvsX* (see reference (43) for review).

DNA strand exchange proteins

Table 1B. Functions of Proteins Involved in Genetic Recombination

GENERAL FUNCTION	PROTEIN	BIOCHEMICAL FUNCTIONS
<i>E. coli</i>		
Initiating protein(s)	RecBCD	ATP-dependent dsDNA and ssDNA exonuclease, ATP-stimulated ssDNA endonuclease, DNA helicase, recombination hotspot χ -recognition
DNA strand exchange	RecA	DNA-dependent ATPase, DNA- and ATP-dependent coprotease, DNA renaturation, DNA strand exchange
ssDNA-binding protein	SSB	ssDNA binding, stimulates DNA strand exchange
Accessory protein(s)	RecF	ssDNA, dsDNA binding, weak ATPase, interacts with RecR protein
	RecO	ssDNA, dsDNA binding; interacts with RecR; RecOR prevents end-dependent disassembly of RecA filaments; interacts with SSB
Branch migration	RecR	Interacts with RecF; RecFR complex attenuates RecA filament extension into dsDNA regions
	RecG	DNA helicase, branch migration of Holliday junctions
	RuvA	Binds to Holliday-, cruciform-, and four-way junctions, interacts with RuvB protein
	RuvB	DNA helicase, branch migration of Holliday junctions, interacts with RuvA protein
Holliday Junction cleavage	RuvC	Binds to four-way junctions, cleaves Holliday junctions
Other proteins	DNA gyrase	Type II topoisomerase
	DNA topoisomerase I	Type I topoisomerase, ω protein
	DNA ligase	DNA ligase
	DNA polymerase I	DNA polymerase, 5'→3' exonuclease, 3'→5' exonuclease
Bacteriophage T4		
Initiating protein(s)	gp46	Interacts with gp47; endo- and exonuclease
	gp47	Interacts with gp46; endo- and exonuclease; stimulates gp46 action
	gp41	DNA-dependent NTPase, ssDNA binding, ATP- or GTP-dependent DNA helicase
	gp59	ssDNA binding, stimulates ATPase and helicase activities of gp41, interacts with gp32 and gp41
DNA strand exchange	UvsX	DNA-dependent ATPase, DNA renaturation, DNA strand exchange
ssDNA-binding protein	gp32	ssDNA binding, stimulates DNA strand exchange, interacts with UvsY and UvsX
Accessory protein(s)	UvsY	Stimulates DNA strand exchange, interacts with UvsX and gp32
Branch migration	Dda	DNA helicase, stimulates branch migration by UvsX protein
	UvsW	DNA helicase, branch migration of Holliday junctions, functional analogue of RecG
	gp41	DNA-dependent NTPase, ssDNA binding, ATP- or GTP-dependent DNA helicase
	gp59	ssDNA binding, stimulates ATPase and helicase activities of gp41, interacts with gp32 and gp41
Holliday Junction cleavage	gp49	Binds to and cleaves Y-junctions and Holliday junctions
<i>S. cerevisiae</i>		
Initiating protein(s)	Mre11	Forms complex with Rad50 and Xrs2 which is possibly responsible for resection of double-strand DNA breaks; with Rad50, ssDNA endo- and 3' to 5' dsDNA exonuclease
	Rad50	Forms complex with Mre11 and Xrs2 which is possibly responsible for resection of double-strand DNA breaks, ATP-dependent binding to dsDNA, contains ATP-binding Motif
	Xrs2	Forms complex with Mre11 and Rad50 which is possibly responsible for resection of double-strand DNA breaks
DNA strand exchange	Spo11	Binds DNA, likely catalytic subunit responsible for double-strand break formation
ssDNA-binding protein	Rad51	DNA-dependent ATPase, DNA strand exchange, interacts with Rad52, Rad54, and Rad55 proteins
Accessory protein(s)	RPA	ssDNA binding, stimulates DNA strand exchange, interacts with Rad52 protein
	Rad52	Stimulates DNA strand exchange, interacts with Rad51 and RPA proteins
	Rad54	Contains both ATP-binding, DNA helicase motifs, interacts with Rad51, hydrolyzes ATP, stimulates DNA strand exchange
	Rad55	Stimulates DNA strand exchange, contains Walker ATP-binding motif, interacts with Rad51 protein, forms stable heterodimer with Rad57 protein; shows homology to Rad51
	Rad57	Stimulates DNA strand exchange, contains Walker ATP-binding motif, forms stable heterodimer with Rad55 protein; shows homology to Rad51
Other proteins	Rad59	Shows homology to Rad52 protein, function is unknown

Table 2. Biochemical Activities of the Prototypic DNA Strand Exchange Proteins

	Activity	RecA protein	UvsX protein	Rad51 protein
ATPase	k_{cat} (min ⁻¹) on ssDNA	18 - 30	145 - 240	0.6 - 0.7
	k_{cat} (min ⁻¹) on dsDNA	19 - 22	5	0.05 - 0.1
	Length dependent	Yes	Yes	-
	Compositional preference	Yes	-	Yes
	Cooperative	Yes	Yes	Yes
	Reaction Products	ADP+P _i	ADP+P _i , AMP+PP _i	ADP+P _i
	Stimulated by	SSB	UvsY; gp32	RPA; Rad52
DNA Binding	ssDNA			
	Stoichiometry (nt:monomer)	3 - 6	3 - 5	3 - 6
	Compositional Preference	Yes (GT>AC)	-	Yes (GT>AC)
	Length dependence	≥ 9 nt	≥ 16 nt	-
	Cooperative	Yes	Yes	Yes
	Stimulated by	SSB	gp32, UvsY	RPA
	Cofactor dependent	No	No	No
	Stabilized by NTP	Yes	Yes (ATPyS)	Yes
	dsDNA			
	Stoichiometry (bp:monomer)	3	3	3
DNA Strand Exchange	pH-dependent	Yes	Not determined	Yes
	Cooperative	Yes	Yes	Yes
	Cofactor required	Yes	Yes	Yes
	Inhibited by Excess dsDNA	No	Yes	Yes
	ATP-binding required	Yes	Yes	Yes
	ATP-hydrolysis essential	No/Yes	Yes	No
	Supported by other NTPs	Yes	No	No
	Coaggregation	Yes	Yes	-
	Polarity	5' to 3'	5' to 3'	3' to 5'
	Stimulated by	SSB; RecF,O,R	gp32, UvsY	RPA; Rad52, 54, 55, 57

UvsX protein (M_r 43,760) has many of the same biochemical activities possessed by RecA protein (table 2): 1) ssDNA-dependent ATP hydrolysis activity (44,45); 2) binding to both ssDNA and dsDNA, with a preference for ssDNA (46,47); 3) formation of a nucleoprotein filament (46-48); 4) ATP-dependent renaturation of complementary ssDNA (45); and 5) ATP-dependent homologous pairing and exchange of complementary DNA strands (44,45,49,50), which, under the appropriate conditions, is stimulated by the two accessory factors, UvsY protein and gene 32 protein (gp32) (47,51).

The third member of the RecA-like proteins that will be discussed in this review is the eukaryotic Rad51 protein. Genetic analyses of the budding yeast *S. cerevisiae* identified many mutants deficient in homologous recombination and DNA double-strand break repair (DSBR). The genes defined by these mutations belong to the RAD52 epistasis group. These mutants were isolated on the basis of their sensitivity to ionizing radiation and sensitivity to methyl methanesulfonate (MMS), and they display a characteristic array of pleiotropic phenotypes (52-54). Mutant strains are defective in meiosis, as evidenced by their inability to produce viable spores (54,55). Mutations in the RAD51, RAD52, and RAD54 genes exhibit additional defects in mitotic recombination and also in mating type switching (54-56). In RAD51, RAD52, RAD55, and RAD57 mutant strains, DNA double-strand breaks (DSBs) with processed tails, which are an early recombination intermediate in *S. cerevisiae* (57,58), form readily but are not further processed to produce viable recombinants (56,59). Thus, subsequent steps at least require these gene products (table 1). Genetic characterization of the RAD52 epistasis group has identified RAD51 as an important member of this group (52,56). The product of the RAD51

gene, Rad51 protein, shows both sequence as well as structural homology to the RecA protein of *E. coli* (59-61). *In vitro*, the Rad51 protein from either *S. cerevisiae* (M_r 42,961; purified from *E. coli* or *S. cerevisiae*) (62-65) or humans (M_r 37,000; purified from *E. coli*) (66,67), possesses those activities which establish it as the eukaryotic homologue of the bacterial RecA protein (table 2). Based on sequence homology at the amino acid level, another RecA-like protein, the Dmc1 protein, is present in yeast (57,58,68). Though potentially another DNA strand exchange protein that is ubiquitous and which functions in meiosis, no biochemical characterization exists, and hence it will not be discussed herein. Like the RecA and UvsX proteins, Rad51 protein functions in concert with a number of accessory proteins that augment its limited DNA strand exchange activity (62,65,69-71).

3. BIOCHEMICAL MODELS FOR HOMOLOGOUS RECOMBINATION

3.1. Introduction

Biochemical mechanisms for homologous recombination can be separated into four distinct stages: initiation, homologous pairing and DNA strand exchange, DNA heteroduplex extension (branch migration), and resolution. Figure 1 depicts a generic model for homologous recombination. This model, the DNA double-strand break repair (DSBR) model (72,73), can be applied to mechanisms of homologous recombination in all organisms examined, since in most cases a DNA double-strand break (DSB) is necessary for initiation of the process (though similar models, involving initiation in ssDNA gaps, can also apply (see below)). The specific steps of this model are as follows. In




Figure 1. General model for homologous recombination. The model depicted is the double-strand break repair (DSBR) model (72,73). Since a double-strand break is involved in most initiation events, this model, in its most general form, likely can apply to mechanisms of homologous recombination in all organisms. Details are discussed in the text. Light blue lines indicate DNA that has been recently synthesized by a DNA polymerase.

step 1, a DSB is introduced into one of the DNA substrates. The formation of this DSB could be due to a programmed *in vivo* process or to environmental circumstances. More specific examples of these will be discussed below. In step 2, the DSB is processed by either a helicase, a nuclease, or a combination of both, to yield a free 3'-ssDNA overhang(s). In step 3, this free 3'-terminal end is homologously paired with its homologue by a RecA-like protein to form an intermediate known as a joint molecule. Following joint molecule formation, pairing of the displaced complementary strands results in the formation of characteristic intermediates known as Holliday junctions (step 4). In step 5, the Holliday junction(s) is extended unidirectionally by branch migration proteins. Finally, in step 6, the Holliday junction is cleaved by a resolvase to yield recombinant products (*e.g.* spliced or patched). Given some minor differences, each of these steps occurs in the major pathways of homologous recombination in *E. coli*, bacteriophage T4, and *S. cerevisiae* (figure 2). The key players in each of these pathways and their respective functions are listed in table 1.

3.1.1. *Escherichia coli*

In *E. coli*, recombination is readily observed following conjugation or transduction. As part of either

process, the DNA is transiently linearized; consequently, most homologous recombination events are thought to initiate at the ends of linear dsDNA (74). In both instances, DNA strand invasion occurs between linear dsDNA (donor) and supercoiled DNA (recipient). A model for the major pathway (the RecBCD-pathway) of genetic recombination in *E. coli* is displayed in figure 2A. Participation of RecA protein in this pathway, as well as most other pathways (4), is absolutely essential as demonstrated by the fact that both conjugation (9) and transduction (75) are reduced by as much as 50,000-fold in *recA* mutant strains.

3.1.2 Bacteriophage T4

At least three major recombination pathways are described for T4 phage. Each pathway is distinguished by the structure of the DNA substrates that enter the recombination process. Two of the pathways occur in only certain regions of the T4 phage genome and are linked to DNA replication (43). The third pathway is the idealized general recombination reaction, which takes place in the absence of DNA replication, and occurs throughout the genome. A model for this pathway is depicted in figure 2B.

3.1.3. *S. cerevisiae*

The genes in the *RAD52* epistasis group of *S. cerevisiae* are required for homologous recombination and for the recombinational repair of DSBs (52-54). Though there are at least three separate *RAD52*-dependent recombination pathways (76), the major pathway for recombination in yeast defines the canonical DSBR model (figure 2C) (72,73). The *RAD51*-dependent pathway is responsible for the majority (95%) of recombination events; this pathway requires the *RAD51*, *RAD52*, *RAD54*, *RAD55*, and *RAD57* genes (76). A functional *RAD52* gene product is absolutely necessary in all pathways, since *rad52* mutants are significantly reduced for recombination (>3,000-fold) (77). It is unclear exactly where the Rad52 protein functions in the pathway but it appears to act after the Rad50 protein (78), and with the Rad51 protein (70,71,79).

3.2. Initiation

Generally, homologous recombination is initiated at a dsDNA end. Subsequently, the linear dsDNA is processed by organism-specific proteins to produce ssDNA suitable for a RecA-like protein function. A dsDNA end can be formed in two general ways: 1) *via* environmental factors (*e.g.* UV light, free radicals, γ -rays, X-rays, and chemical mutagens), and 2) *via* intended *in vivo* processes (for example, in *E. coli*, conjugation and transduction produce dsDNA ends; in *S. cerevisiae*, a protein (Spo11) cleaves DNA to yield dsDNA ends (80,81)). For all of the organisms discussed here, a dsDNA end can also potentially form when the replication machinery encounters a single-strand interruption in the template DNA, causing a collapse of the replication fork (82). The subsequent repair of the interrupted replication fork will then take place *via* the homologous recombination mechanisms discussed below.

3.2.1. *Escherichia coli*

In *E. coli*, formation of the DNA break is a consequence of conjugation, transduction, or direct damage to the DNA. RecBCD enzyme initiates recombination by acting




Figure 2. Biochemical models for homologous recombination. (A) *Escherichia coli*; (B) Bacteriophage T4; (C) *Saccharomyces cerevisiae*. Details are discussed in the text. Dashed lines indicate DNA degraded by a nuclease/helicase and light blue lines indicate DNA recently synthesized by DNA polymerase. The model in panel A is modified from reference (306). The model in panel C is modified from reference (307).

on the break to create a ssDNA substrate suitable for RecA protein activity (4,83,84). The RecBCD enzyme is a combination nuclease/helicase that binds to the end of linear dsDNA, initiates unwinding of the DNA, and simultaneously degrades preferentially the 3'-terminal strand of the DNA (85,86). These concurrent activities continue until RecBCD enzyme interacts with a DNA sequence (5'-GCTGGTGG-3') (87,88) known as Chi (χ ; crossover hotspot instigator (89)). At χ , the nuclease activity of the enzyme is altered so that degradation of the 3'-terminal strand is down-regulated and the nuclease activity of the 5'-terminated strand is up-regulated, but the helicase activity remains unaltered (85,90,91). The effect of this χ -mediated modification is a recombinationally active RecBCD enzyme that retains helicase activity, but has a diminished nuclease activity with an opposite (5'→3') polarity of degradation. After unwinding and degradation, the χ -modified RecBCD enzyme produces resected dsDNA with a 3'-terminal ssDNA tail terminating at the χ sequence (86,88,90-93). During translocation and unwinding, RecBCD enzyme loads RecA protein onto this χ -containing ssDNA tail (94) which, in fact, is the preferred substrate for RecA protein-dependent strand invasion of χ -containing supercoiled DNA (85,90,94,95). The single-stranded DNA binding (SSB) protein also plays an important

role at this step by trapping the unwound ssDNA, preventing it from reannealing and protecting it from other nucleases. Recently, it was shown that another helicase, the RecQ helicase, is also able to function in the initiation of recombination by providing RecA protein with a suitable DNA substrate (96).

3.2.2. Bacteriophage T4

The initiation step in T4 phage recombination also requires processing of at least one DNA partner to produce ssDNA appropriate for UvsX protein function. The recombinogenic ssDNA is believed to be generated *via* cooperation between two reactions. Genetic data suggested that the gene 46 protein (gp46) and gene 47 protein (gp47) form an exonuclease which creates a ssDNA gap in a substrate containing a nick (97); recent *in vitro* results demonstrate that gp46 is both an endo- and an exonuclease, and that these activities are stimulated by gp47 (S. Morrical, personal communication). Subsequently, the gapped substrate is acted upon by a 5'→3' helicase, the gene 41 protein (gp41) (98), and its enhancer, the gene 59 protein (gp59) (99), to produce a free 3'-end that is used by UvsX protein for DNA strand invasion. Gp32 (the analogue of *E. coli* SSB protein) also functions in this unwinding reaction, since it

appears to form a protein complex with gp41 and gp59 (99,100).

3.2.3. *S. cerevisiae*

In the budding yeast *S. cerevisiae*, homologous recombination commonly initiates at DSBs (101). DSBs are assumed to occur spontaneously in mitotically growing cells, but in meiosis the DSBs may be programmed, and they are associated with meiotic recombination hotspot activity (57,102-104). The current view is that DSBs are a consequence of cleavage by a nonspecific endonuclease at sites in the chromatin where the DNA is readily accessible (101,103,105,106). Formation of the DSBs requires at least the products of the *RAD50* and *SPO11* genes (table 1), since their formation is abolished by null mutations in either of these genes (57,107). Recently, it was shown that the Spo11 protein is likely to be the catalytic subunit responsible for formation of the DSBs (80,81). Also, it has become clear that additional gene products are involved in DSB formation (e.g. *MEI4* (108); *MER2* (109); *MRE11* (110,111); *REC102*, *REC104*, *REC114* (112); and *XRS2* (113,114)), since null mutations in any of these genes prevents DSB formation. Following DSB formation, 3'-single-stranded tails of about 600 nucleotides are produced via 5'→3' exonuclease activity (57,58,68). Resection of the 5'-strand termini at DSB's appears to require at least the Rad50, Mre11, and Xrs2 proteins since mutations in the corresponding genes result in cells that cannot process DSBs (106,107,114-116). Recently, all three of these proteins were shown to form a complex with one another (111), further supporting the possibility that they act together to form the nuclease responsible for resection of the DSB's.

3.3. Homologous pairing and DNA strand exchange

Following the production of a free 3'-ssDNA overhang, the following steps occur: 1) the RecA-like protein cooperatively binds to the free 3'-ssDNA end to form a nucleoprotein filament; 2) a quick and efficient search for homology occurs within the dsDNA recipient that results in the formation of a joint molecule; and 3) once the two DNA molecules are homologously aligned, the RecA-like protein then catalyzes the nascent exchange of strands to produce a Holliday junction.

3.3.1. *Escherichia coli*

In *E. coli*, the RecA protein binds ATP, and then in combination with SSB protein, forms a continuous presynaptic filament on the ssDNA (a filament formed on dsDNA is shown in figure 5; discussed in Section V). The presence of SSB protein permits the formation of a continuous filament of RecA protein by destabilizing secondary structure (117-122). This nucleoprotein complex acts as the functional unit to perform a rapid and efficient search for homology within the dsDNA recipient, resulting in the formation of a joint molecule or D-loop. In the presence of SSB protein, the RecF, RecO, and RecR proteins (table 1), stimulate formation of the joint molecules, by helping RecA protein overcome the inhibition of binding by the SSB protein (123,124). The 3'-terminal end is more invasive in RecA protein-dependent pairing reactions because RecA protein polymerizes along ssDNA in a 5'→3' direction; thus, the 3'-end is more likely to be covered (125,126). Following joint molecule formation, RecA protein, perhaps with the aid of

other proteins (for example, DNA polymerase I and/or a topoisomerase, table 1), catalyzes pairing of the remaining two DNA strands to produce a Holliday junction (for further details on the potential roles of DNA polymerase I and topoisomerases in homologous recombination, see reference (4)).

3.3.2. Bacteriophage T4

In T4 phage, UvsX protein forms a nucleoprotein complex with ssDNA (a filament formed on dsDNA is shown in figure 5). This nucleoprotein filament subsequently pairs with dsDNA in the presence of ATP (44,45). Homologous pairing is stimulated by the presence of the accessory factors, UvsY protein and gp32 (47,51) (figure 2B). The gp32 protein is envisioned to bind ssDNA first. Then, the UvsY protein binds to the ssDNA and helps load UvsX protein onto the gp32-ssDNA complex by assisting UvsX protein in displacing the bound gp32 (127). UvsY protein possesses two important properties that are important to this process: its intrinsic affinity for ssDNA is greater than that of gp32 (127) and it exists predominantly in hexameric form, the form which binds ssDNA (128). It was proposed that binding of ssDNA by the UvsY hexamer changes the conformation of the ssDNA (perhaps by wrapping the ssDNA around the hexamer). This change either permits nucleation of the UvsX protein presynaptic filament formation and/or destabilization of gp32-ssDNA interactions so that gp32 is more readily displaced by UvsX protein (128). In addition, interactions between UvsY protein, gp32 and UvsX protein are important, but not essential, for optimal UvsX activity (129).

Once UvsX protein begins to cooperatively bind to the ssDNA, gp32 is gradually displaced. Following homologous pairing, joint molecules are converted to crossover intermediates (Holliday junctions) when UvsX protein exchanges strands between the two homologously paired substrates (44,45,49). The interactions between the gp32, UvsX, and UvsY proteins are a prototype for those between the SSB, RecA, and RecFOR proteins described above for *E. coli*, and for the interactions between the heterotrimeric replication protein A (RPA), Rad51, Rad52, Rad55, and Rad57 proteins described below for *S. cerevisiae*.

3.3.3. *S. cerevisiae*

It is becoming clear that a number of proteins are involved at this step in *S. cerevisiae* (table 1 and figure 2C). Less obvious, however, is the precise role for each protein in this process. The yeast Rad51 protein has considerable homology to the *E. coli* RecA protein (59-61) and, recently, it was demonstrated that Rad51 protein, along with yeast RPA (the yeast ssDNA-binding protein), can promote DNA strand exchange *in vitro* (62-65). Similar to the RecA and UvsX proteins, Rad51 protein forms a nucleoprotein filament, which is the active species in DNA strand exchange (a filament formed on dsDNA is shown in figure 5; discussed in Section V). Therefore, it is likely that *RAD51* functions at this step of recombination. However, also important to this step, are the products of the *RAD52*, *RAD54*, *RAD55*, *RAD57*, and *RAD59* genes. Using a chromosomal inverted-repeat assay to study mitotic recombination in haploid yeast cells, Symington and coworkers showed that mutations in these genes reduce recombination efficiency by >3,000-, 25-, 20-, 33-, and 5-fold, respectively (76,77,130). In addition, the products of the

RAD51 and *RAD59* genes appear to act synergistically, since double mutants reduce recombination by 1200-fold (130). Finally, mutations in the gene of the large subunit of RPA (RPA1) produce a defective recombination phenotype (131,132). Recent *in vitro* studies begin to define the specific function of these accessory proteins: RPA stimulates formation of the Rad51 nucleoprotein filaments (65); Rad52 protein, in combination with RPA, stimulates DNA strand exchange by the Rad51 protein (70,71,133); and, a heterodimer composed of the Rad55 and Rad57 proteins also stimulates Rad51-protein dependent homologous pairing, particularly when RPA must be displaced from the ssDNA (69). Finally, Rad54 protein interacts directly with the Rad51 protein and, in the presence of RPA, produces a significant stimulation in the Rad51 protein-dependent pairing rate, to a level comparable to that of the RecA and UvsX proteins (134).

Several studies provide direct evidence for interactions between a number of these proteins. First, the Rad51 and Rad52 proteins interact both *in vitro* (59) and *in vivo* (135). Second, interactions between the Rad51 and Rad55 proteins, and between the Rad55 and Rad57 proteins, were demonstrated (69,136). Third, there is now both *in vivo* and *in vitro* evidence that the Rad51 and Rad54 proteins directly interact (134,137,138). Finally, there is genetic evidence that the large subunit of RPA interacts with Rad52 protein (131,132). Existence of these interactions lead to the proposal that a complex of these proteins (termed a "recombinosome" (136) or "protein machine" (137)) is responsible for enacting homologous pairing and DNA strand exchange; though, it is also possible that there is not a stable complex *per se*, but rather a sequential progression of interactions between groups of proteins and/or individual proteins, similar to that described for the T4 system.

Though the manner by which they are formed and resolved is unclear, recently, double Holliday junctions were identified as the *in vivo* products of the pairing stage (139), lending additional support for the double-strand break repair model proposed earlier (see figures 1 and 2), since the model predicts such structures as intermediates (72,73).

3.4. DNA heteroduplex extension

Subsequent to Holliday junction formation, the nascent region of DNA heteroduplex undergoes branch migration. Although each of the RecA-like proteins discussed in this review can catalyze DNA heteroduplex extension, it is now appreciated that a class of specialized DNA helicases is more efficient than the RecA-like proteins in catalyzing this reaction (140-143). Therefore, it is possible that the RecA-like proteins play a limited role in branch migration.

3.4.1. *Escherichia coli*

Figure 2A shows that in *E. coli* there are several alternative means to extend the heteroduplex DNA. First, RecA protein has branch migration activity which is unidirectional and proceeds at 2-10 bp/sec (18). Second, a complex of the RuvA and RuvB proteins comprises a unique motor protein that also has branch migration activity (144,145), but which is approximately five-fold faster than that of the RecA protein (146,147). However, in contrast to

RecA protein, the RuvAB proteins catalyze branch migration in either direction (146). Finally, the RecG protein, which is both a DNA and DNA/RNA hybrid helicase, can bind to and migrate Holliday junctions (148,149). Curiously, the RecG protein appears to antagonize the heteroduplex extension activities of RecA protein and the RuvAB proteins (150).

3.4.2. Bacteriophage T4

In T4 phage, DNA heteroduplex extension can be mediated in several different ways (figure 2B). Just as with RecA protein, the UvsX protein itself promotes polar branch migration at a rate of 12-15 bp/sec (45,50). In addition, the Dda protein can stimulate branch migration mediated by UvsX protein (151). The Dda protein is an ATP-dependent DNA helicase (152) which, like the RecG protein, has the ability to both extend and disrupt nascent joint molecules (153). Recently, however, the significance of this stimulation by Dda protein in the physiological process was brought into question when it was demonstrated that UvsX protein-mediated branch migration is strongly inhibited by the UvsY protein under physiological conditions (154). Under these conditions, branch migration is driven by the combined actions of gp41 and gp59, rather than by UvsX protein (154). The gp59 loads gp41 onto the strand exchange complex *via* direct interactions with gp32 (100,155). Finally, recent *in vitro* evidence shows that another protein, UvsW, has the potential to act at this step (156). UvsW protein, partially purified as a GST-fusion protein, displays ssDNA-dependent ATPase activity, and is a DNA helicase that acts on a branched DNA substrate. These observations support for the physiological function of UvsW protein being involved in DNA branch migration.

3.4.3. *S. cerevisiae*

In yeast, the Rad51 protein also possesses polar branch migration activity. Interestingly, heteroduplex extension occurs 3'→5' relative to the displaced strand, which is a polarity opposite to that of the RecA protein (63). If indeed a "recombinosome" exists, then one might expect that at least some of the other proteins are also involved at this step. One candidate is the Rad54 protein, which possesses both ATP binding and DNA helicase motifs (137,138); recently, Rad54 protein was shown to stimulate Rad51 protein-mediated DNA strand exchange (134). For the same reasons, the other candidate is the Rad55/57 heterodimeric complex. Though this complex appears to act at an early step of the recombination process (69), a later function remains a possible, additional role.

3.5 Resolution

The final step of recombination requires resolution of the Holliday junctions *via* nucleolytic cleavage. Symmetric cleavage of the junction produces recombinant progeny that are either patched products (exchange of ssDNA strands between the two starting substrates) or spliced products (exchange of flanking markers) (figure 2).

3.5.1. *Escherichia coli*

In *E. coli*, resolution of the Holliday junctions is performed by the RuvC protein (157,158). Since RuvC protein cleaves Holliday junctions symmetrically, both patched and spliced recombinants are produced. RuvC acts in concert with the RuvAB proteins to coordinate the steps of




Figure 3. Kinetic steps of DNA strand exchange. The reaction between circular ssDNA and linear dsDNA is shown. The product is a nicked circle and a displaced single strand of DNA. A similar reaction scheme likely applies to all of the DNA strand exchange proteins discussed but reaction polarity is not indicated; the illustration is adapted from Kowalczykowski (185). The steps shown are: 1-presynapsis; 2 and 3-synapsis; and 4-DNA heteroduplex extension. The green spheres represent the strand exchange protein; the orange squares represent ssDNA binding proteins.

branch migration and Holliday junction resolution. Recently, the late stages (DNA heteroduplex extension via the RuvA and RuvB proteins, and resolution by the RuvC protein) of recombination in *E. coli* were reconstituted *in vitro* (159).

3.5.2. Bacteriophage T4

The most likely candidate for Holliday junction resolution in T4 phage is endonuclease VII (gp49). This enzyme cleaves branched DNA molecules (41) and four-way Holliday junctions (160). Cleavage of synthetic four-way junctions is symmetrical and is organized around the junction point on opposite sides (161,162). While both T4 endonuclease VII and RuvC protein perform the same activity, there is no strong homology shared between the two proteins (163). This could help explain the difference in the specificity of cleavage of the two proteins. RuvC protein is specific for four-way junctions, while T4 endonuclease VII will cleave not only four-way junctions, but also Y-junctions (164) and heteroduplex loops (165) as well.

3.5.3. *S. cerevisiae*

Resolution of the intermediate with two Holliday junctions in yeast would yield either spliced or patched products depending on how the junctions were cleaved (figure 2C). If the two junctions are cleaved in opposite orientations, this would generate spliced products. However, if they are cut in the same orientation, patched products would be produced. At this time, there is no obvious candidate for this activity. Two partially purified enzymes, termed endo-X1 (166) and endo-X2 (167), were shown to cleave cruciform-containing DNA; however, neither enzyme was purified to homogeneity. Subsequently, another gene, *CCE1* (Cruciform Cutting Endonuclease 1), was identified during a screen of yeast mutants which were unable to cleave cruciform structures (168). The product of the *CCE1* gene was shown to be the same as endo-X2 (168). Though Cce1/endo-X2 displays the ability to cleave Holliday junctions *in vitro*, the activity is mitochondrial, rather than nuclear, in origin (169).

4. MECHANISM OF DNA STRAND EXCHANGE

As is evident from the preceding, RecA protein and its analogues mediate a reaction, DNA strand exchange, that is central to homologous recombination. The enzymatic properties of the *in vitro* strand exchange activity of RecA protein have been studied in great detail (4,28,170) and a summary of the biochemical properties of the DNA strand exchange proteins discussed herein is shown in table 2. Below, we detail the steps of protein-promoted DNA strand exchange.

DNA strand exchange catalyzed by RecA protein occurs by a number of kinetically distinct phases that can be subdivided into at least three experimentally distinguishable steps: (1) presynapsis, (2) synapsis and (3) DNA heteroduplex extension (figure 3). Presynapsis is the step where RecA protein assembles onto the ssDNA to form the nucleoprotein species that is active in the homology search. Synapsis is characterized by initially random non-homologous contacts occurring between the presynaptic complex and naked dsDNA (termed coaggregates), the search for homology, homologous pairing, and finally conversion to either

DNA strand exchange proteins

paranemic or plectonemic joint molecules (discussed below in the synapsis section). DNA heteroduplex extension, as discussed earlier, is the polar migration of the nascent DNA heteroduplex joint.

DNA strand exchange occurs between two DNA molecules provided that the following criteria are met: (1) one of the DNA molecules contains a region of ssDNA; (2) the ssDNA region occurs at a site homologous to the other dsDNA molecule; and (3) for topological reasons, one of the substrate molecules has an end. A reaction that has become a model for mechanistic studies involves the exchange between circular ssDNA (ϕ X174 or M13 phage) and linear dsDNA (figure 3). The products are nicked, circular dsDNA and linear ssDNA (171,172).

4.1 Presynapsis (figure 3, Step 1)

Presynapsis is the ordered assembly of the RecA protein homologue on ssDNA to produce a nucleoprotein complex that is the active species in DNA strand exchange (filaments formed on dsDNA are shown in figure 5 and discussed in Section V). This process occurs by either of two pathways: first, the single-stranded DNA binding protein (SSB; RPA; UvsY and/or gp32) binds to ssDNA (figure 3, step 1A); this is followed by the subsequent binding of the DNA strand exchange protein (RecA; UvsX or Rad51) and displacement of the single-stranded DNA binding protein to form the presynaptic filament (figure 3, steps 1B and 1C, left panel). In the second pathway, the DNA-strand exchange protein binds to the ssDNA substrate, forming an incomplete nucleoprotein filament due to limitations imposed by DNA secondary structure (step 1A, right panel). The single-stranded DNA binding protein binds, removes the secondary structure and is then displaced by further binding of the DNA strand exchange protein (steps 1B and C). The net result of either of these two pathways, is the formation of a complete nucleoprotein filament that involves direct contact by the primary DNA-binding site of the strand exchange protein, and results in the generation of a contiguous secondary DNA-binding site that is essential for homologous pairing and DNA strand exchange (173,174).

4.1.1. *Escherichia coli*

Filament formation by RecA protein is dependent on DNA length, DNA composition, and the amount of secondary structure within the ssDNA substrate (95,175-178). The ATP-bound form of the protein is needed to constitute the active form of the nucleoprotein filament. ATP greatly stabilizes the complex, and transforms RecA protein into a DNA-binding state with high affinity for ssDNA (120). The RecA nucleoprotein filament is capable of hydrolyzing ATP with a k_{cat} of 18-30 min⁻¹, a value that is affected by ssDNA length and sequence (117,177-179). However, DNA serves only as a scaffold to promote filament formation since RecA protein can be activated by high salt concentrations to form both filaments and to hydrolyze ATP (180-182). Though RecA protein is an ATPase, ATP hydrolysis is not essential for DNA strand exchange (64,183,184). Stoichiometric amounts of RecA protein, relative to ssDNA concentrations, are required for filament formation, with maximum DNA strand exchange rates occurring when filaments are formed at a ratio of at least 1 RecA monomer to 3 nucleotides of ssDNA. This stoichiometric requirement indicates the

importance of a saturated, contiguous RecA protein-ssDNA filament in the DNA strand exchange process (185).

In addition to ATP, ATP- γ -S and dATP also support presynaptic filament formation and are capable of transforming RecA protein into the high affinity ssDNA-binding state, the active form of the protein (179,186). This complex, the presynaptic filament, is the physiologically relevant species in RecA protein-mediated cellular processes. Filament assembly onto ssDNA occurs in a 5'→3' direction (187), which is the same direction as branch migration. The resulting filament has a regular structure and the prominent feature is a large helical groove, with 6.2 monomers per helical turn and each monomer interacting with 3 bases of DNA (figure 5) (188).

The SSB protein effects the binding of RecA protein to ssDNA, greatly stimulating the DNA strand exchange process (117,189). The role of SSB protein in presynapsis is to remove secondary structure from ssDNA, which is inhibitory to the formation of the saturated presynaptic complex (117,119-122). This is consistent with the role of the protein as a helix-destabilizing protein and the observation that other helix destabilizing proteins can substitute for SSB protein in DNA strand exchange *in vitro* (119).

4.1.2. Bacteriophage T4

Presynaptic complex formation by the T4 phage proteins is similar, but features an added complexity, the UvsY protein. During presynapsis, UvsX binds to ssDNA cooperatively with a stoichiometry of one UvsX monomer for every 3 to 5 nucleotides of ssDNA to form a nucleoprotein filament (46,47). In contrast to RecA protein, the binding of ATP does not stabilize the UvsX protein-ssDNA presynaptic filament under steady-state conditions for hydrolysis of ATP; however, the binding of ATP- γ -S does (190). This suggests that ATP- γ -S increases the equilibrium binding affinity of UvsX protein for ssDNA which is very similar to what is observed for RecA protein (179,191). However, the UvsX protein-ssDNA complex is more vigorous in hydrolyzing ATP (the rate of ATP hydrolysis is 8 to 10-fold higher) (44), making its presynaptic filament more dynamic; this may explain why ATP does not stabilize the UvsX-presynaptic complex, (*i.e.*, the UvsX-ssDNA-ATP complex is short lived) and why an accessory protein like UvsY protein is needed. In addition, UvsX protein is unique among RecA protein analogues studied to date, in that it produces both ADP + P_i and AMP + PP_i (44). The significance of the AMP production is presently unknown, however.

The formation of the UvsX protein presynaptic filament is facilitated by the presence of both gp32 and UvsY protein (a UvsX-dsDNA nucleoprotein filament is shown in figure 5B and is discussed in Section V). Gp32 enhances filament formation by removing secondary structure native to the ssDNA (45,51). UvsY facilitates filament formation by stabilizing the presynaptic complex itself (51,192,193). Although gp32 is the T4 phage counterpart to the *E. coli* SSB protein, there is no structural homologue of the UvsY protein in the *E. coli* recombination system; however, the RecF, O, and R proteins may comprise a functional homologue (123,124,194-198).

4.1.3. *S. cerevisiae*

Presynaptic filament formation by Rad51 protein results in a stoichiometric nucleoprotein filament that resembles the filament formed by both the RecA and UvsX proteins (figure 5C, discussed in Section V) (63). This complex is the active species in DNA strand exchange (63). Optimal reaction rates are achieved with a stoichiometry of 3 nucleotides per Rad51 monomer (63,199). Binding to ssDNA is stabilized by either ATP (59) or the non-hydrolyzable analogues of ATP, ATP- γ -S and AMP-PNP (64). Rad51 protein filament formation is stimulated by the RPA heterotrimer (63,65), which is the eukaryotic analogue of the *E. coli* SSB and T4 gp32 proteins. In contrast to both the RecA and UvsX proteins, very little ATP is hydrolyzed by the Rad51 nucleoprotein complex ($k_{\text{cat}} = 0.73 \text{ min}^{-1}$) during presynapsis or during DNA strand exchange (64); interestingly, some of the enzymatic characteristics of Rad51 protein more closely parallel those of a mutant RecA protein, RecA K72R, which has a reduced level of ATP hydrolysis ($\sim 600\text{--}850$ -fold reduced to a k_{cat} of 0.032 min^{-1}) yet nonetheless, is able to promote DNA strand exchange.

4.2 Synapsis (figure 3, Steps 2, 3)

Once the presynaptic filament has assembled on ssDNA, synapsis ensues. In this stage of the reaction, a dsDNA molecule must be bound to the filament, homology to the ssDNA within the filament located within the dsDNA, and a plectonemic heteroduplex joint formed. For this phase of DNA strand exchange to occur, a second DNA molecule must bind in a sequence-independent fashion to the secondary DNA binding site of the nucleoprotein filament; presumably, the close approach of this second DNA molecule is facilitated by shielding of the negatively charged phosphodiester backbones of the two DNA molecules by the DNA strand exchange protein. Once dsDNA is bound, the search for homology takes place. The search is rapid, and requires that the binding of dsDNA to the secondary site be both weak and transient. The recognition of homology takes place when the ssDNA within the presynaptic filament hydrogen bonds, *via* non-Watson-Crick base pairing, to either the major or minor groove of the bound dsDNA, in a mechanism that does not require triplex intermediate formation. This alignment provides a signal to the strand exchange protein that homology has been located; base switching occurs, and the heteroduplex dsDNA occupies the primary site whereas the displaced strand occupies the secondary site. If the location of homology occurs in the center of the dsDNA, a paranemic joint forms; if this occurs at the dsDNA molecule ends, a plectonemic joint forms. Due to their inherent instability, paranemic joints must be converted to plectonemic joints in order to survive and become recombinant DNA structures.

4.2.1. *Escherichia coli*

Once an active presynaptic complex forms, the complex can then pair the ssDNA within the filament to dsDNA. On the basis of probability, this initial contact must be at a site of non-homology. Indeed, RecA protein is capable of forming a complex between ss- and dsDNA, or between ssDNA molecules with little or no sequence complementarity (173,200-203). These complexes, called coaggregates, contain many DNA molecules arranged together in a three-dimensional network that can be sedimented in a low speed centrifuge (204,205). They form rapidly, and kinetically

precede joint molecule formation. Thus, it has been inferred that coaggregates, or a related complex, of RecA protein and ss- and dsDNA are intermediates on the pathway of the strand exchange reaction (204,205).

Although RecA protein binds ssDNA in a sequence-independent manner (though compositional preferences exist (95,176,206,207)), the problem of the homology search is analogous to the problem that sequence-specific DNA binding proteins face in locating their target sequence (208). However, for the case of RecA protein and its analogues, the scale of the search problem is much larger: the binding protein is the entire nucleoprotein filament (which can consist of thousands of protein monomers, depending on the ssDNA length), and the target is the unique complementary sequence within the entire genome. Although the entire nucleoprotein filament is involved in the search for homology, the minimum length of homology required for recognition is as low as 15 nucleotides *in vitro* (95), which is somewhat less than that needed *in vivo*, where homologous recombination requires minimally, about 23-40 bp of homology (209,210).

For homologous alignment to occur, a second DNA molecule must bind to the secondary DNA binding site of the RecA protein filament. This binding, although not sequence specific, does have a hierarchy: optimal binding occurs with oligopyrimidinic DNA while poorer binding is observed with oligoadenylic DNA (173). This hierarchy of binding is not due to the sequence of the ssDNA present in the primary site, but rather is due to an intrinsic property of RecA protein (173). Thus, the proposal that homologous recognition occurred by a novel form of non-Watson-Crick base-pairing called “self-recognition” (211,212) is not substantiated (173,174).

How then does RecA protein “sense” when homology has been located? Though many details of the homology search remain unknown, several steps are clear. During the search, the binding of a second DNA molecule to the nucleoprotein filament must, by necessity, be both weak and transient, to facilitate a rapid search. When the homologous locus is found, the dsDNA is unwound, and one of the strands of the duplex pairs with its complement (which is bound to the primary, original DNA-binding site of the pairing protein). The other strand is displaced into the secondary DNA-binding site, where it is bound more tightly than the duplex DNA. During the search for homology, the DNA duplex remains base-paired; it is only when homology is located, that strand switching occurs. The recognition of homology between the ssDNA within the filament and either the minor groove of dsDNA (213-215), or the major groove of the dsDNA, provides the signal to RecA protein that homology has been located (173). Immediately after base-pair switching, the heteroduplex dsDNA product would occupy the primary site, whereas the displaced ssDNA would occupy the secondary site; there is evidence to support this scenario (174,214,216,217).

Once the two DNA molecules are homologously aligned, RecA protein then catalyzes the nascent exchange of

strands. In the prototypical *in vitro* reaction depicted in figure 3, this requires a local denaturation of the dsDNA molecule and the subsequent exchange of the identical single strands of DNA. These steps may be simultaneous or separated in time, but the result is the production of an intermediate known as a joint molecule. Two types of joint molecules may form, depending on the topological constraints of the DNA: either paranemic or plectonemic. A paranemic joint is one in which the individual complementary strands do not intertwine, producing a molecule that is base-paired but not topologically linked; whereas, a plectonemic joint is one in which the incoming single strand is intertwined around its complement as in native dsDNA. In the reaction displayed in figure 3, paranemic joints will form in the interior of the duplex substrate, and plectonemic joints will form at the ends of the duplex substrate. An experimental distinction between paranemic and plectonemic joints is that paranemic joints require RecA protein (or its counterpart) for stability whereas plectonemic do not. Since paranemic joints are statistically more probable than plectonemic, paranemic molecules are likely intermediates on the reaction pathway to the formation of more stable plectonemic joint molecules (218,219).

Both the type and efficiency of joint molecule formation are affected by SSB protein. If SSB protein is omitted, the displaced ssDNA is used by RecA protein to form a second joint molecule with another dsDNA molecule; this results in the formation of complex, homology-dependent networks of joint molecules. In the presence of SSB protein, RecA protein does not form these networks, since SSB protein prevents re-invasion events by binding to the ssDNA displaced from the joint molecules; in addition, the yield of joint molecules is greater (220). However, under reaction conditions where RecA protein is better able to compete with SSB protein (*i.e.* in the presence of the volume excluding agents polyethylene glycol or polyvinyl alcohol; or when dATP is used as cofactor), networks are readily formed (186,221,222). Thus SSB protein acts both at the pre- and post-synaptic steps of DNA strand exchange.

4.2.2. Bacteriophage T4

Some aspects of synapsis are similar for UvsX protein and RecA protein: 1) UvsX protein coaggregates non-homologous ssDNA and dsDNA (47); and 2) UvsX protein catalyzes formation of approximately equal numbers of paranemic and plectonemic joint molecules (223). Differences also exist between the two proteins. First, while both RecA protein and UvsX protein form homology-dependent DNA networks in the absence of a single-stranded DNA-binding protein, UvsX protein also forms networks even in the presence of gp32, but this may simply reflect an increased ability of UvsX to displace gp32 (50). Second, since UvsX protein, unlike RecA protein, binds readily to either ssDNA or dsDNA under normal strand exchange conditions (46), joint molecule formation is reduced due to UvsX protein binding to the dsDNA (223); this behavior more closely resembles that of Rad51 protein (see below). This limitation is overcome *in vitro* by using limiting amounts of UvsX protein relative to the DNA (45).

4.2.3. *S. cerevisiae*

As for UvsX protein, joint molecule formation is inhibited if Rad51 protein is allowed to bind dsDNA and can

be avoided by assembling presynaptic filaments using exactly stoichiometric amounts of Rad51 protein, or by using excess dsDNA (63). This inhibition of joint molecule formation results from the rapid and kinetically stable binding of Rad51 protein to dsDNA, a situation that is very different from the behavior of RecA protein.

Rad51 protein initiates joint molecule formation if the linear dsDNA contains a single-stranded overhang at least 2 nucleotides in length that is complementary to the ssDNA within the presynaptic filament (199). Both joint molecules and nicked circle products are reported to form more efficiently with dsDNA containing a 3'-overhang than with a 5'-overhang (199).

4.3 Branch migration (figure 3, Step 4)

Once the plectonemic joint has formed, the branch migration phase of DNA strand exchange commences. During this phase, the nascent, heteroduplex joint is extended until complete exchange of single strands of DNA occurs, resulting in a nicked, double-stranded circle. Though kinetically distinct, branch migration may not be a mechanistically separate step, but rather may represent a continuation of plectonemic joint molecule formation (219,224).

4.3.1. *Escherichia coli*

For RecA protein, branch migration proceeds in a 5'→3' direction relative to the incoming single strand (the same direction as RecA protein polymerization), at a rate of 2-10 bp sec⁻¹ (171,225), requires ATP hydrolysis (19,226) and induces torsional stress in the dsDNA (227,228). Branch migration is relatively tolerant of nucleotide sequence mismatches (229,230) and DNA lesions (231,232), although the reaction is inhibited 60% by heterologous insertions of 140 bp within the dsDNA substrate and by insertions of 1000 nucleotides within the ssDNA substrate (230,231).

The ability of RecA protein to not only promote DNA heteroduplex extension, but also to extend the DNA heteroduplex beyond limited regions of sequence non-homology (heterology), may serve a valuable biological function, despite the existence of specialized branch migration helicases like the RuvAB proteins (233). For this reason, the mechanism by which RecA protein facilitates the bypass of DNA sequence heterology is an important issue. Several studies established the key requirements for this heterology-bypass (234,235) (D.A. Kitchell and S.C.K., unpublished observations). RecA protein can facilitate bypass through a heterology as large as 140 bp within the dsDNA substrate (230,231), provided that these mismatched sequences are beyond the point at which DNA strand exchange initiates (*i.e.*, the heterology needs to be beyond the 5'-end of the strand being displaced from the dsDNA). However, RecA protein is unable to bypass an insert of only 22 base pairs that is at the distal end of the dsDNA (234). In all cases, the ability of RecA protein to promote DNA heteroduplex extension progressively diminishes as the length of homology distal to the heterology decreases. Furthermore, when the length of the heterology increases, the amount of distal homology required to bypass this heterology increases, apparently to compensate for the longer heterologous distance that must be traversed.

Thus, to bypass a heterology, DNA strand exchange must initiate in a region of DNA sequence homology, and there must be additional homology beyond the heterology. ATP- γ -S could not substitute for ATP in these bypass experiments, demonstrating that ATP hydrolysis was necessary, presumably to permit some dissociation/redistribution of the RecA nucleoprotein filament.

Several mechanisms were proposed to explain these important characteristics. One envisioned that once the heterology was encountered, lateral slippage of the exchanging strand within the RecA nucleoprotein filament would permit re-alignment of the homologous sequences (234). A second mechanism envisioned that facilitated rotation of the DNA substrates about each other mediated by the RecA protein filament in an ATP-hydrolysis coupled mechanism, provided the "motor" that rotated the DNA past the heterology; the specific details of this model were elegantly elaborated previously to which the reader is referred for details (235). Here we would like to present an alternative idea, based on these collective data and unpublished considerations (D.A. Kitchell and S.C.K., unpublished observations).

Figure 4 shows how topological stress, resulting from local unwinding of dsDNA that is constrained in a closed topological domain, can facilitate bypass through a heterology. The basic premise of the topological model is that, homologous pairing involves intermediates (the synaptic complexes) in which the dsDNA is highly unwound. Furthermore, in the presence of ATP, the RecA nucleoprotein filament that participates in DNA strand exchange is dynamic, and it is frequently binding and releasing dsDNA in its quest for proper DNA sequence alignment. Electron microscopy shows that the RecA nucleoprotein filament can envelop kilobase pair-lengths of homologous dsDNA (216), and that the binding of dsDNA by the RecA nucleoprotein filament produces a topological unwinding of the dsDNA from the canonical 10.4 bp/turn to 18.6 bp/turn (236-241,242 (figure 4)). Thus, in the static situation, when the RecA nucleoprotein filament envelops dsDNA, the dsDNA including the heterology is extensively unwound by up to 8.2 bp/turn resulting in 18.6 bp of dsDNA/filament turn versus 10.4 bp turn in B-form dsDNA (figure 4C). However, in the dynamic case, in the presence of ATP, if the RecA nucleoprotein filament were to release a medial region of the dsDNA while the two flanking regions of the dsDNA remained bound to the filament, then the binding energy that maintained the bound dsDNA in its unwound form would be absent, and the released dsDNA segment would be both topologically constrained and highly underwound (figure 4D, left side); in fact, the release of dsDNA from just 2 turns of the filament would result in 16.4 bp of unwinding. Thus, the transient release of dsDNA from 20 turns of the filament would produce a domain that is unwound by 164 bp relative to B-form DNA. If this unwinding occurred in a region of dsDNA heterology, then there would be sufficient, locally trapped unwinding to induce transient strand separation of the dsDNA (figure 4D, right side), and to bypass the heterology in a kinetically stochastic way. With the region of dsDNA heterology locally melted, bypass would readily occur upon subsequent re-polymerization of the filament (figure 4E);

however, if a nick were located in the region of heterology, then the superhelical strain is released and no bypass occurs. This model explains the key experimental requirements for bypass (the requirement for increasingly longer lengths of homology distal to the heterology, the need for ATP, and the inhibitory effect of a dsDNA nick), and it lends insight into how RecA protein, and presumably all of the other DNA strand exchange proteins (since they all extend dsDNA when bound) can mediate bypass of short regions of DNA sequence heterology without the need for specialized branch migrations proteins.

4.3.2. Bacteriophage T4

UvsX protein, like RecA protein, catalyzes branch migration such that DNA strand displacement is 5'→3' relative to either the incoming presynaptic filament or the displaced strand (45,50). However, UvsX protein catalyzes branch migration at a rate of 12-15 bp sec⁻¹ (50,151), which is faster than the rate catalyzed by RecA protein (see above).

4.3.3. *S. cerevisiae*

In contrast to RecA and UvsX proteins, Rad51 protein promotes strand exchange with the opposite polarity (i.e. 3'→5' relative to the incoming single strand (63)) at a rate of 0.5 bp sec⁻¹, and has no requirement for ATP hydrolysis at any stage (64). In addition, the rate of formation of both joint molecule intermediates and products in the reaction is significantly reduced compared to that of either the RecA (2-10 bp sec⁻¹) or UvsX proteins (12-15 bp sec⁻¹) (62-64). Also, this stage of the reaction is inhibited by a 458 bp heterologous insertion in the dsDNA substrate (63).

4.4. ATP hydrolysis in DNA strand exchange

DNA strand exchange requires the presence of a nucleoside triphosphate cofactor, usually ATP. Under standard conditions, ATP hydrolysis coincides with the pairing and exchange of strands of DNA. Initially, it was thought that ATP hydrolysis was a requirement for DNA strand exchange. It is now known that neither the hydrolysis of ATP (183,226) nor the presence of a high-energy phosphate bond (184) are necessary for DNA strand exchange to occur. It is the formation of the high-affinity ssDNA-binding state of RecA protein that is necessary for DNA strand exchange; the formation of which is brought about by ATP binding to RecA protein bound to DNA. The binding of the non-hydrolyzable analogue of ATP, ATP- γ -S, or the non-covalent complex of ADP-AlF₄⁻ can also induce the high-affinity DNA-binding state of RecA protein. Although the hydrolysis of ATP is not required for the exchange of DNA strands, it is required at phases of DNA strand exchange that require the dissociation of RecA protein that is induced by ADP, the product of ATP hydrolysis.

4.4.1. *Escherichia coli*

RecA protein is a DNA-dependent ATPase with a single active site for the binding and hydrolysis of ATP and other nucleoside triphosphates (17,206,243-245). The enzyme is able to hydrolyze ATP with either ss- or dsDNA as cofactor, resulting in a k_{cat} with ssDNA of 18 - 30 min⁻¹ and with dsDNA of 19 to 22 min⁻¹ (178,246,247). The active species in ATP hydrolysis is the nucleoprotein filament, with ATP being hydrolyzed uniformly throughout the filament and




Figure 4. Model for heterology bypass by RecA protein. A pairing event involving ssDNA (drawn in blue) and dsDNA (drawn in red/pink) containing a region of heterology, is shown. The spheres represent RecA protein; the large black arrow indicates the direction of migration of the heteroduplex joint. Details of this model are discussed in the text.

with no detectable enhancement at filament ends (117,177,248). Hydrolysis of ATP results in one or more conformational changes in RecA protein. ATP and ADP serve to modulate RecA between the "high-affinity" DNA-binding and "low-affinity" DNA-binding states, respectively (179,191). It may not be the hydrolysis *per se* that modulates the two states, but rather the release of P_i , which is extremely rapid (249). This proposal is supported by experiments utilizing the ground state analogue, ADP- AlF_4^- , which maintains RecA protein in the high-affinity ssDNA-binding state in the absence of a high-energy bond (184,250).

Although ATP hydrolysis occurs throughout DNA strand exchange promoted by RecA protein, it is not required for all of the stages (19,184,219,226,251-254). ATP hydrolysis is not required for the exchange of DNA strands (184,250,255). However, ATP hydrolysis is required to dissociate RecA protein from the heteroduplex products once the reaction is complete (226,252,256), to facilitate the bypass of structural barriers such as heterologous sequences (253,254), and to maintain reaction polarity (125,257). In addition, ATP hydrolysis is required for DNA strand exchange involving four strands (258).

Wild-type RecA protein is able to promote strand exchange using the non-hydrolyzable analogue ATP- γ -S (226). In addition, a mutant RecA protein, K72R, in which ATP hydrolysis is reduced by approximately 600-850-fold, is able to promote homologous pairing and exchange of up to 1.5 kilobase pairs of DNA (183). In both of these situations, extension of the heteroduplex is blocked (*i.e.* joint molecule intermediates form but are not converted to gapped or nicked dsDNA products) and the reaction is bi-directional (257), which is in contrast to that of the ATP reaction, which has a distinct polarity (171). However, a nucleoside triphosphate is not an essential component of the pairing and exchange process, because a complex of ADP and AlF_4^- activates all of the activities of RecA protein (184,250,255).

4.4.2. Bacteriophage T4

UvsX protein, in contrast to RecA protein, has an absolute requirement for ATP hydrolysis in DNA strand exchange (259). The rate of ATP hydrolysis by UvsX protein is 8- to 10-fold higher than that of RecA protein, with a k_{cat} of 145 - 240 min^{-1} (44), and this activity is inhibited by both ADP and ATP- γ -S (259). In the presence of 2 mM ATP- γ -S, UvsX protein is unable to promote strand exchange; however at 16 and 80 μ M ATP- γ -S, UvsX protein can form intermediates (259). This may simply reflect the situation that existed for RecA protein prior to the realization that the ATP- γ -S reaction had a unique reaction optimum (184). Substitution of ATP- γ -S for ATP allows UvsX protein to bind ssDNA, unwind dsDNA (259), and promote DNA strand annealing (45). Addition of ATP- γ -S to an ongoing DNA strand exchange reaction, initiated with ATP, causes a transient increase in the rate of branch migration (50). This transient stimulation, which occurs due to inhibition of ATP hydrolysis, is attributed to a kinetic stabilization of the active filamentous form of UvsX protein. The high turnover of the UvsX protein nucleoprotein filament appears to limit the steady-state rate of branch migration (259).

4.4.3. *S. cerevisiae*

Positioned at the opposite end of the ATP hydrolysis spectrum is the eukaryotic Rad51 protein. Although Rad51 protein does not require ATP hydrolysis at any step of DNA strand exchange *in vitro* or *in vivo* (64), it does hydrolyze ATP with a k_{cat} of 0.73 min^{-1} (64). Rad51 protein is able to promote both intermediate formation and directional DNA strand exchange using either ATP- γ -S or AMP-PNP, two non-hydrolyzable analogues of ATP (64). In addition, a Rad51 mutant protein, K191R, (the analogue of the RecA K72R mutant protein), is defective for ATPase activity *in vitro*, yet it can promote polar DNA strand exchange (64). Interestingly, the mutant Rad51 protein is phenotypically normal when expressed in yeast cells, demonstrating that ATP hydrolysis is unnecessary for Rad51 protein function (64).

This raises the question of why such a large difference exists between these three proteins in the rates of ATP hydrolysis during DNA strand exchange? Since ATP hydrolysis is not needed for DNA strand exchange, at least two possible explanations exist. First, there may be a relationship between the requirement for ATP hydrolysis and the genetic complexity of the organism. Such a relationship would exist if the presynaptic filament was required to persist for the longer times needed to find DNA sequence homology in the more complex organism: those with the most complex genome would require the more stable presynaptic complex which would exist if the ATP turnover were lower. In accord with this idea, T4 phage which has the least complex genome, has the highest rate of ATP hydrolysis; *E. coli* which has an intermediate complexity, has a corresponding intermediate rate of ATP hydrolysis; and *S. cerevisiae*, which has a more complex genome, has little or no ATP hydrolysis. Alternatively, the requirement for ATP hydrolysis may be related to the length of the organism's cell cycle: T4 undergoes rapid growth after infection; *E. coli* has a doubling time of 20 to 30 minutes; whereas *S. cerevisiae* has a doubling time ranging from 90 to 140 minutes. Since disassembly of the presynaptic complex is also presumably important to a cell's metabolism, the cells with the fastest growth rate should have a pairing protein with the highest ATP turnover. Both views may be correct: since it should take a longer time to identify homology in a more complex genome, it is reasonable to expect that the presynaptic filament should be kinetically more stable. Furthermore, since ATP hydrolysis is also used to remove the DNA strand exchange protein from the DNA heteroduplex product, the more slowly growing organism may simply have the luxury of waiting for a slower dissociation process.

5. STRUCTURAL COMPARISON

In this section, we compare the form of the nucleoprotein filaments for each protein. We also compare the location of conserved amino acid sequences within the context of the crystal structure of RecA protein, using homology-dependent structural modeling for the Rad51 and UvsX proteins. Although the three proteins do form structurally similar nucleoprotein filaments (figure 5), and they do have significant homology at the amino acid level, they are significantly different in key regions, making each protein unique.




Figure 5. Electron micrographs of nucleoprotein filaments formed on dsDNA. (A) RecA nucleoprotein filament assembled on nicked, circular dsDNA. Filaments were formed by incubating RecA protein with the dsDNA substrate in the presence of ATP for 20 minutes, followed by the addition of ATP γ S for an additional 60 minutes to stabilize the complex. The bar in this figure represents 0.1 nm. The striations visible are due to the helical nature of the filament, with the darker bands corresponding to groove in the filament where dsDNA and repressor binding occurs (237). Reprinted with permission from Nature 299, 185-186, Copyright (1982) Macmillan Magazines Limited. (B) UvsX nucleoprotein filament assembled on linear M13 dsDNA. The complexes were formed by incubating UvsX protein with the DNA substrate in the presence of ATP for 8 minutes, followed by fixation and staining with uranyl acetate (46). The scale is not the same as in (A); though the bar represents 0.1 nm. Reprinted with permission from the Journal of Biological Chemistry 260:4484-4491, Copyright (1985) The American Society for Biochemistry and Molecular Biology. (C) Rad51 nucleoprotein filament assembled on nicked, circular ϕ X dsDNA. Filaments were formed in the presence of ATP for 60 minutes, followed by the addition of ADP-AlF $_4$ for an additional 30 minutes to stabilize filaments. The filament shown was negatively stained with uranyl acetate (263). The tobacco mosaic virus particles (indicated by the arrows), are approximately 200 Å in diameter. Reprinted (abstracted/excerpted) with permission from Science 259:1896-1899. Copyright (1993) American Association for the Advancement of Science.

5.1. Filament structure

5.1.1. *Escherichia coli*

In the presence of either ATP, ATP- γ S, or ADP-AlF $_4$, RecA protein polymerizes onto either ss- or dsDNA in a highly cooperative, polar fashion (187,236,260,261). On dsDNA, the resulting helical nucleoprotein filament (figure 5A) contains 6.2 RecA protein monomers per turn, has a pitch of 95 Å and is approximately 110 Å in diameter (188). The stoichiometry, as determined either by mass per unit length measurements (188), or by fluorescence measurements and sedimentation analysis (262), is one RecA monomer per three base-pairs.

The RecA protein filament is arranged so that a deep helical groove exists (the striation in figure 5A). Although it is not visible in the figure shown, one side of the groove is smooth, while the other is penetrated by the protrusion of the individual monomers (263). These protrusions, or lobes, arise from residues 270 to 328, which corresponds to the highly acidic, carboxyl-terminus of the protein. A similar groove exists for Rad51 protein although no corresponding lobes appear to exist, and this is most likely due to the fact that the Rad51 protein sequence, when aligned against that of RecA protein, terminates at amino acids 266-267. This would suggest that RecA protein should possess additional functions, which may be attributed to this lobe region. Truncated RecA proteins, lacking from 20 to 45 C-terminal residues, are capable of promoting LexA repressor cleavage and DNA strand exchange, which suggests that this region is not involved in these activities (264-266). This region was suggested to be involved in filament-filament interactions that may be involved in keeping RecA protein in an inactive form (267). A more recent study presents evidence that the conserved basic amino acid residues within this region have a direct role in binding dsDNA, and that they form part of a “gateway” for homologous recognition (268).

Within the nucleoprotein filament, the conformation of both ss- and dsDNA is significantly different from that of native B-form DNA. Both DNA types are extensively stretched (approximately 150% of B-form) and, in the case of dsDNA, the helix is also unwound (237). For dsDNA, there are approximately 18.6 base-pairs per turn and the axial rise per base-pair increases from 3.4 Å to 5.1 Å (236,269) while the twist changes from 35° to 20° per base-pair (237,238,262). Electron micrographic analysis and neutron scattering reveal that the DNA is located near the axis of the nucleoprotein filament (270,271). This means that in order for DNA strand exchange to occur, the second DNA molecule must somehow find its way into this complex.

5.1.2. Bacteriophage T4

Similar to RecA protein, UvsX protein also forms helical filaments on both ss- and dsDNA in the presence of a nucleotide cofactor (e.g. ATP, ATP- γ S, or Br $^{\delta}$ -ATP) (46,48). Also, resembling RecA protein, UvsX protein forms a helical filament on ssDNA with a distinctly different structure in the absence of ATP (46,48). Figure 5B shows an electron micrograph of UvsX protein on linear dsDNA in the presence of ATP. The helical pitch of the extended UvsX protein-DNA filaments depends on the nucleotide cofactor present. When ATP is present, the pitch is about 120 Å (46).

However, when ATP- γ -S is employed, the mean pitch is 88.6 Å (48). This is regarded as similar to the helical pitch of 95 Å obtained for RecA protein-DNA complexes in the presence of ATP- γ -S (272). Within the UvsX protein-DNA filament, the DNA is stretched such that the axial rise is 5 Å per base or base-pair and untwisted to about 19 bases or base-pairs per helical turn (48). These DNA parameters are very similar to those of the DNA within the RecA protein filament (272). Although there are many similarities between the two filaments, there is one significant difference: when the filaments are formed in the absence of a nucleoside triphosphate cofactor, UvsX protein forms an extended filament (48), whereas RecA protein does not (272).

5.1.3. *S. cerevisiae*

The Rad51 protein from *S. cerevisiae* forms helical filaments on DNA similar to those of RecA and UvsX proteins, and it induces a similar conformation in the DNA (263). On dsDNA, Rad51 protein forms helical nucleoprotein filaments with a helical pitch of 99 Å and 6.2 subunits per turn (figure 5C). The binding stoichiometry is three base-pairs per Rad51 protein monomer, which is consistent with the value for RecA protein (188). In addition, the rise per base-pair induced by the binding of Rad51 protein also increases from 3.4 Å to 5.1 Å; thus, the DNA is also stretched by a factor of 1.5 while having its helicity reduced from 10.5 to 18.6 base-pairs per turn. Sung and Robberson obtained similar results, although they obtained a helical pitch of 80 Å, 17.6 bp per helical repeat and a predicted DNA length that was extended to only 134% that of B-form DNA (63). Although Rad51 protein is able to form filaments on duplex DNA, this nucleoprotein filament is not active for DNA strand exchange (63).

In the presence of RPA, Rad51-ssDNA nucleoprotein filaments resembled those of RecA protein assembled on ssDNA, having a right-handed helical appearance with a pitch of 94 Å (63). Within the complex, the ssDNA is stretched to approximately 154% of native B-form DNA. Filaments formed between Rad51 protein and DNA are less regular than those of RecA (263) and individual lobes of Rad51 monomers are also less visible than those of RecA protein. This is due to structural differences between the Rad51 and RecA protein filaments and not due to disorder within the Rad51 protein filament (263).

5.2. Primary amino acid sequence and crystal structure

There is considerable similarity at the amino acid level between these proteins (29,30,59,79,273-275). We divided the primary amino acid sequence of the three proteins into three sections: an amino-terminal domain, a central "core" region, and a carboxyl-terminal domain; the proteins were then aligned at the highly conserved A and B Walker motifs (figure 6) (274). This alignment is colored; the colors correspond to the 8 highly conserved motifs identified by the Karlin laboratory (29,274). The same colors are used in both the RecA protein structure and the theoretical structures presented for the UvsX and Rad51 proteins (figure 7). The models for Rad51 and UvsX proteins are not intended to imply genuine 3-dimensional structures; instead they show the potential positions of the conserved domains within the crystal structure scaffolding of the *E. coli* RecA protein. In

figure 8A, we highlight the residues that are identical in UvsX and RecA proteins, and in figure 8B, we highlight the residues that are identical in Rad51 and RecA proteins (29,30).

5.2.1. *Escherichia coli*

RecA protein is a 352 amino acid (aa) protein (276). The sequence is highly conserved in prokaryotes (43 to 98% at the amino acid level) (28,30), but less well conserved across genera (29,274). The amino-terminal domain consists of aa 1 through 33; the central "core" region extends from aa 34 to 240, and the carboxyl-terminal domain contains residues 241 through 352 (Figures 6 and 10).

The crystal structure for RecA protein was solved to 2.3 Å resolution in the absence of DNA and in the presence of ADP that was diffused into the crystal (figure 7A, 8 - 10; and references (267,277)). This structure contains a major "core" domain with regions that bind ADP and are proposed to bind both ss- and dsDNA. In addition, it contains two smaller domains at the amino- and carboxyl-termini that protrude from the structure and stabilize the helical polymer (267); the amino-terminal domain is important in forming the monomer-monomer interface in the nucleoprotein filament and also in regulating the self-assembly process (278), whereas the carboxy-terminal domain is involved in controlling dsDNA binding (265,268).

The "core" region of RecA protein extending from aa 34-240 (figure 6, 7A), consists of 8 sub-domains, designated B1, B2, C1, C2, D1, E1, E2 and F1 (274). These sub-domains are highly conserved among eubacterial RecA proteins and also among eukaryotic RecA protein analogues (29). This region contains residues which constitute the ATP binding domain (224), the two highly conserved Walker motifs A and B, which correspond to residues 66-73 and 140-144, respectively (279), and the proposed DNA binding regions, designated Loops 1 (aa 157-164) and 2 (aa 195-209) (figure 9) (267). In the crystal structure, this domain contains eight β -sheets flanked by α -helices. By diffusing ADP into crystals, it was possible to identify residues involved in nucleotide binding (267,277). Amino acids D100, Y103 and G265 all make contacts with the base of the nucleoside. The two highly conserved Walker motifs, designated A and B, also make contact with the nucleoside diphosphate (figure 9). Motif A interacts with the PP_i moiety of ATP (277) and is essentially invariant among eubacterial RecA proteins (29), but is less well conserved among the three proteins compared here (figure 6). Residue K72 (RecA protein numbering) is an invariant residue, which when mutated to arginine, abolishes ATP hydrolysis (but does not affect ATP binding), indicating that Motif A is part of the catalytic domain for hydrolysis (183). A similar result was observed for Rad51 protein where mutation of the equivalent residue to arginine, producing the Rad51 K191R protein, abolished ATPase activity completely (64). Motif B, which corresponds to the N-terminal portion of sub-domain E1, is also highly conserved across all RecA protein sequences. This Motif consists of four successive hydrophobic residues followed by an aspartic acid at position 144, which is the key residue in Motif B (267,277) and is identical in all RecA protein sequences (29,30). This residue coordinates the Mg²⁺ ion at the active site between the β and γ phosphates (277). The

DNA strand exchange proteins




Figure 6. Sequence alignment of RecA, Rad51, and UvsX proteins. The three proteins are aligned at the conserved ATP binding motifs A and B (dotted lines). Numbers immediately to the right of the Rad51 and UvsX proteins are the sequence similarity and identity, respectively, when the entire protein sequence is compared to RecA protein. Numbers in green above the core domain of Rad51 and UvsX proteins are the sequence similarity and identity as compared to RecA protein, for the core regions only. The assignment of the 8 sub-domains (B1, B2, C1, C2, D1, E1, E2, and F1) is the result of an alignment by Karlin and coworkers (274). The sequence for each sub-domain is shown. The order of the sequences presented is RecA protein, Rad51 protein, and UvsX protein. The numbers flanking each sequence block indicate the residue numbers for the RecA, Rad51, and UvsX protein sequences, respectively. The numbers immediately to the right of each block of sequences indicate the percent similarity as compared to *E. coli* RecA protein (obtained using the program Bestfit, which is part of the GCG software package). The numbers in black immediately above sub-domains B1 and F1 are the first and last residues respectively, of the core domains of each protein.

region immediately C-terminal of Motif B (aa 145 to 149) is invariant among eubacterial sequences and less conserved among other RecA analogue sequences; the serine at 145 and leucine at 149 are invariant, however (figure 6 and reference (29)). With regard to ATP hydrolysis, other important residues are E96 (invariant in all RecA protein analogues) which may be involved in the activation of a water molecule for the attack on ATP leading to hydrolysis, and Y264, which cross-links to 8-azido-ATP (280).

For RecA protein and its analogues to function in DNA strand exchange, they need to accommodate at least three DNA strands: the first strand bound to the primary site, and the second and third bound as dsDNA to the secondary (and perhaps tertiary) binding sites. Although the precise location of these sites remains unclear, two are proposed to be in this central core domain, and the third is proposed to be within the C-terminal domain (268). The two regions within

this core domain that are proposed to form the DNA binding sites, are the disordered regions in the crystal structure which are designated loops L1 and L2 (figure 9) and which are comprised of residues 157-164 and 195-209, respectively (267). These loops lie close to the polymer axis, the location of the DNA substrate (267). L1 forms the inner, upper lip of the polymer and was proposed to be involved in duplex DNA binding, the secondary site for RecA protein (267). L2 projects outward from the carboxyl end of β -strand 5 and lies toward the innermost face of the filament, directly above the ATP binding site. L1 is strongly conserved in eubacterial sequences, but only at the first five amino acids of the region (29). L2 is substantially conserved, but only among eubacterial RecA proteins (29,274). There is little sequence homology with either Rad51 or UvsX proteins in this region (29,273,274). Several lines of evidence, however, suggest that Loops 1 and 2 are involved in DNA binding, though the data for Loop 1 are conflicting. The isolated L1 loop peptide




Figure 7. Structures for the core regions of RecA, Rad51, and UvsX proteins. (A) The core domain for RecA protein. The position of ADP as a reference point is shown in white. (B) The expected structure for the core region of UvsX protein. Note that the sub-domain E2, the most poorly conserved region of UvsX, is incapable of forming an α -helix and also disrupts the sub-domain D1. (C) The expected structure for the core region of Rad51 protein. The two model structures were obtained by modeling the RecA protein structure with each of the 8 sub-domains in the RecA protein sequence replaced by either UvsX or Rad51 protein sequences (296-298). Details of the modeling procedure are discussed in the text. The colors used for each of the sub-domains match those used in figure 6. For each predicted structure, the regions shown in light green/aquamarine are residues from the α -carbon backbone of the RecA protein sequence that was used as a scaffold for the homology modeling. The letters for each sub-domain, described previously (274), are indicated in the same color used for figure 6.

cannot bind DNA (281); it exhibits a large mutational flexibility (282); and chemical modification data show that H163 is protected more by dsDNA than by ssDNA (283), suggesting involvement in dsDNA binding. In contrast, the isolated Loop 2 peptide can bind ssDNA (281) and, surprisingly, a mutant variant of the Loop 2 peptide can promote DNA strand invasion of supercoiled DNA by a single-stranded 53-mer (284), suggesting that Loop 2 may be responsible for binding ssDNA providing charge neutralization, and facilitating the close approach to dsDNA. Further, the effects of mutations in Loop 2 (residues 195-209) and the adjacent helix (helix G, residues 210-221) are consistent with this region being involved in DNA binding (285).

Using photochemical cross-linking, Wang and Adzuma demonstrated that when RecA protein was in excess over ssDNA, cross-linking occurred to M164, suggesting that this region is part of the primary site involved in ssDNA binding (286). Also under conditions of RecA protein excess, Mallkov and Camerini-Otero were able to cross-link to both Loops 1 and 2 using d(IU)-containing substrates, suggesting that both loops comprise the primary site responsible for ssDNA binding (287). Finally, Morimatsu and Horii showed that with a 55-mer oligonucleotide, cross-linking occurred at Y65, Y103, Loop 2 and Y264 (288). In cross-linking experiments using conditions of DNA in excess and d(IU)-containing substrates as the second DNA to probe for the second DNA binding site, the majority of cross-linking occurred to M202 and F203, located in Loop2 (286). In separate studies, cross-linking experiments also using conditions of DNA in excess, which should identify both DNA-binding sites 1 and 2, revealed additional positions that may be involved in, or are close to, the DNA binding sites: aa 61-72 (Motif A involved in ATP binding) (289), aa 178-183 (289,290), and aa 233-243 (partially outside the core region) (289). Finally, in a study using stoichiometric amounts of RecA protein and ssDNA, cross-linking to Y103 was observed (290); this amino acid is well outside Loops 1 and 2. The positions of the residues discussed above are shown in figure 9 (maroon side-chains). The discrepancy between the observed sites of DNA cross-linking and the proposed DNA binding sites in the crystal structure may be due to the fact that the structure determined is of the inactive, ADP-bound, DNA-free RecA protein. Thus in the DNA-bound form of the protein, amino acids removed from the DNA binding site may be in close proximity. Additionally, cross-linking may not have occurred with residues in real contact with the DNA, but rather with residues proximal to the binding site. This characteristic was observed for residue Y264 which cross-links to 8-azido-ATP (figure 9 and (280)) but which is not directly involved in ATP contact in the crystal structure, although it is located within the ATP binding cleft (277).

The carboxyl terminus of RecA protein, which spans aa 241 through 352, is highly acidic, containing ~14% of the acidic residues of the protein and is considerably variable, even among prokaryotic RecA protein analogues (29). This region was proposed to be involved in filament-filament interactions (267), in the interaction with repressor




Figure 8. Conserved residues. (A) The crystal structure of RecA protein is shown in grey with the side chains of the residues that are identical to those in the UvsX protein sequence shown in orange. (B) The crystal structure of RecA protein shown in grey with the side chains of the residues that are identical to those in the Rad51 protein sequence shown in orange. ADP is shown in transparent purple. Labels for the α -helices (white letters) and β -strands (yellow numbers) are from Story *et al.* (267).

proteins (29,291,292), in the control of dsDNA binding (265), and in the formation of the secondary DNA binding site (267,292,293). Partial deletion of this region produces truncated RecA proteins 300 amino acids in length that are able to cleave LexA protein and function in DNA strand exchange (264-266). Mutagenesis of a sub-section of this C-terminal domain (aa 270-328) shows it is involved in binding of dsDNA to nucleoprotein filaments formed on ssDNA (268). The replacement of 9 basic amino acid residues in this region showed that mutations in the two conserved residues, K286 and K302, caused deficiencies in recovery following UV-irradiation, whereas mutations in the non-conserved basic residues had little or no effect (268). Purified K286 and K302 mutant RecA proteins were able to form presynaptic complexes but these complexes, once formed, were defective in binding to and pairing with dsDNA (secondary DNA

binding). These results are consistent with the involvement of this region in dsDNA binding, and its proposed role as a “gateway” for homologous recognition where the dsDNA interacts with the ssDNA inside of the nucleoprotein filament (268). In addition to dsDNA binding, the binding of a secondary DNA molecule (which is critical to DNA strand exchange) and LexA repressor cleavage are competitive processes; these findings are consistent with the dual role for this region in dsDNA and repressor binding (291,294).

5.2.2. Bacteriophage T4

The UvsX protein sequence is 390 amino acids in length. When the amino acid sequences of *E. coli* RecA protein and T4 UvsX protein are compared, the two proteins are 52% similar and 24% identical (Figures 6, 8A and reference (295)). The sub-division of the primary amino acid sequence of UvsX protein is as follows: the N-terminus from aa 1-27; the core region from aa 28-225, and the C-terminus from aa 226 to 390 (figure 6). The most significant difference between the sequences takes place in the carboxyl-terminal region (273), while the most similarity exists within the core region where there is 51% similarity and 28% identity (figure 6). The core region can also be divided into 8 sub-domains (figure 6 and 7B). However, there is poor alignment at sub-domains E2 and F1; alignment only becomes possible when large gaps are introduced into the RecA protein sequence, as shown in the alignment of Story *et al.* (273). Even though there is low sequence similarity between UvsX and RecA proteins in these two sub-domains, we included them in the modeling process using the sequence alignment shown in figure 6. Modeling of the 8 sub-domains, using the RecA protein crystal structure as a scaffold, was conducted as follows: the positions of the 8 sub-domains were located in the primary amino acid sequence of RecA protein; these residues were then replaced with the corresponding sequence blocks from the UvsX protein primary sequence. This hybrid sequence was submitted to Swiss-Model, which is an Automated Protein Modeling Server running at the Geneva Biomedical Research Institute, Glaxo Wellcome Research and Development S.A., Switzerland (available at <http://expasy.hcuge.ch/swissmod/SWISS-MODEL.html>) (296-298). The structure for this hybrid protein, showing only the core region, with conserved blocks in color and with RecA protein residues in aquamarine/green is presented in figure 7B.

Modeling of UvsX protein on the RecA protein scaffold shows that the sub-domains used in the modeling assemble into a structure which closely resembles that of the *E. coli* RecA protein. These include the Walker binding motifs A (sub-domain B2) and B (sub-domain E1), located at aa 60-68 and aa 139-143, respectively. As for the RecA and Rad51 proteins, the invariant lysine is present in Motif A at position 66 and the four hydrophobic residues followed by an aspartic acid at position 143 are also present in Motif B (273). The proposed DNA binding domains in RecA protein, Loops 1 and 2, are not well conserved in the UvsX primary sequence, with 50% similarity in Loop 1 and 37% similarity in Loop 2. Interestingly, in Loop 2, residues V207 and M208, which are identical to those of RecA protein at positions 201 and 202, respectively, are conserved. This suggests that the region containing this amino acid may be in close proximity to one of the DNA binding sites.




Figure 9. Crystal Structure of RecA protein monomer. The crystal structure of RecA protein showing the side chains of relevant residues as discussed in the text. The potential location of Loops 1 and 2 (which are described in the crystal) are shown in yellow; the amino acid side chains of the Walker motifs A and B are shown in orange, with the side-chain of residue K72 highlighted in green; amino acid side-chains that are in close proximity to DNA, as identified by photochemical cross-linking, are indicated in maroon.

5.2.3. *S. cerevisiae*

The various Rad51 proteins range in length from 339 (human, chicken and mouse) to 400 (*S. cerevisiae*) amino acids in length (31,59,66,299-302). All of the eukaryotic analogues have extended N-terminal domains and lack the extended C-terminal domain present in their prokaryotic counterparts. Specifically, the *S. cerevisiae* Rad51 protein is 120 residues longer at the amino terminus and 90 amino acids shorter in the carboxyl terminus than RecA protein (figure 6 and references (31,59,61)). Our division of the primary sequence of *S. cerevisiae* Rad51 protein into three regions is as follows: aa 1-154 constitute the amino-terminal portion; the core extends from residues 155 to 375 and the carboxyl domain comprises aa 376 to 400 (figure 6). Alignment of the complete sequences of the two proteins shows that they are 54% similar and 29% identical (59). Within the core region, RecA and Rad51 proteins are even more alike, with 61% similarity and 35% identity (figure 6 and reference (59)).

The amino-terminal domain (which is unique to each eukaryotic species) was proposed to be involved in monomer-monomer interactions, and in forming species specific interactions with Rad52 protein (303,304). Rad51 and Rad52 proteins from either *S. cerevisiae* or humans interact both *in vitro* (59,305) and *in vivo* (135). In *S. cerevisiae*, the amino acids responsible for this interaction map to a region that includes aa 4 through 187 in Rad51 protein (303); in the case of the human proteins, the domain was not mapped. The *S. cerevisiae* Rad51 protein interacts *via* the carboxyl-terminal one third of the Rad52 protein

(135), whereas the human Rad51 protein interacts *via* aa 270-330 of Rad52 protein (305).

As for RecA and UvsX proteins, the Rad51 protein core region also contains 8 sub-domains; there is significantly more homology within the 8 sub-domains than across the entire sequence (274) (figure 6). We have also modeled the Rad51 core domain on the crystal structure of RecA protein. This modeling was done in the same way as for UvsX protein. The model is presented in figure 7C and shows that the sub-domains used in the modeling assemble into a structure, which closely resembles that of both the *E. coli* RecA protein and the T4 UvsX protein.

The core region contains the two Walker motifs: Motif A is located at aa 185 to 192 and Motif B is located at aa 276 to 280. This region may also be involved in interaction(s) with other members of the RAD52 epistasis group that were demonstrated using the two-hybrid system (136). It was noted that the core region may contain a leucine zipper motif (L-X₆-L-X₆-L-X₆-F) at aa 296-317, present in sub-domain F1 (63). The RecA and Rad51 proteins share the most homology within sub-domain E1, which contains Motif B (274) (figure 6 and 8B). Mutation of the conserved lysine at position 191 to alanine results in deficiencies in DNA repair and recombination (59). In contrast, conservative mutation of the same residue to arginine, produces the Rad51 K191R protein which is the analogue of the K72R RecA mutant protein, and which retains biological activity *in vivo*, is defective in ATPase activity *in vitro*, but is still able to promote DNA strand exchange *in vitro* (64). The core region also contains the proposed DNA binding sites of RecA protein, Loops 1 and 2. The modeling of this region using *E. coli* RecA protein residues as spacers between the conserved sub-domains, demonstrates that this region is able to assemble into a similar structure to that of RecA protein. Those regions where sequence similarity is low (sub-domains B2, E2 and F1), have dissimilar predicted structures (figure 7C), while those where sequence similarity is high (sub-domains B1, C, D and E1), have a more similar 3D structure which is almost identical to that within RecA protein (figure 7C).

The C-terminus extends from aa 375 to 400. Within this region, the RecA and Rad51 proteins share very little sequence similarity. The absence of the extended C-terminal domain that is found in the RecA and UvsX proteins may account for the relatively high dsDNA binding activity of Rad51 protein that distinguishes the eukaryotic family from its prokaryotic counterparts. Though inhibitory to DNA strand exchange *in vitro* with naked substrates, this enhanced dsDNA binding may facilitate homologous pairing with dsDNA bound by nucleosomes *in vivo*.

6. PERSPECTIVE

The three proteins reviewed here exhibit many similarities and differences. They show significant homology at the amino acid level; they form similar nucleoprotein filaments on both ss- and dsDNA, and they assume similar structures in their highly conserved central core regions. All three proteins promote DNA strand exchange *in vitro*, which is consistent with the conservation of a central core domain containing all of the sub-domains required for this activity.




Figure 10. Location of the highly conserved core domain within the RecA protein hexamer. Two views of a single turn of a RecA filament are presented. The orientation in (A) is similar to that of Story *et al.* (267) and shows a single turn of the RecA protein filament from the side. (B): The same turn of the RecA filament viewed from the top. For this view, the hexamer in (A) was rotated 90 degrees toward the viewer. The numbers in white indicate the six monomers making up the turn. The core domain of each of the monomers is represented as a solvent surface; these are coloured alternately in orange and green. The α -carbon backbones of the non-conserved regions, the N- and C-termini, are displayed as stick diagrams in yellow. The position of ADP within each monomer is shown as a reference point (red solvent surface).

The differences between these proteins exist in their NTP requirements in DNA strand exchange, and their rates of ATP hydrolysis and their polarity of DNA strand exchange. From a comparison of the amino acid sequences and homology modeling, it is apparent that the N- and C-terminal domains of these proteins are significantly dissimilar. This likely reflects the differences in the additional functions that these proteins perform in their respective hosts, and in the unique protein-protein interactions required for these functions. Thus, though defining the ubiquitous class of DNA strand exchange proteins, these three representative proteins nevertheless display an intriguing diversification of biochemical functions that will require further scrutiny to unravel fully.

7. ACKNOWLEDGMENTS

We thank Drs. E. Egelman, J. Griffith, and A. Stasiak for allowing us to use their electron micrograph figures and Drs. K. Knight, S. Morrical and P. Sung for their careful reading of the manuscript. We also thank the following members of our laboratory for their helpful discussions and critical comments on the manuscript: Dan Anderson, Joel Brockman, Frederic Chedin, Jason Churchill, Deana Haddox, Frank Harmon, Noriko Kantake, Julie Kleiman, Alex Mazin, Jim New, Erica Seitz, Tomohiko Sugiyama and Eugene Zaitsev. This work was supported by NIH grants AI-18987 and GM-41347 to SCK.

8. REFERENCES

1. Clark, A.J. and Low, K.B., Pathways and systems of homologous recombination in *Escherichia coli*, in *The Recombination of Genetic Material*, Low, K.B., Ed., San Diego, Academic Press, 155-215, (1988)
2. Mahajan, S.K., Pathways of homologous recombination in *Escherichia coli*, in *Genetic Recombination*, Kucherlapati, R. and Smith, G.R., Eds., Washington, D.C., American Society for Microbiology, 87-140, (1988)
3. Smith, G.R., Homologous recombination in *E. coli*: Multiple pathways for multiple reasons. *Cell* 58, 807-809, (1989)
4. Kowalczykowski, S.C., Dixon, D.A., Eggleston, A.K., Lauder, S.D. and Rehrauer, W.M., Biochemistry of homologous recombination in *Escherichia coli*. *Microbiol. Rev.* 58, 401-465, (1994)
5. Clark, A.J., *rec* genes and homologous recombination proteins in *Escherichia coli*. *Biochimie* 73, 523-532, (1991)
6. Lloyd, R.G. and Sharples, G.J., Genetic analysis of recombination in prokaryotes. *Curr. Opin. Genet. Dev.* 2, 683-690, (1992)
7. Smith, G.R., Homologous recombination in procaryotes. *Microbiol. Rev.* 52, 1-28, (1988)
8. Clark, A.J. and Margulies, A.D., Isolation and characterization of recombination-deficient mutants of *Escherichia coli* K12. *Proc. Natl. Acad. Sci. USA* 53, 451-459, (1965)

DNA strand exchange proteins

9. Clark, A.J., Recombination deficient mutants of *E. coli* and other bacteria. *Annu. Rev. Genet.* 7, 67-86, (1973)
10. Walker, G.C., Inducible DNA repair systems. *Annu. Rev. Biochem.* 54, 425-457, (1985)
11. Clark, A.J., *RecA* mutants of *E. coli* K12: a personal turning point. *Bioessays* 18, 767-772, (1996)
12. Roberts, J.W., Roberts, C.W., Craig, N.L. and Phizicky, E.M., Activity of the *Escherichia coli RecA*-gene product. *Cold Spring Harbor Symp. Quant. Biol.* 43, 917-920, (1978)
13. Little, J.W., Edmiston, S.H., Pacelli, L.Z. and Mount, D.W., Cleavage of the *Escherichia coli* *lexA* protein by the *RecA* protease. *Proc. Natl. Acad. Sci. USA* 77, 3225-3229, (1980)
14. Bryant, F.R. and Lehman, I.R., On the mechanism of renaturation of complementary DNA strands by the *RecA* protein of *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 82, 297-301, (1985)
15. McEntee, K., Kinetics of DNA renaturation catalyzed by the *RecA* protein of *Escherichia coli*. *Biochemistry* 24, 4345-4351, (1985)
16. Menge, K.L. and Bryant, F.R., Effect of nucleotide cofactor structure on *RecA* protein-promoted DNA pairing. 2. DNA renaturation reaction. *Biochemistry* 31, 5158-5165, (1992)
17. Weinstock, G.M., McEntee, K. and Lehman, I.R., ATP-dependent renaturation of DNA catalyzed by the *RecA* protein of *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 76, 126-130, (1979)
18. Cox, M.M. and Lehman, I.R., Directionality and polarity in *RecA* protein-promoted branch migration. *Proc. Natl. Acad. Sci. USA* 78, 6018-6022, (1981)
19. Cox, M.M. and Lehman, I.R., *RecA* protein of *Escherichia coli* promotes branch migration, a kinetically distinct phase of DNA strand exchange. *Proc. Natl. Acad. Sci. USA* 78, 3433-3437, (1981)
20. Cox, M.M. and Lehman, I.R., *RecA* protein-promoted DNA strand exchange: Stable complexes of *RecA* protein and single-stranded DNA formed in the presence of ATP and single-stranded DNA binding protein. *J. Biol. Chem.* 257, 8523-8532, (1982)
21. DasGupta, C. and Radding, C.M., Polar branch migration promoted by *RecA* protein: Effect of mismatched base pairs. *Proc. Natl. Acad. Sci. USA* 79, 762-766, (1982)
22. DasGupta, C., Shibata, T., Cunningham, R.P. and Radding, C.M., The topology of homologous pairing promoted by *RecA* protein. *Cell* 22, 437-446, (1980)
23. McEntee, K., Weinstock, G.M. and Lehman, I.R., Initiation of general recombination catalyzed *in vitro* by the *RecA* protein of *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 76, 2615-2619, (1979)
24. Shibata, T., DasGupta, C., Cunningham, R.P. and Radding, C.M., Purified *Escherichia coli RecA* protein catalyzes homologous pairing of superhelical DNA and single-stranded fragments. *Proc. Natl. Acad. Sci. USA* 76, 1638-1642, (1979)
25. Soltis, D.A. and Lehman, I.R., *RecA* protein promoted DNA strand exchange: Isolation of *RecA* protein-DNA complexes formed in the presence of single-stranded DNA binding protein. *J. Biol. Chem.* 258, 6073-6077, (1983)
26. West, S.C., Cassuto, E. and Howard-Flanders, P., Heteroduplex formation by *RecA* protein: Polarity of strand exchanges. *Proc. Natl. Acad. Sci. USA* 78, 6149-6153, (1981)
27. Mosig, G., The essential role of recombination in phage T4 growth. *Annu. Rev. Genet.* 21, 347-371, (1987)
28. Roca, A.I. and Cox, M.M., The *RecA* protein: structure and function. *Crit. Rev. Biochem. Mol. Biol.* 25, 415-456, (1990)
29. Karlin, S. and Brocchieri, L., Evolutionary conservation of *recA* genes in relation to protein structure and function. *J. Bacteriol.* 178, 1881-1894, (1996)
30. Karlin, S., Weinstock, G.M. and Brendel, V., Bacterial classifications derived from *RecA* protein sequence comparisons. *J. Bacteriol.* 177, 6881-6893, (1995)
31. Ogawa, T., Shinohara, A., Nabetani, A., Ikeya, T., Yu, X., Egelman, E.H. and Ogawa, H., *RecA*-like recombination proteins in eukaryotes: Function and structures of *RAD51* genes. *Cold Spring Harb Symp Quant Biol* 58, 567-576, (1993)
32. Sandler, S.J., Satin, L.H., Samra, H.S. and Clark, A.J., *RecA*-like genes from three archaean species with putative protein products similar to *Rad51* and *Dmc1* proteins of the yeast *Saccharomyces cerevisiae*. *Nucleic Acids Res.* 24, 2125-2132, (1996)
33. Seitz, E.M., Brockman, J.P. Sandler, S.J., Clark, A.J., and Kowalczykowski, S.C., *Genes and Development*, in press, (1998)
34. Harm, W., Mutants of Phage T4 with Increased Sensitivity to Ultraviolet. *Virology* 19, 66-71, (1963)
35. Hamlett, N.V. and Berger, H., Mutations altering genetic recombination and repair of DNA in bacteriophage T4. *Virology* 63, 539-567, (1975)
36. Boyle, J.M. and Symonds, N., Radiation-sensitive mutants of T4D. I. T4y: a new radiation-sensitive mutant; effect of the mutation on radiation survival, growth and recombination. *Mutat Res* 8, 431-439, (1969)

37. Cunningham, R.P. and Berger, H., Mutations affecting genetic recombination in bacteriophage T4D. I. Pathway analysis. *Virology* 80, 67-82, (1977)
38. Conkling, M.A. and Drake, J.W., Thermal rescue of UV-irradiated bacteriophage T4 and biphasic mode of action of the WXY system. *Genetics* 107, 525-536, (1984)
39. Bernstein, C. and Wallace, S.S., DNA repair, in *The Bacteriophage T4*, Mathews, C.K., Kutter, E.M., Mosig, G. and Berget, P.B., Eds., Washington, DC, American Society for Microbiology, 1983, 138-151.
40. Dewey, M.J. and Frankel, F.R., Two suppressor loci for *gene 49* mutations of bacteriophage T4. I. Genetic properties and DNA synthesis. *Virology* 68, 387-401, (1975)
41. Minagawa, T. and Ryo, Y., Substrate specificity of *gene 49*-controlled deoxyribonuclease of bacteriophage T4: special reference to DNA packaging. *Virology* 91, 222-233, (1978)
42. Nishimoto, H., Takayama, M. and Minagawa, T., Purification and some properties of deoxyribonuclease whose synthesis is controlled by *gene 49* of bacteriophage T4. *Eur J Biochem* 100, 433-440, (1979)
43. Mosig, G., Homologous Recombination, in *Molecular biology of bacteriophage T4*, Karam, J.D., Ed., Washington, D.C., American Society of Microbiology, 1994, 54-82.
44. Formosa, T. and Alberts, B.M., Purification and characterization of the T4 bacteriophage *uvsX* protein. *J. Biol. Chem.* 261, 6107-6118, (1986)
45. Yonesaki, T. and Minagawa, T., T4 phage *gene uvsX* product catalyzes homologous DNA pairing. *EMBO J.* 4, 3321-3327, (1985)
46. Griffith, J. and Formosa, T., The *uvsX* protein of bacteriophage T4 arranges single-stranded and double-stranded DNA into similar helical nucleoprotein filaments. *J. Biol. Chem.* 260, 4484-4491, (1985)
47. Harris, L.D. and Griffith, J.D., Formation of D loops by the *UvsX* protein of T4 bacteriophage: a comparison of the reaction catalyzed in the presence or absence of *gene 32* protein. *Biochemistry* 27, 6954-6959, (1988)
48. Yu, X. and Egelman, E.H., DNA conformation induced by the bacteriophage T4 *UvsX* protein appears identical to the conformation induced by the *Escherichia coli* RecA protein. *J. Mol. Biol.* 232, 1-4, (1993)
49. Hinton, D.M. and Nossal, N.G., Cloning of the bacteriophage T4 *uvsX* gene and purification and characterization of the T4 *uvsX* recombination protein. *J. Biol. Chem.* 261, 5663-5673, (1986)
50. Kodadek, T., Wong, M.L. and Alberts, B.M., The mechanism of homologous DNA strand exchange catalyzed by the bacteriophage T4 *uvsX* and *gene 32* proteins. *J. Biol. Chem.* 263, 9427-9436, (1988)
51. Yonesaki, T. and Minagawa, T., Synergistic action of three recombination gene products of bacteriophage T4, *uvsX*, *uvsY*, and *gene 32* proteins. *J. Biol. Chem.* 264, 7814-7820, (1989)
52. Game, J.C. and Mortimer, R.K., A genetic study of X-ray sensitive mutants in yeast. *Mutation Research* 24, 281, (1974)
53. Game, J.C., DNA double-strand breaks and the *RAD50-RAD57* genes in *Saccharomyces*. *Semin Cancer Biol* 4, 73-83, (1993)
54. Petes, T.D., Malone, R.E. and Symington, L.S., Recombination in yeast, in *The Molecular and Cellular Biology of the Yeast Saccharomyces: Genome Dynamics, Protein Synthesis, and Energetics*, Vol. I, Broach, J.R., Jones, E. and Pringle, J., Eds., New York, Cold Spring Harbor Laboratory, 1991, 407-521.
55. Resnick, M.A., Investigating the genetic control of biochemical events in meiotic recombination, in *Meiosis*, Moens, P.B., Ed., New York, Academic Press Inc., 1987, 157-210.
56. Game, J.C., Radiation-sensitive mutants and repair in yeast, in *Yeast Genetics, Fundamental and Applied Aspects*, Spencer, J.F.T., Spencer, D. and Smith, A.R.W., Eds., New York, Springer Verlag, 1983, 109-137.
57. Cao, L., Alani, E. and Kleckner, N., A pathway for generation and processing of double-strand breaks during meiotic recombination in *S. cerevisiae*. *Cell* 61, 1089-1101, (1990)
58. Sun, H., Treco, D. and Szostak, J.W., Extensive 3'-overhanging, single-stranded DNA associated with the meiosis-specific double-strand breaks at the *ARG4* recombination initiation site. *Cell* 64, 1155-1161, (1991)
59. Shinohara, A., Ogawa, H. and Ogawa, T., Rad51 protein involved in repair and recombination in *S. cerevisiae* is a RecA-like protein. *Cell* 69, 457-470, (1992)
60. Aboussekhr, A., Chanet, R., Adjiri, A. and Fabre, F., Semidominant suppressors of *Srs2* helicase mutations of *Saccharomyces cerevisiae* map in the *RAD51* gene, whose sequence predicts a protein with similarities to procaryotic RecA proteins. *Mol. Cell. Biol.* 12, 3224-3234, (1992)
61. Basile, G., Aker, M. and Mortimer, R.K., Nucleotide sequence and transcriptional regulation of the yeast recombinational repair gene *RAD51*. *Mol. Cell. Biol.* 12, 3235-3246, (1992)
62. Sung, P., Catalysis of ATP-dependent homologous DNA pairing and strand exchange by yeast Rad51 protein. *Science* 265, 1241-1243, (1994)
63. Sung, P. and Roberson, D.L., DNA strand exchange mediated by a Rad51-ssDNA nucleoprotein filament with polarity opposite to that of RecA. *Cell* 82, 453-461, (1995)

64. Sung, P. and Stratton, S.A., Yeast Rad51 recombinase mediates polar DNA strand exchange in the absence of ATP hydrolysis. *J Biol Chem* 271, 27983-27986, (1996)
65. Sugiyama, T., Zaitseva, E.M. and Kowalczykowski, S.C., A single-stranded DNA-binding protein is needed for efficient presynaptic complex formation by the *Saccharomyces cerevisiae* Rad51 protein. *J. Biol. Chem.* 272, 7940-7945, (1997)
66. Benson, F.E., Stasiak, A. and West, S.C., Purification and characterization of the human Rad51 protein, an analogue of *E. coli* RecA. *Embo J* 13, 5764-5771, (1994)
67. Gupta, R.C., Bazemore, L.R., Golub, E.I. and Radding, C.M., Activities of human recombination protein Rad51. *Proc. Natl. Acad. Sci. USA* 94, 463-468, (1997)
68. Bishop, D.K., Park, D., Xu, L. and Kleckner, N., *DMC1*: a meiosis-specific yeast homolog of *E. coli* RecA required for recombination, synaptonemal complex formation, and cell cycle progression. *Cell* 69, 439-456, (1992)
69. Sung, P., Yeast RAD55 and RAD57 proteins form a heterodimer that functions with replication protein A to promote DNA strand exchange by Rad51 recombinase. *Genes Dev.* 11, 1111-1121, (1997)
70. New, J.H., Sugiyama, T., Zaitseva, E. and Kowalczykowski, S.C., Rad52 protein stimulates DNA strand exchange by Rad51 and replication protein A. *Nature* 391, 407-410, (1998)
71. Sung, P., Function of yeast *RAD52* protein as a mediator between replication protein A and the Rad51 recombinase. *J. Biol. Chem.* 272, 28194-28197, (1997)
72. Resnick, M.A., The repair of double-strand breaks in DNA: A model involving recombination. *J. Theor. Biol.* 59, 97-106, (1976)
73. Szostak, J.W., Orr-Weaver, T.L., Rothstein, R.J. and Stahl, F.W., The double-strand break repair model for recombination. *Cell* 33, 25-35, (1983)
74. Thaler, D.S. and Stahl, F.W., DNA double-chain breaks in recombination of phage lambda and of yeast. *Annu. Rev. Genet.* 22, 169-197, (1988)
75. Clark, A.J., Sandler, S.J., Willis, D.K., Chu, C.C., Blamar, M.A. and Lovett, S.T., Genes of the *RecE* and *RecF* pathways of conjugational recombination in *Escherichia coli*. *Cold Spring Harbor Symp. Quant. Biol.* 49, 453-462, (1984)
76. Rattray, A.J. and Symington, L.S., Multiple pathways for homologous recombination in *Saccharomyces cerevisiae*. *Genetics* 139, 45-56, (1995)
77. Rattray, A.J. and Symington, L.S., Use of a chromosomal inverted repeat to demonstrate that the *RAD51* and *RAD52* genes of *Saccharomyces cerevisiae* have different roles in mitotic recombination. *Genetics* 138, 587-595, (1994)
78. Sugawara, N. and Haber, J.E., Characterization of double-strand break-induced recombination: homology requirements and single-stranded DNA formation. *Mol Cell Biol* 12, 563-575, (1992)
79. Shinohara, A., Ogawa, H., Matsuda, Y., Ushio, N., Ikeo, K. and Ogawa, T., Cloning of human, mouse and fission yeast recombination genes homologous to Rad51 and RecA. *Nat. Genet.* 4, 239-243, (1993)
80. Keeney, S., Giroux, C.N. and Kleckner, N., Meiosis-specific DNA double-strand breaks are catalyzed by *SPO11*, a member of a widely conserved protein family. *Cell* 88, 375-384, (1997)
81. Bergerat, A., de Massy, B., Gadelle, D., Varoutas, P.C., Nicolas, A. and Forterre, P., An atypical topoisomerase II from Archaea with implications for meiotic recombination, *Nature* 386, 414-417, (1997)
82. Kuzminov, A., Collapse and repair of replication forks in *Escherichia coli*. *Mol. Microbiol.* 16, 373-384, (1995)
83. Smith, G.R., Hotspots of homologous recombination. *Experientia* 50, 234-241, (1994)
84. Myers, R.S. and Stahl, F.W., Chi and the RecBCD D enzyme of *Escherichia coli*. *Annu. Rev. Genet.* 28, 49-70, (1994)
85. Dixon, D.A. and Kowalczykowski, S.C., Homologous pairing *in vitro* stimulated by the recombination hotspot, Chi. *Cell* 66, 361-371, (1991)
86. Dixon, D.A. and Kowalczykowski, S.C., The recombination hotspot χ is a regulatory sequence that acts by attenuating the nuclease activity of the *E. coli* RecBCD enzyme. *Cell* 73, 87-96, (1993)
87. Smith, G.R., Kunes, S.M., Schultz, D.W., Taylor, A. and Triman, K.L., Structure of Chi hotspots of generalized recombination. *Cell* 24, 429-436, (1981)
88. Bianco, P.R. and Kowalczykowski, S.C., The recombination hotspot Chi is recognized by the translocating RecBCD enzyme as the single strand of DNA containing the sequence 5'-GCTGGTGG-3'. *Proc. Natl. Acad. Sci. USA* 94, 6706-6711, (1997)
89. Lam, S.T., Stahl, M.M., McMillin, K.D. and Stahl, F.W., Rec-mediated recombinational hot spot activity in bacteriophage lambda. II. A mutation which causes hot spot activity. *Genetics* 77, 425-433, (1974)
90. Dixon, D.A. and Kowalczykowski, S.C., Role of the *Escherichia coli* recombination hotspot, χ , in RecABCD-dependent homologous pairing. *J. Biol. Chem.* 270, 16360-16370, (1995)
91. Anderson, D.G. and Kowalczykowski, S.C., The recombination hot spot χ is a regulatory element that switches the polarity of DNA degradation by the RecBCD enzyme. *Genes Dev.* 11, 571-581, (1997)

92. Ponticelli, A.S., Schultz, D.W., Taylor, A.F. and Smith, G.R., Chi-dependent DNA strand cleavage by recBC enzyme. *Cell* 41, 145-151, (1985)
93. Smith, G.R., Schultz, D.W., Taylor, A.F. and Triman, K., Chi sites, recBC enzyme, and generalized recombination. *Stadler Symp.* 13, 25-37, (1981)
94. Anderson, D.G. and Kowalczykowski, S.C., The translocating RecBCD enzyme stimulates recombination by directing RecA protein onto ssDNA in a χ -regulated manner. *Cell* 90, 77-86, (1997)
95. Tracy, R.B. and Kowalczykowski, S.C., *In vitro* selection of preferred DNA pairing sequences by the *Escherichia coli* RecA protein. *Genes Dev.* 10, 1890-1903, (1996)
96. Harmon, F.G. and Kowalczykowski, S.C., RecQ helicase, in concert with RecA and SSB proteins, initiates and disrupts DNA recombination. *Genes and Development* 12, 1134-1144, (1998)
97. Prashad, N. and Hosoda, J., Role of genes 46 and 47 in bacteriophage T4 reproduction. II. Formation of gaps on parental DNA of polynucleotide ligase defective mutants. *J Mol Biol* 70, 617-635, (1972)
98. Richardson, R.W. and Nossal, N.G., Characterization of the bacteriophage T4 gene 41 DNA helicase. *J. Biol. Chem.* 264, 4725-4731, (1989)
99. Tarumi, K. and Yonesaki, T., Functional interactions of gene 32, 41, and 59 proteins of bacteriophage T4. *J Biol Chem* 270, 2614-2619, (1995)
100. Morrical, S.W., Hempstead, K. and Morrical, M.D., The gene 59 protein of bacteriophage T4 modulates the intrinsic and single-stranded DNA-stimulated ATPase activities of gene 41 protein, the T4 replicative DNA helicase. *J. Biol. Chem.* 269, 33069-33081, (1994)
101. Wu, T.C. and Lichten, M., Meiosis-induced double-strand break sites determined by yeast chromatin structure. *Science* 263, 515-518, (1994)
102. Sun, H., Treco, D., Schultes, N.P. and Szostak, J.W., Double-strand breaks at an initiation site for meiotic gene conversion. *Nature* 338, 87-90, (1989)
103. Ohta, K., Shibata, T. and Nicolas, A., Changes in chromatin structure at recombination initiation sites during yeast meiosis. *EMBO J.* 13, 5754-5763, (1994)
104. Fan, Q., Xu, F. and Petes, T.D., Meiosis-specific double-strand DNA breaks at the *HIS4* recombination hot spot in the yeast *Saccharomyces cerevisiae*: control in cis and trans. *Mol Cell Biol* 15, 1679-1688, (1995)
105. Xu, L. and Kleckner, N., Sequence non-specific double-strand breaks and interhomolog interactions prior to double-strand break formation at a meiotic recombination hot spot in yeast. *Embo J* 14, 5115-5128, (1995)
106. Liu, J., Wu, T.C. and Lichten, M., The location and structure of double-strand DNA breaks induced during yeast meiosis: evidence for a covalently linked DNA-protein intermediate. *Embo J* 14, 4599-4608, (1995)
107. Alani, E., Padmore, R. and Kleckner, N., Analysis of wild-type and *RAD50* mutants of yeast suggests an intimate relationship between meiotic chromosome synapsis and recombination. *Cell* 61, 419-436, (1990)
108. Menees, T.M., Ross-MacDonald, P.B. and Roeder, G.S., *MEI4*, a meiosis-specific yeast gene required for chromosome synapsis. *Mol Cell Biol* 12, 1340-1351, (1992)
109. Rockmill, B., Engebrecht, J.A., Scherthan, H., Loidl, J. and Roeder, G.S., The yeast *MER2* gene is required for chromosome synapsis and the initiation of meiotic recombination. *Genetics* 141, 49-59, (1995)
110. Ajimura, M., Leem, S.H. and Ogawa, H., Identification of new genes required for meiotic recombination in *Saccharomyces cerevisiae*. *Genetics* 133, 51-66, (1993)
111. Johzuka, K. and Ogawa, H., Interaction of *MRE11* and *RAD50*: two proteins required for DNA repair and meiosis-specific double-strand break formation in *Saccharomyces cerevisiae*. *Genetics* 139, 1521-1532, (1995)
112. Bullard, S.A., Kim, S., Galbraith, A.M. and Malone, R.E., Double strand breaks at the *HIS2* recombination hot spot in *Saccharomyces cerevisiae*. *Proc Natl Acad Sci U S A* 93, 13054-13059, (1996)
113. Ivanov, E.L., Korolev, V.G. and Fabre, F., *XRS2*, a DNA repair gene of *Saccharomyces cerevisiae*, is needed for meiotic recombination. *Genetics* 132, 651-664, (1992)
114. Ivanov, E.L., Sugawara, N., White, C.I., Fabre, F. and Haber, J.E., Mutations in *XRS2* and *RAD50* delay but do not prevent mating-type switching in *Saccharomyces cerevisiae*. *Mol Cell Biol* 14, 3414-3425, (1994)
115. Keeney, S. and Kleckner, N., Covalent protein-DNA complexes at the 5' strand termini of meiosis-specific double-strand breaks in yeast. *Proc Natl Acad Sci U S A* 92, 11274-11278, (1995)
116. Ogawa, H., Johzuka, K., Nakagawa, T., Leem, S.H. and Hagihara, A.H., Functions of the yeast meiotic recombination genes, *MRE11* and *MRE2*. *Adv Biophys* 31, 67-76, (1995)
117. Kowalczykowski, S.C. and Krupp, R.A., Effects of the *Escherichia coli* SSB protein on the single-stranded DNA-dependent ATPase activity of *Escherichia coli* RecA protein: Evidence that SSB protein facilitates the binding of RecA protein to regions of secondary structure within single-stranded DNA. *J. Mol. Biol.* 193, 97-113, (1987)

118. Morrical, S.W., Lee, J. and Cox, M.M., Continuous association of *Escherichia coli* single-stranded DNA binding protein with stable complexes of RecA protein and single-stranded DNA. *Biochemistry* 25, 1482-1494, (1986)
119. Muniyappa, K., Shaner, S.L., Tsang, S.S. and Radding, C.M., Mechanism of the concerted action of RecA protein and helix destabilizing proteins in homologous recombination. *Proc. Natl. Acad. Sci. USA* 81, 2757-2761, (1984)
120. Roman, L.J. and Kowalczykowski, S.C., Relationship of the physical and enzymatic properties of *Escherichia coli* RecA protein to its strand exchange activity. *Biochemistry* 25, 7375-7385, (1986)
121. McEntee, K., Weinstock, G.M. and Lehman, I.R., RecA protein-catalyzed strand assimilation: Stimulation by *Escherichia coli* single-stranded DNA-binding protein. *Proc. Natl. Acad. Sci. USA* 77, 857-861, (1980)
122. Cox, M.M., Soltis, D.A., Livneh, Z. and Lehman, I.R., On the role of single-stranded DNA binding protein in RecA protein-promoted DNA strand exchange. *J. Biol. Chem.* 258, 2577-2585, (1983)
123. Umezū, K., Chi, N.W. and Kolodner, R.D., Biochemical interaction of the *Escherichia coli* RecF, RecO, and RecR proteins with RecA protein and single-stranded DNA binding protein. *Proc. Natl. Acad. Sci. USA* 90, 3875-3879, (1993)
124. Umezū, K. and Kolodner, R.D., Protein interactions in genetic recombination in *Escherichia coli*. Interactions involving RecO and RecR overcome the inhibition of RecA by single-stranded DNA-binding protein. *J. Biol. Chem.* 269, 30005-30013, (1994)
125. Konforti, B.B. and Davis, R.W., ATP hydrolysis and the displaced strand are two factors that determine the polarity of RecA-promoted DNA strand exchange. *J. Mol. Biol.* 227, 38-53, (1992)
126. Thresher, R.J., Christiansen, G. and Griffith, J.D., Assembly of presynaptic filaments. Factors affecting the assembly of RecA protein onto single-stranded DNA. *J. Mol. Biol.* 201, 101-113, (1988)
127. Sweezy, M.A. and Morrical, S.W., Single-stranded DNA binding properties of the uvsY recombination protein of bacteriophage T4. *J. Mol. Biol.* 266, 927-938, (1997)
128. Beernink, H.T.H. and Morrical, S.W., The UvsY recombination protein of bacteriophage T4 forms hexamers in the presence and absence of single-stranded DNA. *Biochemistry* in press, (1998)
129. Yassa, D.S., Chou, K.M. and Morrical, S.W., Characterization of an amino-terminal fragment of the bacteriophage T4 uvsY recombination protein. *Biochimie* 79, 275-285, (1997)
130. Bai, Y. and Symington, L.S., A RAD52 homolog is required for RAD51-independent mitotic recombination in *Saccharomyces cerevisiae*. *Genes Dev.* 10, 2025-2037, (1996)
131. Firmenich, A.A., Elias-Arnanz, M. and Berg, P., A novel allele of *Saccharomyces cerevisiae* RFA1 that is deficient in recombination and repair and suppressible by RAD52. *Mol. Cell. Biol.* 15, 1620-1631, (1995)
132. Smith, J. and Rothstein, R., A mutation in the gene encoding the *Saccharomyces cerevisiae* single-stranded DNA-binding protein Rfa1 stimulates a RAD52-independent pathway for direct-repeat recombination. *Mol. Cell. Biol.* 15, 1632-1641, (1995)
133. Shinohara, A. and Ogawa, T., Stimulation by Rad52 of yeast Rad51-mediated recombination. *Nature* 391, 404-407, (1998)
134. Petukhova, G., Stratton, S. and Sung, P., Catalysis of homologous DNA pairing by yeast Rad51 and Rad54 proteins. *Nature* 393, 91-94, (1998)
135. Milne, G.T. and Weaver, D.T., Dominant negative alleles of RAD52 reveal a DNA repair/recombination complex including RAD51 and RAD52. *Genes Dev.* 7, 1755-1765, (1993)
136. Hays, S.L., Firmenich, A.A. and Berg, P., Complex formation in yeast double-strand break repair: participation of Rad51, Rad52, Rad55, and Rad57 proteins. *Proc. Natl. Acad. Sci. USA* 92, 6925-6929, (1995)
137. Jiang, H., Xie, Y., Houston, P., Stemke-Hale, K., Mortensen, U.H., Rothstein, R. and Kodadek, T., Direct association between the yeast Rad51 and Rad54 recombination proteins. *J. Biol. Chem.* 271, 33181-33186, (1996)
138. Clever, B., Interthal, H., Schmuckli-Maurer, J., King, J., Sigrist, M. and Heyer, W.D., Recombinational repair in yeast: functional interactions between Rad51 and Rad54 proteins. *EMBO J.* 16, 2535-2544, (1997)
139. Schwacha, A. and Kleckner, N., Identification of double Holliday junctions as intermediates in meiotic recombination. *Cell* 83, 783-791, (1995)
140. West, S.C., The processing of recombination intermediates: Mechanistic insights from studies of bacterial proteins. *Cell* 76, 9-15, (1994)
141. Tsaneva, I.R. and West, S.C., Targeted versus non-targeted DNA helicase activity of the RuvA and RuvB proteins of *Escherichia coli*. *J. Biol. Chem.* 269, 26552-26558, (1994)
142. Mitchell, A.H. and West, S.C., Role of RuvA in branch migration reactions catalyzed by the RuvA and RuvB proteins of *Escherichia coli*. *J. Biol. Chem.* 271, 19497-19502, (1996)

143. West, S.C., Processing of recombination intermediates by the RuvABC proteins. *Annu Rev Genet* 31, 213-244, (1997)
144. Shiba, T., Iwasaki, H., Nakata, A. and Shinagawa, H., SOS-inducible DNA repair proteins, RuvA and RuvB, of *Escherichia coli*: functional interactions between RuvA and RuvB for ATP hydrolysis and renaturation of the cruciform structure in supercoiled DNA. *Proc. Natl. Acad. Sci. USA* 88, 8445-8449, (1991)
145. Iwasaki, H., Takahagi, M., Nakata, A. and Shinagawa, H., *Escherichia coli* RuvA and RuvB proteins specifically interact with Holliday junctions and promote branch migration. *Genes Dev.* 6, 2214-2220, (1992)
146. Tsaneva, I.R., Muller, B. and West, S.C., ATP-dependent branch migration of Holliday junctions promoted by the RuvA and RuvB proteins of *E. coli*. *Cell* 69, 1171-1180, (1992)
147. West, S.C. and Connolly, B., Biological roles of the *Escherichia coli* RuvA, RuvB and RuvC proteins revealed. *Mol. Microbiol.* 6, 2755-2759, (1992)
148. Lloyd, R.G. and Sharples, G.J., Dissociation of synthetic Holliday junctions by *E. coli* RecG protein. *EMBO J.* 12, 17-22, (1993)
149. Lloyd, R.G. and Sharples, G.J., Processing of recombination intermediates by the RecG and RuvAB proteins of *Escherichia coli*. *Nucleic Acids Res.* 21, 1719-1725, (1993)
150. Whitby, M.C., Ryder, L. and Lloyd, R.G., Reverse branch migration of Holliday junctions by RecG protein: a new mechanism for resolution of intermediates in recombination and DNA repair. *Cell* 75, 341-350, (1993)
151. Kodadek, T. and Alberts, B.M., Stimulation of protein-directed strand exchange by a DNA helicase. *Nature* 326, 312-314, (1987)
152. Jongeneel, C.V., Formosa, T. and Alberts, B.M., Purification and characterization of the bacteriophage T4 dda protein: A DNA helicase that associates with the viral helix-destabilizing protein. *J. Biol. Chem.* 259, 12925-12932, (1984)
153. Kodadek, T., Inhibition of protein-mediated homologous pairing by a DNA helicase. *J. Biol. Chem.* 266, 9712-9718, (1991)
154. Salinas, F. and Kodadek, T., Phage T4 homologous strand exchange: a DNA helicase, not the strand transferase, drives polar branch migration. *Cell* 82, 111-119, (1995)
155. Morrical, S.W., Beermink, H.T.H., Dash, A. and Hempstead, K., The gene 59 protein of bacteriophage T4. Characterization of protein-protein interactions with gene 32 protein, the T4 single-stranded DNA binding protein. *J. Biol. Chem.* 271, 20198-20207, (1996)
156. Carles-Kinch, K., George, J.W. and Kreuzer, K.N., Bacteriophage T4 UvsW protein is a helicase involved in recombination, repair and the regulation of DNA replication origins. *Embo J* 16, 4142-4151, (1997)
157. Connolly, B. and West, S.C., Genetic recombination in *Escherichia coli*: Holliday junctions made by RecA protein are resolved by fractionated cell-free extracts. *Proc. Natl. Acad. Sci. USA* 87, 8476-8480, (1990)
158. Connolly, B., Parsons, C.A., Benson, F.E., Dunderdale, H.J., Sharples, G.J., Lloyd, R.G. and West, S.C., Resolution of Holliday junctions *in vitro* requires the *Escherichia coli* *ruvC* gene product. *Proc. Natl. Acad. Sci. USA* 88, 6063-6067, (1991)
159. Eggleston, A.K., Mitchell, A.H. and West, S.C., *In vitro* reconstitution of the late steps of genetic recombination in *E. coli*. *Cell* 89, 607-617, (1997)
160. Mizuuchi, K., Kemper, B., Hays, J. and Weisberg, R.A., T4 endonuclease VII cleaves holliday structures. *Cell* 29, 357-365, (1982)
161. Mueller, J.E., Newton, C.J., Jensch, F., Kemper, B., Cunningham, R.P., Kallenbach, N.R. and Seeman, N.C., Resolution of Holliday junction analogs by T4 endonuclease VII can be directed by substrate structure. *J Biol Chem* 265, 13918-13924, (1990)
162. Parsons, C.A., Kemper, B. and West, S.C., Interaction of a four-way junction in DNA with T4 endonuclease VII. *J Biol Chem* 265, 9285-9289, (1990)
163. Takahagi, M., Iwasaki, H., Nakata, A. and Shinagawa, H., Molecular analysis of the *Escherichia coli* *ruvC* gene, which encodes a Holliday junction-specific endonuclease. *J. Bacteriol.* 173, 5747-5753, (1991)
164. Jensch, F. and Kemper, B., Endonuclease VII resolves Y-junctions in branched DNA *in vitro*. *Embo J* 5, 181-189, (1986)
165. Kleff, S. and Kemper, B., Initiation of heteroduplex-loop repair by T4-encoded endonuclease VII *in vitro*. *Embo J* 7, 1527-1535, (1988)
166. West, S.C. and Korner, A., Cleavage of cruciform DNA structures by an activity from *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA* 82, 6445-6449, (1985)
167. Symington, L.S. and Kolodner, R., Partial purification of an enzyme from *Saccharomyces cerevisiae* that cleaves Holliday junctions. *Proc. Natl. Acad. Sci. USA* 82, 7247-7251, (1985)
168. Kleff, S., Kemper, B. and Sternglanz, R., Identification and characterization of yeast mutants and the gene for a cruciform cutting endonuclease. *Embo J* 11, 699-704, (1992)
169. Lockshon, D., Zweifel, S.G., Freeman-Cook, L.L., Lorimer, H.E., Brewer, B.J. and Fangman, W.L., A role for

recombination junctions in the segregation of mitochondrial DNA in yeast. *Cell* 81, 947-955, (1995)

170. Kowalczykowski, S.C. and Eggleston, A.K., Homologous pairing and DNA strand-exchange proteins. *Annu. Rev. Biochem.* 63, 991-1043, (1994)

171. Cox, M.M., Soltis, D.A., Livneh, Z. and Lehman, I.R., DNA strand exchange promoted by RecA protein and single-stranded DNA-binding protein of *Escherichia coli*. *Cold Spring Harbor Symp. Quant. Biol.* 47, 803-810, (1983)

172. Radding, C.M., Flory, J., Wu, A., Kahn, R., DasGupta, C., Gonda, D., Bianchi, M. and Tsang, S.S., Three phases in homologous pairing: Polymerization of RecA protein on single-stranded DNA, synapsis, and polar strand exchange. *Cold Spring Harbor Symp. Quant. Biol.* 47, 821-828, (1983)

173. Mazin, A.V. and Kowalczykowski, S.C., The specificity of the secondary DNA binding site of RecA protein defines its role in DNA strand exchange. *Proc. Natl. Acad. Sci. USA* 93, 10673-10678, (1996)

174. Mazin, A. and Kowalczykowski, S.C., The function of the secondary DNA-binding site of RecA protein during DNA strand exchange. *EMBO J.* 17, 1161-1168, (1997)

175. Leahy, M.C. and Radding, C.M., Topography of the interaction of RecA protein with single-stranded deoxyoligonucleotides. *J. Biol. Chem.* 261, 6954-6960, (1986)

176. Amaratunga, M. and Benight, A.S., DNA sequence dependence of ATP hydrolysis by RecA protein. *Biochem. Biophys. Res. Commun.* 157, 127-133, (1988)

177. Brenner, S.L., Mitchell, R.S., Morrical, S.W., Neuendorf, S.K., Schutte, B.C. and Cox, M.M., RecA protein-promoted ATP hydrolysis occurs throughout RecA nucleoprotein filaments. *J. Biol. Chem.* 262, 4011-4016, (1987)

178. Bianco, P.R. and Weinstock, G.M., Interaction of the RecA protein of *Escherichia coli* with single-stranded oligodeoxyribonucleotides. *Nucleic Acids Res.* 24, 4933-4939, (1996)

179. Menetski, J.P., Varghese, A. and Kowalczykowski, S.C., Properties of the high-affinity single-stranded DNA binding state of the *Escherichia coli* RecA protein. *Biochemistry* 27, 1205-1212, (1988)

180. Pugh, B.F. and Cox, M.M., High salt activation of RecA protein ATPase in the absence of DNA. *J. Biol. Chem.* 263, 76-83, (1988)

181. Lee, J.W. and Cox, M.M., Inhibition of RecA protein promoted ATP hydrolysis. 1. ATP γ S and ADP are antagonistic inhibitors. *Biochemistry* 29, 7666-7676, (1990)

182. DiCapua, E., Ruigrok, R.W. and Timmins, P.A., Activation of RecA protein: the salt-induced structural transition. *J. Struct. Biol.* 104, 91-96, (1990)

183. Rehrauer, W.M. and Kowalczykowski, S.C., Alteration of the nucleoside triphosphate (NTP) catalytic domain within *Escherichia coli* RecA protein attenuates NTP hydrolysis but not joint molecule formation. *J. Biol. Chem.* 268, 1292-1297, (1993)

184. Kowalczykowski, S.C. and Krupp, R.A., DNA-strand exchange promoted by RecA protein in the absence of ATP: implications for the mechanism of energy transduction in protein-promoted nucleic acid transactions. *Proc. Natl. Acad. Sci. USA* 92, 3478-3482, (1995)

185. Kowalczykowski, S.C., Mechanistic aspects of the DNA strand exchange activity of *E. coli* RecA protein. *Trends Biochem. Sci.* 12, 141-145, (1987)

186. Menetski, J.P. and Kowalczykowski, S.C., Enhancement of *Escherichia coli* RecA protein enzymatic function by dATP. *Biochemistry* 28, 5871-5881, (1989)

187. Register, J.C., III and Griffith, J., The direction of RecA protein assembly onto single strand DNA is the same as the direction of strand assimilation during strand exchange. *J. Biol. Chem.* 260, 12308-12312, (1985)

188. Di Capua, E., Engel, A., Stasiak, A. and Koller, T., Characterization of complexes between RecA protein and duplex DNA by electron microscopy. *J. Mol. Biol.* 157, 87-103, (1982)

189. Kowalczykowski, S.C., Clow, J.C., Somani, R. and Varghese, A., Effects of the *Escherichia coli* SSB protein on the binding of *Escherichia coli* RecA protein to single-stranded DNA: Demonstration of competitive binding and the lack of a specific protein-protein interaction. *J. Mol. Biol.* 193, 81-95, (1987)

190. Ando, R.A. and Morrical, S.W., Single-stranded DNA binding properties of the UvsX recombinase of bacteriophage T4: Binding parameters and effects of nucleotides. *J. Mol. Biol.* submitted, (1998)

191. Menetski, J.P. and Kowalczykowski, S.C., Interaction of RecA protein with single-stranded DNA. Quantitative aspects of binding affinity modulation by nucleotide cofactors. *J. Mol. Biol.* 181, 281-295, (1985)

192. Harris, L.D. and Griffith, J.D., UvsY protein of bacteriophage T4 is an accessory protein for *in vitro* catalysis of strand exchange. *J. Mol. Biol.* 206, 19-27, (1989)

193. Kodadek, T., Gan, D.C. and Stemke-Hale, K., The phage T4 uvsY recombination protein stabilizes presynaptic filaments. *J. Biol. Chem.* 264, 16451-16457, (1989)

194. Madiraju, M.V.V.S., Templin, A. and Clark, A.J., Properties of a mutant RecA-encoded protein reveal a possible role for *Escherichia coli* recF-encoded protein in genetic recombination. *Proc. Natl. Acad. Sci. USA* 85, 6592-6596, (1988)

195. Madiraju, M.V.V.S. and Clark, A.J., Effect of RecF protein on reactions catalyzed by RecA protein. *Nucleic Acids Res.* 19, 6295-6300, (1991)

196. Madiraju, M.V. and Clark, A.J., Evidence for ATP binding and double-stranded DNA binding by *Escherichia coli* RecF protein. *J. Bacteriol.* 174, 7705-7710, (1992)
197. Webb, B.L., Cox, M.M. and Inman, R.B., An interaction between the *Escherichia coli* RecF and RecR proteins dependent on ATP and double-stranded DNA. *J. Biol. Chem.* 270, 31397-31404, (1995)
198. Shan, Q., Bork, J.M., Webb, B.L., Inman, R.B. and Cox, M.M., RecA protein filaments: end-dependent dissociation from ssDNA and stabilization by RecO and RecR proteins. *J. Mol. Biol.* 265, 519-540, (1997)
199. Namsaraev, E. and Berg, P., Characterization of strand exchange activity of yeast Rad51 protein. *Mol. Cell. Biol.* 17, 5359-5368, (1997)
200. Bryant, F.R., Taylor, A.R. and Lehman, I.R., Interaction of the RecA protein of *Escherichia coli* with single-stranded DNA. *J. Biol. Chem.* 260, 1196-1202, (1985)
201. Julin, D.A., Riddles, P.W. and Lehman, I.R., On the mechanism of pairing of single-stranded and double-stranded DNA molecules by the RecA and single-stranded DNA-binding proteins of *Escherichia coli*. *J. Biol. Chem.* 261, 1025-1030, (1986)
202. Gonda, D.K. and Radding, C.M., The mechanism of the search for homology promoted by RecA protein: Facilitated diffusion within nucleoprotein networks. *J. Biol. Chem.* 261, 13087-13096, (1986)
203. Menetski, J.P. and Kowalczykowski, S.C., Transfer of RecA protein from one polynucleotide to another: Kinetic evidence for a ternary intermediate during the transfer reaction. *J. Biol. Chem.* 262, 2085-2092, (1987)
204. Tsang, S.S., Muniyappa, K., Azhderian, E., Gonda, D.K., Radding, C.M., Flory, J. and Chase, J.W., Intermediates in homologous pairing promoted by RecA protein: Isolation and characterization of active presynaptic complexes. *J. Mol. Biol.* 185, 295-309, (1985)
205. Tsang, S.S., Chow, S.A. and Radding, C.M., Networks of DNA and RecA protein are intermediates in homologous pairing. *Biochemistry* 24, 3226-3232, (1985)
206. Weinstock, G.M., McEntee, K. and Lehman, I.R., Hydrolysis of nucleoside triphosphates catalyzed by the RecA protein of *Escherichia coli*: Characterization of ATP hydrolysis. *J. Biol. Chem.* 256, 8829-8834, (1981)
207. Wittung, P., Ellouze, C., Maraboeuf, F., Takahashi, M. and Norden, B., Thermochemical and kinetic evidence for nucleotide-sequence-dependent RecA-DNA interactions. *Eur. J. Biochem.* 245, 715-719, (1997)
208. Berg, O.G., Winter, R.B. and von Hippel, P.H., How do genome-regulatory proteins locate their DNA target sites? *Trends Biochem. Sci.* 7, 52-55, (1982)
209. Shen, P. and Huang, H.V., Homologous recombination in *Escherichia coli*: dependence on substrate length and homology. *Genetics* 112, 441-457, (1986)
210. King, S.R. and Richardson, J.P., Role of homology and pathway specificity for recombination between plasmids and bacteriophage lambda. *Mol. Gen. Genet.* 204, 141-147, (1986)
211. Rao, B.J. and Radding, C.M., Homologous recognition promoted by RecA protein via non-Watson-Crick bonds between identical DNA strands. *Proc. Natl. Acad. Sci. USA* 90, 6646-6650, (1993)
212. Rao, B.J. and Radding, C.M., Formation of base triplets by non-Watson-Crick bonds mediates homologous recognition in RecA recombination filaments. *Proc. Natl. Acad. Sci. USA* 91, 6161-6165, (1994)
213. Baliga, R., Singleton, J.W. and Dervan, P.B., RecA-oligonucleotide filaments bind in the minor groove of double-stranded DNA. *Proc. Natl. Acad. Sci. USA* 92, 10393-10397, (1995)
214. Podyminogin, M.A., Meyer, R.B. and Gamper, H.B., Sequence-specific covalent modification of DNA by cross-linking oligonucleotides. Catalysis by RecA and implication for the mechanism of synaptic joint formation. *Biochemistry* 34, 13098-13108, (1995)
215. Zhou, X. and Adzuma, K., DNA strand exchange mediated by the *Escherichia coli* RecA protein initiates in the minor groove of double-stranded DNA. *Biochemistry* 36, 4650-4661, (1997)
216. Stasiak, A., Stasiak, A.Z. and Koller, T., Visualization of RecA-DNA complexes involved in consecutive stages of an *in vitro* strand exchange reaction. *Cold Spring Harbor Symp. Quant. Biol.* 49, 561-570, (1984)
217. Adzuma, K., Stable synopsis of homologous DNA molecules mediated by the *Escherichia coli* RecA protein involves local exchange of DNA strands. *Genes Dev.* 6, 1679-1694, (1992)
218. Riddles, P.W. and Lehman, I.R., The formation of paranemic and plectonemic joints between DNA molecules by the RecA and single-stranded DNA-binding proteins of *Escherichia coli*. *J. Biol. Chem.* 260, 165-169, (1985)
219. Riddles, P.W. and Lehman, I.R., The formation of plectonemic joints by the RecA protein of *Escherichia coli*. *J. Biol. Chem.* 260, 170-173, (1985)
220. Lavery, P.E. and Kowalczykowski, S.C., A postsynaptic role for single-stranded DNA-binding protein in RecA protein-promoted DNA strand exchange. *J. Biol. Chem.* 267, 9315-9320, (1992)
221. Lavery, P.E. and Kowalczykowski, S.C., Enhancement of RecA protein-promoted DNA strand exchange activity by volume-occupying agents. *J. Biol. Chem.* 267, 9307-9314, (1992)

DNA strand exchange proteins

222. Lavery, P.E. and Kowalczykowski, S.C., Properties of RecA441 protein-catalyzed DNA strand exchange can be attributed to an enhanced ability to compete with SSB protein. *J. Biol. Chem.* 265, 4004-4010, (1990)
223. Harris, L.D. and Griffith, J., Visualization of the homologous pairing of DNA catalyzed by the bacteriophage T4 UvsX protein. *J. Biol. Chem.* 262, 9285-9292, (1987)
224. Kowalczykowski, S.C., Biochemistry of genetic recombination: Energetics and mechanism of DNA strand exchange. *Annu. Rev. Biophys. Biophys. Chem.* 20, 539-575, (1991)
225. Cox, M.M., Morrical, S.W. and Neuendorf, S.K., Unidirectional branch migration promoted by nucleoprotein filaments of RecA protein and DNA. *Cold Spring Harbor Symp. Quant. Biol.* 49, 525-533, (1984)
226. Menetski, J.P., Bear, D.G. and Kowalczykowski, S.C., Stable DNA heteroduplex formation catalyzed by the *Escherichia coli* RecA protein in the absence of ATP hydrolysis. *Proc. Natl. Acad. Sci. USA* 87, 21-25, (1990)
227. Honigberg, S.M. and Radding, C.M., The mechanics of winding and unwinding helices in recombination: torsional stress associated with strand transfer promoted by RecA protein. *Cell* 54, 525-532, (1988)
228. Jwang, B. and Radding, C.M., Torsional stress generated by RecA protein during DNA strand exchange separates strands of a heterologous insert. *Proc. Natl. Acad. Sci. USA* 89, 7596-7600, (1992)
229. Bianchi, M.E. and Radding, C.M., Heteroduplex DNA with large unpaired regions made by RecA protein, in *Mechanisms of DNA Replication and Recombination*, Cozzarelli, N.R., Ed., New York, Alan R. Liss, 1983, 697-707.
230. Bianchi, M.E. and Radding, C.M., Insertions, deletions and mismatches in heteroduplex DNA made by RecA protein. *Cell* 35, 511-520, (1983)
231. Hahn, T.R., West, S. and Howard-Flanders, P., RecA-mediated strand exchange reactions between duplex DNA molecules containing damaged bases, deletions, and insertions. *J. Biol. Chem.* 263, 7431-7436, (1988)
232. Livneh, Z. and Lehman, I.R., Recombinational bypass of pyrimidine dimers promoted by the RecA protein of *Escherichia coli*. *Proc Natl Acad Sci USA* 79, 3171-3175, (1982)
233. Muller, B., Tsaneva, I.R. and West, S.C., Branch migration of Holliday junctions promoted by the *Escherichia coli* RuvA and RuvB proteins. I. Comparison of RuvAB- and RuvB-mediated reactions. *J. Biol. Chem.* 268, 17179-17184, (1993)
234. Morel, P., Stasiak, A., Ehrlich, S.D. and Cassuto, E., Effect of length and location of heterologous sequences on RecA-mediated strand exchange. *J. Biol. Chem.* 269, 19830-19835, (1994)
235. MacFarland, K.J., Shan, Q., Inman, R.B. and Cox, M.M., RecA as a motor protein. Testing models for the role of ATP hydrolysis in DNA strand exchange. *J. Biol. Chem.* 272, 17675-17685, (1997)
236. Stasiak, A., DiCapua, E. and Koller, T., Elongation of duplex DNA by RecA protein. *J. Mol. Biol.* 151, 557-564, (1981)
237. Stasiak, A. and Di Capua, E., The helicity of DNA in complexes with RecA protein. *Nature* 299, 185-186, (1982)
238. Stasiak, A., DiCapua, E. and Koller, T., Unwinding of duplex DNA in complexes with RecA protein. *Cold Spring Harbor Symp. Quant. Biol.* 47, 811-820, (1983)
239. Conley, E.C. and West, S.C., Homologous pairing and the formation of nascent synaptic intermediates between regions of duplex DNA by RecA protein. *Cell* 56, 987-995, (1989)
240. Chiu, S.K., Wong, B.C. and Chow, S.A., Homologous pairing in duplex DNA regions and the formation of four-stranded paranemic joints promoted by RecA protein. Effects of gap length and negative superhelicity. *J. Biol. Chem.* 265, 21262-21268, (1990)
241. Conley, E.C. and West, S.C., Underwinding of DNA associated with duplex-duplex pairing by RecA protein. *J. Biol. Chem.* 265, 10156-10163, (1990)
242. Lindsley, J.E. and Cox, M.M., On RecA protein-mediated homologous alignment of two DNA molecules. Three strands versus four strands. *J. Biol. Chem.* 265, 10164-10171, (1990)
243. Weinstock, G.M., McEntee, K. and Lehman, I.R., Hydrolysis of nucleoside triphosphates catalyzed by the RecA protein of *Escherichia coli*: Steady state kinetic analysis of ATP hydrolysis. *J. Biol. Chem.* 256, 8845-8849, (1981)
244. Weinstock, G.M., McEntee, K. and Lehman, I.R., Hydrolysis of nucleoside triphosphates catalyzed by the RecA protein of *Escherichia coli*: Hydrolysis of UTP. *J. Biol. Chem.* 256, 8856-8858, (1981)
245. Weinstock, G.M., McEntee, K. and Lehman, I.R., Interaction of the RecA protein of *Escherichia coli* with adenosine 5'-O-(3- thiotriphosphate) *J. Biol. Chem.* 256, 8850-8855, (1981)
246. Pugh, B.F. and Cox, M.M., Stable binding of RecA protein to duplex DNA. Unraveling a paradox. *J. Biol. Chem.* 262, 1326-1336, (1987)
247. Lindsley, J.E. and Cox, M.M., Dissociation pathway for RecA nucleoprotein filaments formed on linear duplex DNA. *J. Mol. Biol.* 205, 695-711, (1989)

248. Schutte, B.C. and Cox, M.M., Homology-dependent changes in adenosine 5'-triphosphate hydrolysis during RecA protein promoted DNA strand exchange: evidence for long paranemic complexes. *Biochemistry* 26, 5616-5625, (1987)
249. Cox, M.M., Soltis, D.A., Lehman, I.R., DeBrosse, C. and Benkovic, S.J., ADP-mediated dissociation of stable complexes of RecA protein and single-stranded DNA. *J. Biol. Chem.* 258, 2586-2592, (1983)
250. Moreau, P.L. and Carlier, M.F., RecA protein-promoted cleavage of LexA repressor in the presence of ADP and structural analogues of inorganic phosphate, the fluoride complexes of aluminum and beryllium. *J. Biol. Chem.* 264, 2302-2306, (1989)
251. Honigberg, S.M., Gonda, D.K., Flory, J. and Radding, C.M., The pairing activity of stable nucleoprotein filaments made from RecA protein, single-stranded DNA, and adenosine 5'-(γ -thio)triphosphate. *J. Biol. Chem.* 260, 11845-11851, (1985)
252. Rosselli, W. and Stasiak, A., Energetics of RecA-mediated recombination reactions: Without ATP hydrolysis RecA can mediate polar strand exchange but is unable to recycle. *J. Mol. Biol.* 216, 335-352, (1990)
253. Rosselli, W. and Stasiak, A., The ATPase activity of RecA is needed to push the DNA strand exchange through heterologous regions. *EMBO J.* 10, 4391-4396, (1991)
254. Kim, J.I., Cox, M.M. and Inman, R.B., On the role of ATP hydrolysis in RecA protein-mediated DNA strand exchange. I. Bypassing a short heterologous insert in one DNA substrate. *J. Biol. Chem.* 267, 16438-16443, (1992)
255. Yu, X. and Egelman, E.H., Image analysis reveals that *Escherichia coli* RecA protein consists of two domains. *Biophys. J.* 57, 555-566, (1990)
256. Stasiak, A.Z., Rosselli, W. and Stasiak, A., RecA-DNA helical filaments in genetic recombination. *Biochimie* 73, 199-208, (1991)
257. Shan, Q., Cox, M.M. and Inman, R.B., DNA strand exchange promoted by RecA K72R. Two reaction phases with different Mg^{2+} requirements. *J. Biol. Chem.* 271, 5712-5724, (1996)
258. Kim, J.I., Cox, M.M. and Inman, R.B., On the role of ATP hydrolysis in RecA protein-mediated DNA strand exchange. II. Four-strand exchanges. *J. Biol. Chem.* 267, 16444-16449, (1992)
259. Yonesaki, T., Recombination apparatus of T4 phage. *Adv Biophys* 31, 3-22, (1995)
260. West, S.C., Cassuto, E., Mursalim, J. and Howard-Flanders, P., Recognition of duplex DNA containing single-stranded regions by RecA protein. *Proc. Natl. Acad. Sci. USA* 77, 2569-2573, (1980)
261. McEntee, K., Weinstock, G.M. and Lehman, I.R., Binding of the RecA protein of *Escherichia coli* to single- and double-stranded DNA. *J. Biol. Chem.* 256, 8835-8844, (1981)
262. Dombroski, D.F., Scraba, D.G., Bradley, R.D. and Morgan, A.R., Studies of the interaction of RecA protein with DNA. *Nucleic Acids Res.* 11, 7487-7504, (1983)
263. Ogawa, T., Yu, X., Shinohara, A. and Egelman, E.H., Similarity of the yeast Rad51 filament to the bacterial RecA filament. *Science* 259, 1896-1899, (1993)
264. Rusche, J.R., Konigsberg, W. and Howard-Flanders, P., Isolation of altered RecA polypeptides and interaction with ATP and DNA. *J. Biol. Chem.* 260, 949-955, (1985)
265. Benedict, R.C. and Kowalczykowski, S.C., Increase of the DNA strand assimilation activity of RecA protein by removal of the C terminus and structure-function studies of the resulting protein fragment. *J. Biol. Chem.* 263, 15513-15520, (1988)
266. Tateishi, S., Horii, T., Ogawa, T. and Ogawa, H., C-terminal truncated *Escherichia coli* RecA protein RecA5327 has enhanced binding affinities to single- and double-stranded DNAs. *J. Mol. Biol.* 223, 115-129, (1992)
267. Story, R.M., Weber, I.T. and Steitz, T.A., The structure of the *E. coli* RecA protein monomer and polymer. *Nature* 355, 318-325, (1992)
268. Kurumizaka, H., Aihara, H., Ikawa, S., Kashima, T., Bazemore, L.R., Kawasaki, K., Sarai, A., Radding, C.M. and Shibata, T., A possible role of the C-terminal domain of the RecA protein. A gateway model for double-stranded DNA binding. *J. Biol. Chem.* 271, 33515-33524, (1996)
269. Dunn, K., Chrysogelos, S. and Griffith, J., Electron microscopic visualization of RecA-DNA filaments: Evidence for a cyclic extension of duplex DNA. *Cell* 28, 757-765, (1982)
270. Egelman, E.H. and Yu, X., The location of DNA in RecA-DNA helical filaments. *Science* 245, 404-407, (1989)
271. DiCapua, E., Schnarr, M. and Timmins, P.A., The location of DNA in complexes of RecA protein with double-stranded DNA. A neutron scattering study. *Biochemistry* 28, 3287-3292, (1989)
272. Yu, X. and Egelman, E.H., Structural data suggest that the active and inactive forms of the RecA filament are not simply interconvertible. *J. Mol. Biol.* 227, 334-346, (1992)
273. Story, R.M., Bishop, D.K., Kleckner, N. and Steitz, T.A., Structural relationship of bacterial RecA proteins to recombination proteins from bacteriophage T4 and yeast. *Science* 259, 1892-1896, (1993)
274. Brendel, V., Brocchieri, L., Sandler, S.J., Clark, A.J. and Karlin, S., Evolutionary comparisons of RecA-like proteins

DNA strand exchange proteins

- across all major kingdoms of living organisms. *J. Mol. Evol.* 44, 528-541, (1997)
275. Roca, A.I. and Cox, M.M., RecA protein: structure, function, and role in recombinational DNA repair. *Prog. Nucleic Acid Res. Mol. Biol.* 56, 129-223, (1997)
276. Horii, T., Ogawa, T. and Ogawa, H., Organization of the RecA gene of *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 77, 313-317, (1980)
277. Story, R.M. and Steitz, T.A., Structure of the RecA protein-ADP complex. *Nature* 355, 374-376, (1992)
278. Mikawa, T., Masui, R., Ogawa, T., Ogawa, H. and Kuramitsu, S., N-terminal 33 amino acid residues of *Escherichia coli* RecA protein contribute to its self-assembly. *J. Mol. Biol.* 250, 471-483, (1995)
279. Walker, J.E., Saraste, M., Runswick, M.J. and Gay, N.J., Distantly related sequences in the α - and β -subunits of ATP synthase, myosin, kinases and other ATP-requiring enzymes and a common nucleotide binding fold. *EMBO J.* 1, 945-951, (1982)
280. Knight, K.L. and McEntee, K., Tyrosin 264 in the RecA protein from *Escherichia coli* is the site of modification by the photoaffinity label 8-azido adenosine 5'-triphosphate. *J. Biol. Chem.* 260, 10185-10191, (1985)
281. Gardner, R.V., Voloshin, O.N. and Camerini-Otero, R.D., The identification of the single-stranded DNA-binding domain of the *Escherichia coli* RecA protein. *Eur. J. Biochem.* 233, 419-425, (1995)
282. Nastri, H.G. and Knight, K.L., Identification of residues in the L1 region of the RecA protein which are important to recombination or coprotease activities. *J. Biol. Chem.* 269, 26311-26322, (1994)
283. Takahashi, M. and Norden, B., Accessibility to modification of histidine residues of RecA protein upon DNA and cofactor binding. *Eur. J. Biochem.* 217, 665-670, (1993)
284. Voloshin, O.N., Wang, L. and Camerini-Otero, R.D., Homologous DNA pairing promoted by a 20-amino acid peptide derived from RecA. *Science* 272, 868-872, (1996)
285. Cazaux, C., Larminat, F., Villani, G., Johnson, N.P., Schnarr, M. and Defais, M., Purification and biochemical characterization of *Escherichia coli* RecA proteins mutated in the putative DNA binding site. *J. Biol. Chem.* 269, 8246-8254, (1994)
286. Wang, Y. and Adzuma, K., Differential proximity probing of two DNA binding sites in the *Escherichia coli* RecA protein using photo-cross-linking methods. *Biochemistry* 35, 3563-3571, (1996)
287. Malkov, V.A. and Camerini-Otero, R.D., Photocross-links between single-stranded DNA and *Escherichia coli* RecA protein map to loops L1 (amino acid residues 157-164) and L2 (amino acid residues 195-209) *J. Biol. Chem.* 270, 30230-30233, (1995)
288. Morimatsu, K. and Horii, T., DNA-binding surface of RecA protein photochemical cross-linking of the first DNA binding site on RecA filament. *Eur. J. Biochem.* 234, 695-705, (1995)
289. Rehrauer, W.M. and Kowalczykowski, S.C., The DNA binding site(s) of the *Escherichia coli* RecA protein. *J. Biol. Chem.* 271, 11996-12002, (1996)
290. Morimatsu, K. and Horii, T., The DNA-binding site of the RecA protein. Photochemical cross-linking of Tyr103 to single-stranded DNA. *Eur. J. Biochem.* 228, 772-778, (1995)
291. Harmon, F.G., Rehrauer, W.M. and Kowalczykowski, S.C., Interaction of *Escherichia coli* RecA protein with LexA repressor. II. Inhibition of DNA strand exchange by the uncleavable LexA S119A repressor argues that recombination and SOS induction are competitive processes. *J. Biol. Chem.* 271, 23874-23883, (1996)
292. Yu, X. and Egelman, E.H., The LexA repressor binds within the deep helical groove of the activated RecA filament. *J. Mol. Biol.* 231, 29-40, (1993)
293. DiCapua, E., Cuillel, M., Hewat, E., Schnarr, M., Timmins, P.A. and Ruigrok, R.W., Activation of RecA protein: The open helix model for LexA cleavage. *J. Mol. Biol.* 226, 707-719, (1992)
294. Rehrauer, W.M., Lavery, P.E., Palmer, E.L., Singh, R.N. and Kowalczykowski, S.C., Interaction of *Escherichia coli* RecA protein with LexA repressor. I. LexA repressor cleavage is competitive with binding of a secondary DNA molecule. *J. Biol. Chem.* 271, 23865-23873, (1996)
295. Fujisawa, H., Yonesaki, T. and Minagawa, T., Sequence of the T4 recombination gene, *UvsX*, and its comparison with that of the RecA gene of *Escherichia coli*. *Nucleic Acids Res.* 13, 7473-7481, (1985)
296. Peitsch, M.C. and Jongeneel, C.V., A 3-D model for the CD40 ligand predicts that it is a compact trimer similar to the tumor necrosis factors. *Int. Immunol.* 5, 233-238, (1993)
297. Peitsch, M.C., Protein modeling by E-mail. *BioTechnology* 13, 658-660, (1995)
298. Peitsch, M.C., ProMod and Swiss-Model: Internet-based tools for automated comparative protein modeling. *Biochem Soc Trans* 24, 274-279, (1996)
299. Morita, T., Yoshimura, Y., Yamamoto, A., Murata, K., Mori, M., Yamamoto, H. and Matsushiro, A., A mouse homolog of the *Escherichia coli* RecA and *Saccharomyces cerevisiae* RAD51 genes. *Proc. Natl. Acad. Sci. USA* 90, 6577-6580, (1993)
300. Muris, D.F., Vreeken, K., Carr, A.M., Broughton, B.C., Lehmann, A.R., Lohman, P.H. and Pastink, A., Cloning the

DNA strand exchange proteins

RAD51 homologue of *Schizosaccharomyces pombe*. *Nucleic Acids Res.* 21, 4586-4591, (1993)

301. Bezzubova, O., Shinohara, A., Mueller, R.G., Ogawa, H. and Buerstedde, J.M., A chicken *RAD51* homologue is expressed at high levels in lymphoid and reproductive organs. *Nucleic Acids Res.* 21, 1577-1580, (1993)

302. Maeshima, K., Morimatsu, K., Shinohara, A. and Horii, T., *RAD51* homologues in *Xenopus laevis*: two distinct genes are highly expressed in ovary and testis. *Gene* 160, 195-200, (1995)

303. Donovan, J.W., Milne, G.T. and Weaver, D.T., Homotypic and heterotypic protein associations control *RAD51* function in double-strand break repair. *Genes Dev.* 8, 2552-2562, (1994)

304. Ogawa, T., Shinohara, A. and Ikeya, T., A species-specific interaction of Rad51 and Rad52 proteins in eukaryotes. *Adv. Biophys.* 31, 93-100, (1995)

305. Shen, Z., Cloud, K.G., Chen, D.J. and Park, M.S., Specific interactions between the human Rad51 and Rad52 proteins. *J. Biol. Chem.* 271, 148-152, (1996)

306. Kowalczykowski, S.C., *In vitro* reconstitution of homologous recombination reactions. *Experientia* 50, 204-215, (1994)

307. Gilbertson, L.A. and Stahl, F.W., A test of the double-strand break repair model for meiotic recombination in *Saccharomyces cerevisiae*. *Genetics* 144, 27-41, 144, 27-41, (1996)

Key Words: DNA binding protein, DNA strand exchange, DNA repair, homologous recombination, nuclease, helicase, RecA protein, Rad51 protein, UvsX protein, RecBCD enzyme, SSB protein, RecF, RecO, RecR, RecG, RecQ, RuvAB, RuvC, DNA gyrase, DNA topoisomerase, gp46, gp47, gp41, gp59, gp32, UvsY, Dda, UvsW, gp41, gp59, gp49, Mre11, Rad50, Xrs2, Spo11, RPA, Rad52, Rad54, Rad55, Rad57, Rad59

Send correspondence to: Dr Stephen C. Kowalczykowski, Section of Microbiology, University of California, 1 Shields Ave., 156 Hutchison Hall, Davis, CA 95616, Tel: (530)-752-5938, Fax:(530)-752-5939, E-mail: sckowalczykowski@ucdavis.edu